

## Dual Effect of H<sub>2</sub>O<sub>2</sub> on the Regulation of Cholecystokinin-induced Amylase Release in Rat Pancreatic Acinar Cells

Jeong Mi An, Jin Hak Rhie, and Jeong Taeg Seo\*

Department of Oral Biology, Oral Science Research Center, Yonsei University College of Dentistry, 134 Sinchon-dong, Seodaemoon-gu, Seoul 120-752, Korea

(Received August 17, 2006 ; Accepted September 28, 2006)

H<sub>2</sub>O<sub>2</sub>, a member of reactive oxygen species (ROS), is known to be involved in the mediation of physiological functions in a variety of cell types. However, little has been known about the physiological role of H<sub>2</sub>O<sub>2</sub> in exocrine cells. Therefore, in the present study, the effect of H<sub>2</sub>O<sub>2</sub> on cholecystokinin (CCK)-evoked Ca<sup>2+</sup> mobilization and amylase release was investigated in rat pancreatic acinar cells. Stimulation of the acinar cells with sulfated octapeptide form of CCK (CCK-8S) induced biphasic increase in amylase release. Addition of 30 μM H<sub>2</sub>O<sub>2</sub> enhanced amylase release caused by 10 pM CCK-8S, but inhibited the amylase release induced by CCK-8S at concentrations higher than 100 pM. An ROS scavenger, 10 μM Mn(III)tetrakis(4-benzoic acid)porphyrin chloride, increased amylase release caused by CCK-8S at concentrations higher than 100 pM, although lower concentrations of CCK-8S-induced amylase release was not affected. To examine whether the effect of H<sub>2</sub>O<sub>2</sub> on CCK-8S-induced amylase release was exerted via modulation of intracellular Ca<sup>2+</sup> signaling, we measured the changes in intracellular Ca<sup>2+</sup> concentration ([Ca<sup>2+</sup>]<sub>i</sub>) in fura-2 loaded acinar cells. Although 30 μM H<sub>2</sub>O<sub>2</sub> did not induce any increase in [Ca<sup>2+</sup>]<sub>i</sub> by itself, it increased the frequency and amplitude of [Ca<sup>2+</sup>]<sub>i</sub> oscillations caused by 10 pM CCK-8S. However, 30 μM H<sub>2</sub>O<sub>2</sub> had little effect on 1 nM CCK-8S-induced increase in [Ca<sup>2+</sup>]<sub>i</sub>. ROS scavenger, 1 mM N-acetylcysteine, did not affect [Ca<sup>2+</sup>]<sub>i</sub> changes induced by 10 pM or 1 nM CCK-8S. Therefore, it was concluded that 30 μM H<sub>2</sub>O<sub>2</sub> enhanced low concentration of CCK-8S-induced amylase

release probably by increasing [Ca<sup>2+</sup>]<sub>i</sub> oscillations while it inhibited high concentration of CCK-8S-induced amylase release.

**Keywords:** H<sub>2</sub>O<sub>2</sub>, cholecystokinin, amylase release, pancreatic acinar cells

### Introduction

Reactive oxygen species (ROS), such as superoxide anion, H<sub>2</sub>O<sub>2</sub> and the hydroxyl radical, have been considered as cytotoxic by-products of cellular metabolism. However, recent studies have provided evidences that H<sub>2</sub>O<sub>2</sub> serves as a signaling molecule modulating various physiological functions (Dröge, 2002). Cells possess mechanisms that can rapidly synthesize and destroy H<sub>2</sub>O<sub>2</sub> in response to receptor stimulations. For example, stimulation of membrane receptors of various growth factors, such as transforming growth factor-β1, platelet-derived growth factor and epidermal growth factor, triggers the rapid and transient production of H<sub>2</sub>O<sub>2</sub> (Ohba *et al.*, 1994; Sundaresan *et al.*, 1995; Bae *et al.*, 1997; Bae *et al.*, 2000). H<sub>2</sub>O<sub>2</sub> generated in response to the receptor stimulation has been shown to play an important role in regulating various normal cell functions, such as cell proliferation, platelet aggregation and vasodilation (Sobey *et al.*, 1997; Pignatelli *et al.*, 1998; Varela *et al.*, 2004). In addition to this, exogenous addition of H<sub>2</sub>O<sub>2</sub> at low concentrations affects the functions of various proteins involved in signal transduction and ion channels (Tournier *et al.*, 1997; Akaishi *et al.*, 2004; Varela *et al.*, 2004).

The exocrine pancreas plays an essential role in the synthesis and secretion of major enzymes for the digestion.

\*Corresponding author: Jeong Taeg Seo,

Tel.: +82-2-2228-3054; Fax.: +82-2-364-1085; e-mail: jeong@yumc.yonsei.ac.kr

Control of pancreatic secretion is regulated primarily by cholecystokinin (CCK). The signal transduction pathways linked to CCK receptor activation have been extensively studied in pancreatic acinar cells (Williams, 2001). Stimulation of the CCK receptors, which are coupled to Gq/11-proteins, increases the production of inositol 1,4,5-trisphosphate (IP<sub>3</sub>) through the activation of phospholipase C. IP<sub>3</sub> increases intracellular Ca<sup>2+</sup> concentration ([Ca<sup>2+</sup>]<sub>i</sub>) both by mobilization of Ca<sup>2+</sup> from intracellular Ca<sup>2+</sup> stores and by stimulation of Ca<sup>2+</sup> entry across the plasma membrane. Recently, the existence of other Ca<sup>2+</sup> mobilizing second messengers, such as cyclic ADP ribose and nicotinic acid adenine dinucleotide phosphate (NAADP), was also reported to be involved in the CCK-induced Ca<sup>2+</sup> mobilization (Cancela *et al.*, 2000). An increase in [Ca<sup>2+</sup>]<sub>i</sub> triggers exocytosis of secretory granules containing digestive enzymes (Williams, 2001).

Recently, a great deal of attention has focused on the sensitivity of the mechanisms responsible for Ca<sup>2+</sup> mobilization to changes in the redox state. H<sub>2</sub>O<sub>2</sub> has been shown to enhance the mobilization of Ca<sup>2+</sup> in many cell types by modifying Ca<sup>2+</sup> channels, such as L-type Ca<sup>2+</sup> channels (Akaishi *et al.*, 2004), long transient receptor potential channel 2 (LTRPC2, Wehage *et al.*, 2002), ryanodine-dependent Ca<sup>2+</sup> channels (Favero *et al.*, 1995) and IP<sub>3</sub>-dependent Ca<sup>2+</sup> channels (Hu *et al.*, 2000), Ca<sup>2+</sup> pumps such as sarcoendoplasmic reticulum Ca<sup>2+</sup> ATPase (SERCA, Redondo *et al.*, 2004) and plasma membrane Ca<sup>2+</sup> ATPase (PMCA, Redondo *et al.*, 2004) or enzymes involved in Ca<sup>2+</sup> signaling pathways, such as phospholipase Cγ1 (PLCγ1, Wang *et al.*, 2001) and phospholipase D (PLD, Servitja *et al.*, 2000).

Given the widespread involvement of H<sub>2</sub>O<sub>2</sub> in modulating Ca<sup>2+</sup> signaling cascade, it is tempting to speculate that amylase release from pancreatic acinar cells, which is regulated mainly by Ca<sup>2+</sup> signaling, may be modified by H<sub>2</sub>O<sub>2</sub>.

Therefore, in the present study, we investigated the roles of H<sub>2</sub>O<sub>2</sub> in Ca<sup>2+</sup> signaling and amylase release in rat pancreatic acinar cells. Our results indicate that low, physiologically relevant concentration of H<sub>2</sub>O<sub>2</sub> (30 μM) enhances low concentration of CCK-induced [Ca<sup>2+</sup>]<sub>i</sub> oscillations and amylase release, while it inhibits high concentration of CCK-induced amylase release.

---

## Materials and Methods

### Preparation of Rat Pancreatic Acinar Cells

Male Sprague-Dawley rats (150-250 g) were anesthetized with diethyl ether and killed by decapitation. The pancreata were immediately removed and trimmed of fat on ice. The acini were prepared using 50 U/ml collagenase by a slight modification of the methods described previously (Matozaki *et al.*, 1990; Toescu *et al.*, 1993). The isolated pancreatic acinar cells were suspended in a HEPES-buffered physiolo-

gical solution containing (in mM): NaCl 140, KCl 5, MgCl<sub>2</sub> 1, CaCl<sub>2</sub> 1, HEPES free acid 10, D-glucose 11.1, bovine serum albumin (0.1 mg/ml), soybean trypsin inhibitor (0.15 mg/ml) and pyruvic acid (1 mg/ml) (adjusted to pH 7.4, gassed with 100% O<sub>2</sub>).

### Amylase Assay

Isolated pancreatic acini were centrifuged and then resuspended in fresh HEPES-buffered physiological solution containing 1.5 mM CaCl<sub>2</sub>. One ml aliquots of acinar suspension distributed into 10 ml polycarbonate Erlenmeyer flasks were incubated with secretagogues at 37°C for 30 min under constant shaking at 60 times/min, and the incubation was then terminated with centrifugation at 500 g for 3 min at 4°C. The supernatant was reserved on ice for a subsequent amylase assay, which was carried out by a modified method of Bernfeld using starch as a substrate (Bernfeld, 1955). The amylase release was expressed as the percentage of the total content of amylase in each sample.

### Fura-2 Loading and [Ca<sup>2+</sup>]<sub>i</sub> Measurements

The cells were loaded with fura-2 by incubation with 2 μM acetoxymethyl ester of fura-2 (fura-2/AM) in a HEPES buffered solution equilibrated with 100% O<sub>2</sub> for 40 min at room temperature. They were washed twice and resuspended in a HEPES buffered solution. The cells were allowed to attach to a coverslip that formed the base of a cell chamber mounted on the stage of an inverted microscope and were superfused with the HEPES buffered solution at a flow rate of 2 ml/min. [Ca<sup>2+</sup>]<sub>i</sub> was measured by spectrofluorometry (Photon Technology International, Brunswick, NJ) with excitations at 340 and 380 nm and emission was measured at 510 nm. The values of [Ca<sup>2+</sup>]<sub>i</sub> were calculated from the ratio of fluorescence intensities (F<sub>340/380</sub>) according to Grynkiewicz *et al.* (1985).

### Materials


H<sub>2</sub>O<sub>2</sub>, N-acetylcysteine (NAC), Collagenase (type IV), soybean trypsin inhibitor, HEPES, bovine serum albumin, MEM amino acids, glutamine and sulfated octapeptide form of CCK (CCK-8S) were purchased from Sigma (St. Louis, MO). Mn(III)tetrakis(4-benzoic acid)porphyrin chloride (MnTBAP) was from Calbiochem (San Diego, CA). Fura-2/AM was obtained from Molecular Probes (Eugene, OR).

---

## Results

### Effect of H<sub>2</sub>O<sub>2</sub> on CCK-induced Amylase Release

CCK has been known to produce a typical biphasic dose-response curve for amylase release (Stark *et al.*, 1989). In this study, we also observed that the amylase release increased as CCK-8S concentration increased, reaching a maximum at 100 pM, and then decreased (Fig. 1A). To examine the effect of H<sub>2</sub>O<sub>2</sub> on amylase release, we treated



**Fig. 1.** Dual Effect of H<sub>2</sub>O<sub>2</sub> on CCK-induced amylase release in rat pancreatic acinar cells. Cells were exposed to various concentrations of CCK-8S with/without 30 μM H<sub>2</sub>O<sub>2</sub> or 10 μM MnTBAP at 37°C for 30 min and the amylase assay was carried out as described in *Materials and Methods*. The amylase release was expressed as the percentage of the total content of amylase in each sample. A. CCK-8S induced a biphasic increase in amylase release. B. Addition of 30 μM H<sub>2</sub>O<sub>2</sub> increased 10 pM CCK-8S-induced amylase release, but decreased the amylase release caused by CCK-8S at concentrations higher than 100 pM. C. ROS scavenger 10 μM MnTBAP increased the amylase release caused by CCK-8S at concentrations higher than 100 pM. Values are means ± S.E.M. from 6-9 separate experiments.

cells with 30 μM H<sub>2</sub>O<sub>2</sub> in combination with each concentration of CCK-8S. As shown in Fig. 1B, addition of 30 μM H<sub>2</sub>O<sub>2</sub> increased 10 pM CCK-8S-induced amylase release, but decreased the amylase release caused by CCK-8S at concentrations higher than 100 pM.

#### Effect of ROS Scavenger on CCK-induced Amylase Release

Because stimulation of the pancreatic acinar cells with CCK has been known to produce ROS (Granados *et al.*, 2004), we examined whether scavenging ROS affected the CCK-induced amylase release. Since some scavengers, such as NAC and glutathione, interfered the measurement of optical density (data not shown), we used MnTBAP to measure the amylase release. As shown in Fig. 1C, 10 μM MnTBAP increased amylase release induced by 100 pM or higher concentrations of CCK-8S, suggesting that ROS produced by CCK-8S stimulation may be involved in the inhibition of high concentrations of CCK-induced amylase release in rat pancreatic acinar cells. The amylase release induced by CCK-8S at concentrations lower than 100 pM was not affected by 10 μM MnTBAP.

#### Effect of H<sub>2</sub>O<sub>2</sub> on [Ca<sup>2+</sup>]<sub>i</sub> Increase in Rat Pancreatic Acinar Cells


The effect of H<sub>2</sub>O<sub>2</sub> on Ca<sup>2+</sup> mobilization was examined in fura-2 loaded rat pancreatic acinar cells. As shown in Fig. 2A and B, exposure of the cells to H<sub>2</sub>O<sub>2</sub> at concentrations lower than 100 μM failed to increase [Ca<sup>2+</sup>]<sub>i</sub> at least for 10 min. In contrast, 100 μM H<sub>2</sub>O<sub>2</sub> was shown to increase [Ca<sup>2+</sup>]<sub>i</sub> by 42 ± 11.4 nM and 1 mM H<sub>2</sub>O<sub>2</sub> induced a rapid increase in [Ca<sup>2+</sup>]<sub>i</sub> by 235 ± 32.6 nM. However, the Ca<sup>2+</sup> response to 1 mM H<sub>2</sub>O<sub>2</sub> was irreversible, suggesting that this concentration of H<sub>2</sub>O<sub>2</sub> caused cell damage. H<sub>2</sub>O<sub>2</sub>, at concentrations lower than 1 mM, did not induce significant increase in amylase release (data not shown).

#### Effect of H<sub>2</sub>O<sub>2</sub> on CCK-induced Ca<sup>2+</sup> Mobilization


Since Ca<sup>2+</sup> signaling plays a critical role in the regulation of amylase release in rat pancreatic acinar cells, we checked whether 30 μM H<sub>2</sub>O<sub>2</sub> affected CCK-induced Ca<sup>2+</sup> mobilization. As shown in Fig. 3A, 10 pM CCK-8S induced [Ca<sup>2+</sup>]<sub>i</sub> oscillations and an addition of 30 μM H<sub>2</sub>O<sub>2</sub> enhanced the 10 pM CCK-8S-induced [Ca<sup>2+</sup>]<sub>i</sub> oscillations. On the contrary, 30 μM H<sub>2</sub>O<sub>2</sub> did not affect Ca<sup>2+</sup> mobilization caused by 100 pM or higher concentrations of CCK-8S (Fig. 4A and 4B). Scavenging ROS using 1 mM NAC did not affect CCK-8S-induced Ca<sup>2+</sup> mobilization (Fig. 3B and Fig. 4B).

## Discussion


H<sub>2</sub>O<sub>2</sub> is a member of ROS, which cause oxidative damage to cellular components such as lipids, nucleic acids, and proteins. Therefore, H<sub>2</sub>O<sub>2</sub> has generally been considered to



**Fig. 2.** Dose-dependent  $Ca^{2+}$  mobilization by  $H_2O_2$  in rat pancreatic acinar cells. Cells were loaded with fura-2 as described in *Materials and Methods* and changes in  $[Ca^{2+}]_i$  were measured using ratiometric fluorescence imaging. (A) Cells were consecutively exposed to 30  $\mu$ M, 100  $\mu$ M and 1 mM of  $H_2O_2$ . The trace is representative of 18-20 cells in four independent experiments. (B) The size of  $[Ca^{2+}]_i$  increases in respond to  $H_2O_2$  is plotted as a function of the added  $H_2O_2$  concentrations (10  $\mu$ M, 30  $\mu$ M, 100  $\mu$ M, 1 mM and 10 mM). Results are depicted as mean  $\pm$  S.E.M.



**Fig. 3.** Effect of  $H_2O_2$  or NAC, an ROS scavenger, on low concentration of CCK-induced  $[Ca^{2+}]_i$  oscillations in rat pancreatic acinar cells. Cells were exposed to 10 pM CCK-8S to generate  $[Ca^{2+}]_i$  oscillations and then 30  $\mu$ M  $H_2O_2$  (A) or 1 mM NAC (B) was added. Each trace is representative of 18-20 cells in four independent experiments.



**Fig. 4.** Effect of  $H_2O_2$  or NAC, an ROS scavenger, on high concentration of CCK-induced  $[Ca^{2+}]_i$  increases in rat pancreatic acinar cells. (A) Cells were exposed to 1 nM CCK-8S to induce biphasic  $[Ca^{2+}]_i$  increase. The trace is representative of 18-20 cells in four independent experiments. (B) The peak concentrations of intracellular  $Ca^{2+}$  achieved in respond to 1 nM CCK-8S in the presence of absence of 30  $\mu$ M  $H_2O_2$  or 1 mM NAC were measured. Note that  $[Ca^{2+}]_i$  increases induced by CCK-8S were not affected by 30  $\mu$ M  $H_2O_2$  or NAC, an ROS scavenger. Values are means  $\pm$  S.E.M. from 3-5 separate experiments.

be cytotoxic and hazardous to living organisms. However, H<sub>2</sub>O<sub>2</sub> is also known to play a crucial role in the regulation of various physiological functions as an intracellular messenger (Dröge, 2002). The physiologically relevant concentration range of H<sub>2</sub>O<sub>2</sub>, which causes an acceleration of cellular functions in a variety of cell types, is considered 1-100 μM although it depends on cell type (Gamaley and Klyubin, 1999; Dröge, 2002). Since the main aim of this study was to examine the physiological role of ROS in the regulation of amylase release from pancreatic acinar cells, we used 30 μM H<sub>2</sub>O<sub>2</sub>.

In the present study, we found that exogenous addition of 30 μM H<sub>2</sub>O<sub>2</sub> had a dual effect on amylase release in rat pancreatic acinar cells; 30 μM H<sub>2</sub>O<sub>2</sub> enhanced amylase release caused by low concentration of CCK whereas it inhibited high concentration of CCK-induced amylase release. An article published recently by Granados and colleagues (2006) also showed that H<sub>2</sub>O<sub>2</sub> exerted an inhibitory effect on CCK-induced amylase secretion. However, our findings are contrary to those reported by Granados *et al.*, which demonstrated that H<sub>2</sub>O<sub>2</sub> inhibited amylase releases caused by both low and high concentration of CCK. One possible explanation for this discrepancy would be the use of different animal species. Further study might be necessary for elucidating the differences.

Amylase release triggered by CCK receptor activation is biphasic; CCK increased amylase release in a dose-dependent manner but concentrations above 1 nM caused a reduction in amylase release. This is because the rat pancreatic acinar cells possess two different receptor states, high affinity receptors and low affinity receptors (Williams and Blevins, 1993). The activation of the high affinity CCK receptors enhances amylase release, whereas the low affinity CCK receptor causes the inhibition of amylase release.

Activation of the CCK receptors is also known to induce two different patterns of Ca<sup>2+</sup> responses. The high affinity CCK receptors have been known to elicit [Ca<sup>2+</sup>]<sub>i</sub> oscillations, while the low affinity CCK receptors are responsible for the biphasic increase in [Ca<sup>2+</sup>]<sub>i</sub>, consisting of a rapid increase in [Ca<sup>2+</sup>]<sub>i</sub> followed by a slow decrease to a sustained elevated level (Matozaki *et al.*, 1990).

Ca<sup>2+</sup> plays a pivotal role in the regulation of a diverse range of cellular functions, such as muscle contraction, secretion, synaptic plasticity, cell proliferation and cell death (Berridge *et al.*, 2000). Cellular Ca<sup>2+</sup> signals generally encode information in two different modes; frequency-modulated and amplitude-modulated signals (Berridge, 1997). Frequency-modulated Ca<sup>2+</sup> signaling (i.e. Ca<sup>2+</sup> oscillations) is generally considered to have the highest fidelity, and many cells use this paradigm in response to low physiological concentrations of agonists (Berridge *et al.*, 1998).

Since we also observed that low concentration of CCK-8S induced [Ca<sup>2+</sup>]<sub>i</sub> oscillations and amylase release, we first examined whether H<sub>2</sub>O<sub>2</sub>-enhanced amylase release was due to the increase in the CCK-induced [Ca<sup>2+</sup>]<sub>i</sub> oscillations.

Although 30 μM H<sub>2</sub>O<sub>2</sub> did not mobilize Ca<sup>2+</sup> by itself in our system, it obviously enhanced the CCK-induced [Ca<sup>2+</sup>]<sub>i</sub> oscillations. Pancreatic acinar cells are not unique in this respect. H<sub>2</sub>O<sub>2</sub> was also shown to enhance [Ca<sup>2+</sup>]<sub>i</sub> oscillations induced by histamine in human aortic endothelial cells (Hu *et al.*, 2002). In this cell type and rat astrocytes, H<sub>2</sub>O<sub>2</sub> was shown to induce [Ca<sup>2+</sup>]<sub>i</sub> oscillations by itself. The mechanisms by which H<sub>2</sub>O<sub>2</sub> enhances or induces [Ca<sup>2+</sup>]<sub>i</sub> oscillations in various cells are controversial. In endothelial cells, increasing the sensitivity of endoplasmic reticulum (ER) to IP<sub>3</sub> was suggested to be responsible for the H<sub>2</sub>O<sub>2</sub>-induced [Ca<sup>2+</sup>]<sub>i</sub> oscillations (Hu *et al.*, 2000; 2002). In contrast to this, increased IP<sub>3</sub> production by the activation of PLCγ1 was shown to play a critical role in the H<sub>2</sub>O<sub>2</sub> induced [Ca<sup>2+</sup>]<sub>i</sub> oscillations in cultured rat astrocytes (Hong *et al.*, 2006). Although the mechanism responsible for the enhancement of CCK-induced [Ca<sup>2+</sup>]<sub>i</sub> oscillations by H<sub>2</sub>O<sub>2</sub> was not elucidated in this study, both increased IP<sub>3</sub> production and sensitivity of ER to IP<sub>3</sub> could be involved.

Although H<sub>2</sub>O<sub>2</sub> increased low concentration of CCK-induced amylase release, it did not increase the amylase release induced by high concentrations of CCK. Instead, it decreased the amylase release. In addition to this, MnTBAP was shown to increase amylase release induced by high concentrations of CCK. Since it has been shown that high concentration of CCK accumulates ROS in rat pancreatic acinar cells (Granados *et al.*, 2004), our data suggest that ROS produced by high concentration of CCK may contribute to the inhibition of amylase release. The mechanism for the inhibition of amylase release by H<sub>2</sub>O<sub>2</sub> was not clear, but our data indicated that high concentration of CCK-induced Ca<sup>2+</sup> mobilization was not affected. Therefore, H<sub>2</sub>O<sub>2</sub> might act on the downstream signal of [Ca<sup>2+</sup>]<sub>i</sub> increase, leading to the inhibition of exocytosis of the secretory granules. In support of our observation, Rosado and colleagues reported that H<sub>2</sub>O<sub>2</sub> induced temporal and spatial modifications in actin filament reorganization, which was responsible for the decrease in the CCK-stimulated pancreatic secretion (Rosado *et al.*, 2002). The exact mechanism for the inhibition of high-concentration of CCK-induced amylase release caused by H<sub>2</sub>O<sub>2</sub> remains to be elucidated.

Therefore, it was concluded that, although 30 μM H<sub>2</sub>O<sub>2</sub> did not induce [Ca<sup>2+</sup>]<sub>i</sub> oscillations by itself, it increased the low concentrations of CCK-induced [Ca<sup>2+</sup>]<sub>i</sub> oscillations and amylase release. However, 30 μM H<sub>2</sub>O<sub>2</sub> was shown to inhibit the amylase release caused by high concentrations of CCK and the inhibitory effect did not appear to be mediated by inhibition of Ca<sup>2+</sup> signaling. A further study will be needed to elucidate the inhibitory mechanisms.

---

## Acknowledgments

This work was supported by the Korea Research Foundation Grant (KRF-2003-041-E00076).

## References

- Akaishi, T., Nakazawa, K., Sato, K., Saito, H., Ohno, Y., Ito, Y.: Hydrogen peroxide modulates whole cell  $\text{Ca}^{2+}$  currents through L-type channels in cultured rat dentate granule cells. *Neurosci. Lett.* **356**:25-28, 2004.
- Bae, Y.S., Kang, S.W., Seo, M.S., Baines, I.C., Tekle, E., Chock, P.B., Rhee, S.G.: Epidermal growth factor (EGF)-induced generation of hydrogen peroxide. Role in EGF receptor-mediated tyrosine phosphorylation. *J. Biol. Chem.* **272**:217-221, 1997.
- Bae, Y.S., Sung, J.Y., Kim, O.S., Kim, Y.J., Hur, K.C., Kazlauskas, A., Rhee, S.G.: Platelet-derived growth factor-induced  $\text{H}_2\text{O}_2$  production requires the activation of phosphatidylinositol 3-kinase. *J. Biol. Chem.* **275**:10527-10531, 2000.
- Bernfeld, P.: Amylase alpha and beta. *Meth. Enzymol.* **5**:139-148, 1955.
- Berridge, M.J.: The AM and FM of calcium signalling. *Nature.* **386**:759-760, 1997.
- Berridge, M.J., Bootman, M.D., Lipp, P.: Calcium--a life and death signal. *Nature.* **395**:645-648, 1998.
- Berridge, M.J., Lipp, P., Bootman, M.D.: The versatility and universality of calcium signalling. *Nat. Rev. Mol. Cell. Biol.* **1**:11-21, 2000.
- Cancela, J.M., Gerasimenko, O.V., Gerasimenko, J.V., Tepikin, A.V., Petersen, O.H.: Two different but converging messenger pathways to intracellular  $\text{Ca}^{2+}$  release: the roles of nicotinic acid adenine dinucleotide phosphate, cyclic ADP-ribose and inositol trisphosphate. *EMBO J.* **19**:2549-2557, 2000.
- Dröge, W.: Free radicals in the physiological control of cell function. *Physiol. Rev.* **82**:47-95, 2002.
- Favero, T.G., Zable, A.C., Abramson, J.J.: Hydrogen peroxide stimulates the  $\text{Ca}^{2+}$  release channel from skeletal muscle sarcoplasmic reticulum. *J. Biol. Chem.* **270**:25557-25563, 1995.
- Gamaley, I.A., Klyubin, I.V.: Roles of reactive oxygen species: signaling and regulation of cellular function. *Int. Rev. Cytol.* **188**:203-255, 1999.
- Granados, M.P., Salido, G.M., Gonzalez, A., Pariente, J.A.: Dose-dependent effect of hydrogen peroxide on calcium mobilization in mouse pancreatic acinar cells. *Biochem. Cell Biol.* **84**:39-48, 2006.
- Granados, M.P., Salido, G.M., Pariente, J.A., Gonzalez, A.: Generation of ROS in response to CCK-8 stimulation in mouse pancreatic acinar cells. *Mitochondrion.* **3**(5):285-296, 2004.
- Grynkiewicz, G., Poenie, M., Tsien, R.Y.: A new generation of  $\text{Ca}^{2+}$  indicators with greatly improved fluorescence properties. *J. Biol. Chem.* **260**: 3440-3450, 1985.
- Hong, J.H., Moon, S.J., Byun, H.M., Kim, M.S., Jo, H., Bae, Y.S., Lee, S.I., Bootman, M.D., Roderick, H.L., Shin, D.M., Seo, J.T.: Critical role of PLC $\gamma$ 1 in the generation of  $\text{H}_2\text{O}_2$ -evoked  $[\text{Ca}^{2+}]_i$  oscillations in cultured rat cortical astrocytes. *J. Biol. Chem.* **281**:13057-13067, 2006.
- Hu, Q., Zheng, G., Zweier, J.L., Deshpande, S., Irani, K., Ziegelstein, R.C.: NADPH oxidase activation increases the sensitivity of intracellular  $\text{Ca}^{2+}$  stores to inositol 1,4,5-trisphosphate in human endothelial cells. *J. Biol. Chem.* **275**:15749-15757, 2000.
- Hu, Q., Yu, Z.X., Ferrans, V.J., Takeda, K., Irani, K., Ziegelstein, R.C.: Critical role of NADPH oxidase-derived reactive oxygen species in generating  $\text{Ca}^{2+}$  oscillations in human aortic endothelial cells stimulated by histamine. *J. Biol. Chem.* **277**:32546-32551, 2002.
- Matozaki, T., Goke, B., Tsunoda, Y., Rodriguez, M., Martinez, J., Williams, J.A.: Two functionally distinct cholecystokinin receptors show different modes of actions on  $\text{Ca}^{2+}$  mobilization and phospholipid hydrolysis in isolated rat pancreatic acini. *J. Biol. Chem.* **265**:6247-6254, 1990.
- Ohba, M., Shibamura, M., Kuroki, T., Nose, K.: Production of hydrogen peroxide by transforming growth factor-beta 1 and its involvement in induction of egr-1 in mouse osteoblastic cells. *J. Cell Biol.* **126**:1079-1088, 1994.
- Pignatelli, P., Pulcinelli, F.M., Lenti, L., Gazzaniga, P.P., Violi, F.: Hydrogen peroxide is involved in collagen-induced platelet activation. *Blood.* **91**:484-490, 1998.
- Redondo, P.C., Salido, G.M., Rosado, J.A., Pariente, J.A.: Effect of hydrogen peroxide on  $\text{Ca}^{2+}$  mobilisation in human platelets through sulphhydryl oxidation dependent and independent mechanisms. *Biochem. Pharmacol.* **67**:491-502, 2004.
- Rosado, J.A., Gonzalez, A., Salido, G.M., Pariente, J.A.: Effects of reactive oxygen species on actin filament polymerisation and amylase secretion in mouse pancreatic acinar cells. *Cell Signal.* **14**:547-556, 2002.
- Servitja, J.M., Masgrau, R., Pardo, R., Sarri, E., Picatoste, F.: Effects of oxidative stress on phospholipid signaling in rat cultured astrocytes and brain slices. *J. Neurochem.* **75**:788-794, 2000.
- Sobey, C.G., Heistad, D.D., Faraci, F.M.: Mechanisms of bradykinin-induced cerebral vasodilatation in rats. Evidence that reactive oxygen species activate  $\text{K}^+$  channels. *Stroke.* **28**:2290-2294, 1997.
- Stark, H.A., Sharp, C.M., Sutliff, V.E., Martinez, J., Jensen, R.T., Gardner, J.D.: CCK-JMV-180: a peptide that distinguishes high-affinity cholecystokinin receptors from low-affinity cholecystokinin receptors. *Biochim. Biophys. Acta.* **1010**:145-150, 1989.
- Sundaresan, M., Yu, Z.X., Ferrans, V.J., Irani, K., Finkel, T.: Requirement for generation of  $\text{H}_2\text{O}_2$  for platelet-derived growth factor signal transduction. *Science.* **270**:296-299, 1995.
- Toescu, E.C., Lawrie, A.M., Gallacher, D.V., Petersen, O.H.: The pattern of agonist-evoked cytosolic  $\text{Ca}^{2+}$  oscillations depends on the resting intracellular  $\text{Ca}^{2+}$  concentration. *J. Biol. Chem.* **268**:18654-18658, 1993.
- Tournier, C., Thomas, G., Pierre, J., Jacquemin, C., Pierre, M., Saunier, B.: Mediation by arachidonic acid metabolites of the  $\text{H}_2\text{O}_2$ -induced stimulation of mitogen-activated protein kinases (extracellular-signal-regulated kinase and c-Jun NH $_2$ -terminal kinase). *Eur. J. Biochem.* **244**:587-595, 1997.
- Varela, D., Simon, F., Riveros, A., Jorgensen, F., Stutzin, A.: NAD(P)H oxidase-derived  $\text{H}_2\text{O}_2$  signals chloride channel activation in cell volume regulation and cell proliferation. *J. Biol. Chem.* **279**:13301-13304, 2004.

- Wang, X.T., McCullough, K.D., Wang, X.J., Carpenter, G., Holbrook, N.J.: Oxidative stress-induced phospholipase C-gamma 1 activation enhances cell survival. *J. Biol. Chem.* **276**:28364-28371, 2001.
- Wehage, E., Eisfeld, J., Heiner, I., Jungling, E., Zitt, C., Luckhoff, A.: Activation of the cation channel long transient receptor potential channel 2 (LTRPC2) by hydrogen peroxide. A splice variant reveals a mode of activation independent of ADP-ribose. *J. Biol. Chem.* **277**:23150-23156, 2002.
- Williams, J.A.: Intracellular signaling mechanisms activated by cholecystokinin- regulating synthesis and secretion of digestive enzymes in pancreatic acinar cells. *Annu. Rev. Physiol.* **63**:77-97, 2001.
- Williams, J.A., Blevins, G.T. Jr.: Cholecystokinin and regulation of pancreatic acinar cell function. *Physiol. Rev.* **73**: 701-723, 1993.