

Origins of electromagnetic
hypersensitivity to radiation emitted by
WCDMA mobile phones

Min Kyung Kwon

Department of Medical Science

The Graduate School, Yonsei University

Origins of electromagnetic
hypersensitivity to radiation emitted by
WCDMA mobile phones

Directed by Professor Deok Won Kim

The Master's Thesis
submitted to the Department of Medical Science,
the Graduate School of Yonsei University
in partial fulfillment of the requirements for the degree
of Master of Medical Science

Min Kyung Kwon

December 2012

This certifies that the Master's Thesis of
Min Kyung Kwon is approved.

Thesis Supervisor : Deok Won Kim

Thesis Committee Member#1 : Jong-Gwan Yook

Thesis Committee Member#2 : Se Joo Kim

The Graduate School
Yonsei University

December 2012

ACKNOWLEDGEMENTS

의공학을 전공한지 벌써 7년이라는 시간이 흘렀습니다. 이 기간 동안 YBIL 연구실에서의 추억들이 가장 기억에 남을 것 같습니다. 힘들 때마다 아버지처럼 힘이 되어 주시고, 때론 엄격하게 지도해주신 김덕원 교수님께 가장 먼저 감사 드립니다. 연구가 더 나은 방향으로 나아갈 수 있도록 도와 주신 육종관 교수님, 김세주 교수님께도 감사 드립니다. 또한 제가 고민될 때 마다 아낌없이 조언해 주시는 저의 멘토 조종만 교수님께도 감사 드립니다.

석사과정 동안 가족같이 생활했던 연구실 식구들 모두에게도 감사합니다. 처음 연구실 생활을 시작으로 영원히 우정을 함께할 삼총사 재림오빠와 경환 오빠 정말 고마워. 그리고 랩장으로서 최선을 다하는 준열이, 즐거운 수다를 함께한 성권 오빠, 경미와 결혼해서 행복해 하는 정모 오빠, 연구를 진심으로 좋아하는 똑똑한 태근이, 고민을 잘 들어주는 잘생긴 완형 오빠, 앞으로 랩을 책임질 수범 오빠 모두 고마워.

대학교 때부터 함께한 유쾌하고 즐거운 리나, 똑부러지는 보경이, 합창에 푹 빠진 박지은, 코드가 비슷한 이지은! 우리 우정 변치 말자. 그리고 연구를 하면서 힘들 때마다 서로 의지하고, 아낌없이 도와줬던 혜정이와 연희야! 정말 고마워. 정모 오빠와 결혼해서

행복해하는 경미, 서울 생활을 공유한 은혜, 태성이
엄마 정미야! 고등학교 때부터 이어온 우리 우정 변치
말자. 그리고 항상 즐거움을 주고, 힘들 때 마다
아낌없이 격려해준 귀여운 윤성 오빠! 너무 고마워.
또한 학문적, 인생적 선배로서 아낌없이 조언해 주신
성우 선배, 혜진 언니, 동인 선배, 충근 선배, 성현
선배, 이지현 선생님께도 정말 감사합니다.

마지막으로 사랑하는 가족들이 있었기에 제가
행복하게, 걱정 없이 연구할 수 있었습니다. 제가 힘들
때마다 묵묵히 기다려 주고, 하고자 하는 일에
아낌없이 지지해 주시는 아빠, 영원한 나의 베스트
프렌드 엄마, 그리고 하나뿐이 없는 내 동생 기범이!
진심으로 사랑하고 감사합니다.

2012년 12월
권민경 올림

TABLE OF CONTENTS

ABSTRACT	1
I. INTRODUCTION.....	2
II. MATERIALS AND METHODS	4
1. Subjects.....	4
2. Experimental setup	6
3. Experimental procedures	9
4. Physiological measurements.....	10
5. Subjective symptoms and perception of EMF	12
6. Data analysis	13
III. RESULTS.....	15
1. EHS and non-EHS groups	15
2. Physiological variables	15
3. Subjective symptoms	17
4. Perception percentages	19
IV. DISCUSSION	21
V. CONCLUSION	23
REFERENCES	24
ABSTRACT(IN KOREAN).....	28
PUBLICATION LIST	29

LIST OF FIGURES

Figure 1. The measured SAR distribution of WCDMA module	7
Figure 2. Block diagram of exposure setups	8
Figure 3. Experimental procedures for measuring physiological changes and investigating symptoms and perception	10
Figure 4. Percentage of belief of being exposed in EHS and non-EHS groups for sham and real exposure sessions	14

LIST OF TABLES

Table 1. Demographic data of subjects	5
Table 2. Descriptive and statistical tests for heart rate, respiration rate, and LFP/HFP among stage, exposure, and Interaction.....	16
Table 3. All subjective symptoms of the each stage for the real and sham sessions in EHS group.....	17
Table 4. All subjective symptoms of the each stage for the real and sham sessions in non-EHS group	18
Table 5. Percentages of those who believed they were being exposed during pre-exposure, exposure and post-exposure periods, and P-values for sham and real exposures in EHS and non-EHS groups	20

ABBREVIATION

ANOVA: Analysis of variance

ECG: Electrocardiogram

EHS: Electromagnetic hypersensitivity

ELF: Extremely low frequency

ERP: Ear reference point

GSM: Global system for Mobile Communication

HFP: High-frequency power

HRV: Heart rate variability

h/d: Hour per day

IEEE: Institute of Electrical and Electronics Engineers

LFP: Low-frequency power

Max: Maximum

Min: Minimum

N: Number

RF-EMFs: Radio frequency-electromagnetic fields

SAR: Specific absorption rate

SD: Standard deviation

WCDMA: Wideband code division multiple access

yr: Year

YUHS: Yonsei University Hospital System

3G: Third generation

<ABSTRACT>

Origins of electromagnetic
hypersensitivity to radiation emitted
by WCDMA mobile phones

Min Kyung Kwon

*Department of Medical Science
The Graduate School, Yonsei University*

(Directed by Professor Deok Won Kim)

With the use of the third generation (3 G) mobile phones on the rise, social concerns have arisen concerning the possible health effects of radio frequency-electromagnetic fields (RF-EMFs) emitted by wideband code division multiple access (WCDMA) mobile phones in humans. The number of people with self-reported electromagnetic hypersensitivity (EHS), who complain of various subjective symptoms such as headache, dizziness and fatigue, has also increased. However, the origins of EHS remain unclear.

In this double-blind study, two volunteer groups of 17 EHS and 20 non-EHS subjects were simultaneously investigated for physiological changes (heart rate, heart rate variability, and respiration rate), eight subjective symptoms, and perception of RF-EMFs during real and sham exposure sessions. Experiments were conducted using a dummy phone containing a WCDMA module (average power, 24 dBm at 1950 MHz; specific absorption rate, 1.57 W/kg) within a headset placed on the head for 32 min.

WCDMA RF-EMFs generated no physiological changes or subjective symptoms in either group. There was no evidence that EHS subjects perceived RF-EMFs better than non-EHS subjects.

Considering the analyzed physiological data, the subjective symptoms surveyed, and the percentages of those who believed they were being exposed, 32 min of RF radiation emitted by WCDMA mobile phones demonstrated no effects in either EHS or non-EHS subjects.

Key words: provocation, physiological changes, HRV, subjective symptoms, EMF perception

Origins of electromagnetic
hypersensitivity to radiation emitted
by WCDMA mobile phones

Min Kyung Kwon

*Department of Medical Science
The Graduate School, Yonsei University*

(Directed by Professor Deok Won Kim)

I. INTRODUCTION

With the increasing use of third generation (3 G) mobile phones, social concerns have arisen concerning the possible health effects of radio frequency-electromagnetic fields (RF-EMFs) emitted by mobile phones in humans¹. On the basis of limited evidence from both human and animal studies, the World Health Organization has classified RF-EMFs as possibly carcinogenic to humans (Group 2B)². A number of people have self-reported electromagnetic hypersensitivity (EHS), characterized by a variety of non-specific symptoms that differ from individual to individual. Cross-sectional survey studies in different countries have reported that EHS subjects experience non-specific subjective symptoms (e.g., headache, dizziness, fatigue, sleep disorder) associated with EMF exposure: 1.5% in Sweden³, 3.2% in California⁴, and 5% in Switzerland⁵. For some individuals, the symptoms can have lifestyle-changing consequences⁶.

Although numerous studies have examined the effects of Global System for Mobile Communications (GSM) on humans between EHS and non-EHS groups, only a few provocation studies involving WCDMA have simultaneously evaluated physiological changes, subjective symptoms, and EMF perception.

Furubayashi et al. measured psychological and cognitive parameters during pre- and post-exposure⁷. They also monitored physiological parameters, such as skin temperature, heart rate and local blood flow, and asked participants (EHS and non-EHS women) to report on their subjective perception of EMF emitted by WCDMA devices. They concluded that EHS and non-EHS groups did not differ in their responses to real or sham EMF exposure with respect to any psychological, cognitive, or autonomic parameter.

Electromagnetic sensibility in the context of subjective symptoms and perception refers to the ability to perceive EMF without necessarily developing non-specific health symptoms attributable to EMF exposure⁸. Mueller et al. reported no significant differences in the ability to detect EMF between EHS and non-EHS groups⁹. In a study by Hietanen et al., in which EHS subjects were examined for their ability to perceive EMF, none of the subjects could distinguish real EMF exposure from sham exposure¹⁰. Kwon et al. reported that there was no evidence to indicate that EHS subjects could detect EMF exposure¹¹. However, Leitgeb et al. reported that a subset of EHS subjects with significantly increased electromagnetic sensibility could be differentiated from non-EHS groups⁸. Therefore, a comprehensive study is necessary to understand whether EHS is actually caused by exposure to RF-EMFs.

II. MATERIALS AND METHODS

1. Subjects

Because determination of EHS subjects was crucial to this provocation study⁵, we utilized the EHS screening tool developed by Eltiti et al.¹². We adopted the following criteria to identify EHS individuals: (1) a total symptom score greater than or equal to 26 out of a maximum score of 228 (57 symptoms, each ranked from 0 for “not at all” to 4 for “a great deal”); (2) individuals who explicitly attribute their symptoms to exposure to only 3 G mobile phones; and (3) individuals whose current symptoms cannot be explained by a pre-existing chronic illness.

The experiment was performed as a double-blind study with a total of 45 subjects. Initially, 19 EHS and 26 non-EHS subjects were screened; however, two EHS subjects and six non-EHS subjects were excluded. The two EHS subjects were excluded because they were considered outliers in respiration rate, which was greater than two standard deviations from the median (extreme outlier) or 20 beats per min higher than normal without exposure. In the non-EHS group, one subject was excluded because of some drowsiness and motion artifacts during the experiment; three subjects were excluded because they were outliers with respect to heart rate; and two subjects were eliminated because of abnormal electrocardiogram (ECG). None of the EHS or non-EHS subjects failed to attend the second day after attending the first day. Therefore, data from a total of 37 subjects—17 EHS and 20 non-EHS—were analyzed in this study. As shown in Table 1, there were no significant differences in male–female ratio, age, height, weight, body-mass index, nonsmoker-smoker ratio, computer usage time, TV viewing time, or mobile phone usage between the two groups.

The subjects were advised not to consume caffeine, smoke or exercise, and to

sleep enough before the experimental day in order to minimize confounding factors. All subjects, who were recruited by advertisements at the Yonsei University Hospital System (YUHS), were informed of the purpose and procedure of the experiment and were required to give written consent to participate in this study. The Institutional Review Board of the YUHS approved the protocol of this study (project number: 1-2010-0030).

Table 1. Demographic data of subjects¹³

	EHS	Non-EHS	P-value
No. of subjects (n)	17	20	-
Male: female	8: 9	11: 9	0.75
Age (yr)	30.1 ± 7.6	29.4 ± 5.2	0.87
Height (cm)	167.9 ± 7.5	167.6 ± 8.0	0.71
Weight (kg)	63.2 ± 11.9	60.3 ± 11.5	0.44
BMI (kg/m ²)	22.3 ± 2.9	21.3 ± 2.3	0.24
Nonsmoker: smoker	15:2	18:2	1.00
Computer usage time (h/d)	4.4 ± 2.9	5.0 ± 3.8	0.99
TV viewing time (h/d)	1.6 ± 1.3	1.5 ± 1.1	0.96
Mobile phone usage periods (yr)	10.9 ± 3.0	11.6 ± 2.6	0.33

2. Experimental setup

The laboratory was used exclusively for this experiment, and all other electrical devices were unplugged except for our instruments in order to minimize background field levels. Background extremely low frequency (ELF) fields at the level of the head in the laboratory were measured to ensure that they did not influence the subjects. The average ELF electric and magnetic fields were determined to be 1.8 ± 0.0 V/m and 0.02 ± 0.01 μ T, respectively, measured using an electric and magnetic field analyzer (EHP-50C, NARDA-STES, Milano, Italy). The RF field was determined to be 0.05 ± 0.00 V/m with a microwave frequency range from 1920 to 1980 MHz, measured using a radiation meter (SRM 3000, Narda GmbH, Pfullingen, Germany).

To achieve better control over exposure, we used WCDMA modules with Qualcomm chipsets (baseband: MSM6290, RF: RFR6285, power management: PM6658, San Diego, CA, USA) to generate WCDMA RF-EMFs instead of a regular smart phone. The WCDMA modules continuously transmitted at a mean output power of 24 dBm at 1950 MHz, which was measured using a wireless communication test set (E5515C, Agilent, Santa Clara, CA, USA). The modules were inserted into a dummy phone¹⁴, and the location of the module was varied to meet the recommended general public specific absorption rate (SAR)_{1g} of 1.6 W/kg according to the IEEE Standard¹⁵. The SAR measurements were made with a DASY 4 measurement system (SPEAG, Zurich, Switzerland), and a Twin SAM (specific anthropomorphic mannequin) phantom was filled with head tissue-equivalent liquid (mass density, 1000 kg/m³) as specified by the Federal Communications Commission. The measured dielectric properties of the liquid were $\sigma = 1.41$ S/m and $\epsilon_r = 39.7$ for the WCDMA frequency range. When the antenna of the module was positioned 67.5 mm from the ear reference point (ERP) of the dummy, the averaged peak spatial SAR_{1g} was determined to be 1.57 W/kg at 1950 MHz at the left cheek position¹⁶. The electric field and

power drift at the ERP were 6.9 V/m and -0.001 dB, respectively. The measured SAR distribution is shown in Figure 1.




Figure 1. The measured SAR distribution of the WCDMA module on the left side¹³

The module was connected via a 5-m USB cable and a USB type ammeter to a portable laptop computer (X-note R500, LG Electronics, Korea), which controlled the module and monitored electrical current to check exposure conditions (Figure 2). The laptop computer was remotely controlled from another outside desktop computer to satisfy the double-blind study design. The dummy phone was attached to the subject's head using an earplug and headset to fix it at the ERP next to the cheek¹⁷. The phone was held at a distance of 3 mm from the ear using a piece of wood for insulation to prevent battery-generated heat from providing subjects with an indication that the phone was working. The apparatus was constructed from plastic and rubber only, without any metal^{17, 18}.




Figure 2. Block diagram of exposure setups¹³

3. Experimental procedures

No information was given to the subjects except that they would be asked about symptoms and RF-EMFs perception at the beginning of the first experimental day. Sham and real sessions were conducted as a double-blind test to minimize any test bias resulting from a subject and an experimenter recognizing the operational state of the WCDMA module. The experiment was performed for two days, one day for a real session and a second day for a sham session (or vice versa). No matter which came first, sham or real exposure, the second session was always conducted at approximately the same time of the day as the first session in order to maintain the subjects' physiological rhythm. The order of sham and real sessions for each subject was randomly assigned and counterbalanced on our automatic exposure control program using MATLAB 2008a (Mathworks Inc. Natick, MA) to minimize experimental bias. Nine subjects in the EHS group and 11 in the non-EHS group received sham exposure session first. Time duration between sessions was a minimum of one day and a maximum of ten days.

Room temperature and relative humidity, which could considerably affect outcomes, were recorded and maintained. For the non-EHS group, room temperature showed no significant differences between real ($24.4^{\circ}\text{C} \pm 0.9^{\circ}\text{C}$; Min = 23°C , Max = 26°C) and sham ($24.5^{\circ}\text{C} \pm 0.8^{\circ}\text{C}$; Min = 23°C , Max = 26°C) sessions ($P=0.627$). Humidity also showed no significant differences between real ($40.0\% \pm 2.2\%$; Min = 35%, Max = 45%) and sham ($40.8\% \pm 3.3\%$; Min = 35%, Max = 45%) sessions ($P=0.161$). For the EHS group, room temperature showed no significant differences between real ($24.1^{\circ}\text{C} \pm 0.9^{\circ}\text{C}$; Min = 23°C , Max = 26°C) and sham ($24.2^{\circ}\text{C} \pm 1.1^{\circ}\text{C}$; Min = 23°C , Max = 27°C) sessions ($P=0.682$). Humidity also showed no significant differences between real ($40.0\% \pm 2.4\%$; Min = 32%, Max = 45%) and sham ($39.7\% \pm 2.7\%$; Min = 36%, Max = 46%) sessions ($P=0.732$).

4. Physiological measurements

The duration of each exposure session was 64 min, as shown in Figure 3. Before the experiment, subjects were instructed to rest in a sitting position for at least 10 min. Physiological data were collected for 5 min each for four different stages: pre-exposure (stage I), after 11 min of exposure (stage II), after 27 min of exposure (stage III), and post-exposure (stage IV). At each stage, ECG and respiration were simultaneously measured for 5 min (the minimum data requirement for HRV)¹⁹. Heart rate, HRV, and respiration rate were obtained with a computerized polygraph (PolyG-I, Laxtha, Daejeon, Korea) with a sampling frequency of 512 Hz. The data were transferred to a nearby laptop computer (LG Electronics) and analyzed using data acquisition (Telescan 0.9) and analysis (Complexity software) software (Laxtha). The PolyG-I recorded ECG through Ag-AgCl electrodes (2223; 3 M, St. Paul, MN, USA) placed on both arms and the right leg of participants.




Figure 3. Experimental procedures for measuring physiological changes and investigating symptoms and perception. The four shaded areas are periods during which the subjects were questioned regarding the eight symptoms. “o” indicates timing of the inquiries for perception¹³.

Some studies have indicated that EHS subjects may exhibit abnormal autonomic nervous system regulation^{20,21}. Therefore, we first obtained heart rate from ECGs and then acquired HRV and the power spectrum of HRV. High-frequency power (HFP) is reflective of the effects on respiratory sinus arrhythmia, an index of parasympathetic nerve activity, whereas low-frequency power (LFP) is reflective of the effects on both sympathetic and parasympathetic nerves²². In this study, the LFP/HFP ratio was used as an index of autonomic nerve activity balance. Respiratory inductance plethysmography, with an excitation frequency of 3 MHz, was used to measure respiration rate. Subjects wore a coiled band around their upper abdomen for measurement of inductance changes resulting from cross-sectional change.

5. Subjective symptoms and perception of EMF

The four shaded areas in Figure 3 denote periods during which subjects were questioned regarding the eight symptoms; each period lasted approximately 1 min. The eight subjective symptoms of throbbing, itching, warmth, fatigue, headache, dizziness, nausea, and palpation were evaluated through verbal surveys, which were graded on a 4-point scale ranging from 1 (no sensation) to 4 (strong sensation)²³. In addition, perception of EMF exposure was investigated every 5 min throughout the entire session, denoted by an “o” in Figure 3. Subjects were asked to answer the question “Do you believe that you are exposed right now?” nine times during each session. Percentages of those who believed they were being exposed were calculated for pre-exposure, exposure, and post-exposure periods. The total number of inquiries was 185 (5×37) during real exposure and 481 (13×37) during non-exposure; the total number of subjects was 37 ($17 + 20$).

6. Data analysis

A repeated two-way analysis of variance (ANOVA) was performed using SPSS software (SPSS 18, SPSS, Chicago, IL, USA) to investigate differences in heart rate, respiration rate, and relative change in LFP/HFP with exposure and stage for EHS and non-EHS groups. A P-value < 0.05 was considered statistically significant. Subjective symptoms, which are ordered paired data, were analyzed using a non-parametric Wilcoxon signed-rank test. A total of 64 P-values ($4 \text{ stages} \times 8 \text{ symptoms} \times 2 \text{ groups}$) were obtained for the real and sham exposure sessions for the eight symptoms at four stages in both groups. The significance level was adjusted to 0.0125 ($0.05/4$) because testing was performed in four stages.

There were two exposure sessions for each participant, and nine perception inquiries for each session, as shown in Figure 3. For each session, there was one inquiry during pre-exposure, five inquiries during sham or real exposure, and three inquiries during post-exposure. In both groups, the percentages of those who believed they were being exposed were obtained and evaluated for significant differences between real and sham sessions using the McNemar's test. The pre-exposure period of the sham sessions was compared with that of the real sessions to test whether the conditions before sham and real exposures of subjects were the same. The sham exposure period was compared with the real exposure period to test whether the subjects could detect the fields. The post-exposure period after sham exposure was compared with the post-exposure period after real exposure to test whether the real exposure influenced the perception of exposure in the post-exposure period. The Chi-square test was applied to evaluate differences in the percentages of those who answered "yes", which were ordinal data, as shown in Figure 4.




Figure 4. Percentage of belief of being exposed in EHS and non-EHS groups for sham (A) and real (B) exposure sessions. Asterisks indicate statistical significance in perception percentages between EHS and non-EHS groups. Bars indicate standard errors¹³.

III. RESULTS

1. EHS and non-EHS groups

The symptom scores for EHS and non-EHS groups obtained using the Eltiti scale were 53.9 ± 28.5 and 9.3 ± 7.4 (mean \pm S.D), respectively, and they were significantly different ($P < 0.001$). The most typical symptoms reported in the EHS group ($n = 17$) among 57 subjective symptoms on the questionnaire (multiple answers allowed) were fatigue ($n = 17$), headaches ($n = 17$), heaviness in the head ($n = 17$), exhaustion ($n = 15$), migraine ($n = 15$), sleep disturbance ($n = 15$), vertigo ($n = 14$), and difficulty in focusing attention ($n = 14$). The most typical symptoms reported in the non- EHS group ($n = 20$) were fatigue ($n = 14$), blurry vision ($n = 10$), difficulty in concentration ($n = 10$), heaviness in the head ($n = 9$), difficulty in focusing attention ($n = 8$), headaches ($n = 6$), migraine ($n = 6$), and pain/warmth in the head ($n = 6$).

2. Physiological variables

Heart rate, respiration rate, and LFP/HFP ratios of the non-EHS and EHS groups during real and sham exposure are shown in the top section of Table 2. For analysis of the relative changes in LFP/HFP, LFP/HFP values for real and sham were expressed relative to the corresponding stage I values (defined as 100%) because of large individual variation. A repeated two-way ANOVA showed no significant differences in heart rate, respiration rate, or LFP/HFP for stage and exposure in either group, except for LFP/HFP for stage in both groups, as shown in the bottom section of Table 2. For the non-EHS group, LFP/HFP showed no significant difference between real and sham exposures ($P=0.552$), but did show a significant difference among stages ($P=0.001$). For the EHS group, LFP/HFP was also not significantly different between real and sham

exposures (P=0.079), but was significantly different among stages (P=0.048).

Table 2. Descriptive and statistical tests for heart rate, respiration rate, and LFP/HFP among stage, exposure, and interaction¹³

	Heart rate (bpm)				Respiration rate (bpm)				LFP/HFP (%)			
	Non-EHS		EHS		Non-EHS		EHS		Non-EHS		EHS	
	Real	Sham	Real	Sham	Real	Sham	Real	Sham	Real	Sham	Real	Sham
Stage: mean (standard error)												
I	76.0 (1.7)	75.6 (2.5)	77.0 (2.8)	77.2 (2.8)	17.2 (0.6)	17.3 (0.6)	17.4 (0.6)	18.0 (0.8)	100.0 (0.0)	100.0 (0.0)	100.0 (0.0)	100.0 (0.0)
II	75.5 (1.6)	75.3 (2.6)	77.8 (2.9)	77.2 (2.8)	17.3 (0.7)	17.9 (0.5)	17.6 (0.6)	17.0 (0.7)	143.9 (27.0)	165.6 (12.8)	133.8 (15.0)	122.7 (17.0)
III	75.2 (1.7)	74.4 (2.2)	76.4 (2.7)	77.6 (2.9)	16.9 (0.7)	17.6 (0.5)	17.5 (0.6)	17.3 (0.6)	151.0 (31.5)	167.6 (23.4)	198.3 (32.8)	110.6 (13.7)
IV	75.1 (1.6)	73.3 (2.1)	76.9 (2.8)	77.6 (2.9)	18.4 (0.7)	17.7 (0.5)	17.1 (0.7)	17.5 (0.7)	131.3 (23.5)	178.0 (19.9)	178.5 (31.0)	141.5 (23.5)
Factor (P-value)												
Exposure	0.629		0.815		0.772		0.754		0.552		0.079	
Stage	0.166		0.727		0.205		0.614		0.001*		0.048*	
Interaction (exposure & stage)	0.621		0.226		0.518		0.431		0.428		0.055	

* P < 0.05, bpm; beats per min

3. Subjective symptoms

Neither the EHS nor the non-EHS group showed significant differences in any of the eight subjective symptoms surveyed (throbbing, itching, warmth, fatigue, headache, dizziness, nausea, and palpitation) between sham and real sessions in any of the four stages (Table 3, 4).

Table 3. Eight subjective symptom scores of the each stage for the real and sham sessions in EHS group¹³

Symptoms	Stage	Sham (M±S.D)	Real (M±S.D)	P-value	Symptoms	Stage	Sham (M±S.D)	Real (M±S.D)	P-value
Itching	I	1.1±0.3	1.3±0.4	0.083	Headache	I	1.4±0.7	1.4±0.6	0.655
	II	1.3±0.4	1.4±0.5	0.317		II	1.8±1.0	1.5±0.6	0.132
	III	1.4±0.6	1.4±0.6	0.705		III	1.8±0.9	1.7±0.9	0.705
	IV	1.3±0.4	1.3±0.6	1.000		IV	2.0±1.0	1.9±1.0	0.608
Throbbing	I	1.6±0.6	1.4±0.5	0.257	Dizziness	I	1.4±0.6	1.4±0.5	0.739
	II	1.8±0.7	1.4±0.6	0.096		II	1.6±0.8	1.6±0.6	0.739
	III	1.9±0.9	1.6±0.6	0.272		III	1.3±0.6	1.8±0.8	0.035
	IV	2.0±1.0	2.0±1.0	0.914		IV	1.7±0.7	1.8±0.8	0.527
Warmth	I	1.8±0.9	1.5±0.6	0.157	Nausea	I	1.1±0.5	1.1±0.3	0.705
	II	1.8±0.7	1.6±0.6	0.366		II	1.2±0.5	1.2±0.4	1.000
	III	1.8±0.8	2.1±0.9	0.234		III	1.2±0.5	1.4±0.8	0.739
	IV	1.6±0.6	1.8±1.0	0.557		IV	1.3±0.6	1.4±0.9	0.257
Fatigue	I	1.7±0.8	1.8±0.8	0.557	Palpitation	I	1.4±0.5	1.5±0.7	0.480
	II	2.0±0.7	2.0±0.6	1.000		II	1.6±0.7	1.4±0.6	0.083
	III	2.2±0.5	2.1±0.8	0.710		III	1.6±0.6	1.6±0.7	0.739
	IV	2.2±0.8	2.1±0.7	1.000		IV	1.6±0.6	1.4±0.6	0.655

M±S.D; Mean±Standard Deviation

Table 4. Eight subjective symptom scores of the each stage for the real and sham sessions in non-EHS group¹³

Symptoms	Stage	Sham (M±S.D)	Real (M±S.D)	P- value	Symptoms	Stage	Sham (M±S.D)	Real (M±S.D)	P- Value
Itching	I	1.0±0.2	1.0±0.0	0.317	Headache	I	1.0±0.2	1.0±0.0	0.317
	II	1.0±0.2	1.0±0.0	0.317		II	1.1±0.3	1.0±0.2	0.157
	III	1.0±0.2	1.0±0.0	0.317		III	1.1±0.3	1.0±0.2	0.564
	IV	1.0±0.2	1.0±0.0	0.317		IV	1.2±0.4	1.2±0.5	1.000
Throbbing	I	1.0±0.0	1.0±0.0	1.000	Dizziness	I	1.1±0.3	1.0±0.0	0.157
	II	1.0±0.0	1.0±0.0	1.000		II	1.1±0.3	1.0±0.2	0.317
	III	1.1±0.3	1.0±0.2	0.317		III	1.1±0.3	1.1±0.3	1.000
	IV	1.0±0.0	1.0±0.2	1.000		IV	1.1±0.4	1.1±0.5	0.655
Warmth	I	1.1±0.3	1.0±0.2	0.564	Nausea	I	1.0±0.0	1.0±0.0	1.000
	II	1.0±0.2	1.1±0.3	0.317		II	1.0±0.2	1.0±0.0	0.317
	III	1.1±0.3	1.1±0.4	0.564		III	1.1±0.4	1.0±0.2	0.317
	IV	1.0±0.2	1.1±0.5	0.257		IV	1.1±0.4	1.1±0.5	1.000
Fatigue	I	1.0±0.0	1.0±0.0	0.317	Palpitation	I	1.1±0.3	1.0±0.0	0.564
	II	1.0±0.2	1.1±0.5	0.414		II	1.0±0.0	1.0±0.0	0.317
	III	1.0±0.2	1.2±0.5	0.414		III	1.0±0.0	1.0±0.0	1.000
	IV	1.1±0.5	1.2±0.5	0.655		IV	1.0±0.0	1.0±0.0	0.317

M±S.D; Mean±Standard Deviation

4. Perception percentages

Table 5 shows the percentage of subjects who believed they were being exposed during pre-exposure, exposure (real or sham), and post-exposure in the EHS and non-EHS groups. To compare the percentages of those perceiving exposure during experimental sessions, we applied the McNemar's test and found no significant difference between real and sham exposures in the EHS ($P=0.572$) or non-EHS ($P=0.375$) groups. To test whether there were any delayed effects of real exposure on post-exposure perception, we applied the same test and found no significant difference in the percentages of those who believed they were being exposed following real and sham exposures in the EHS ($P=1.000$) or non-EHS ($P=1.000$) groups. There was also no significant difference during pre-exposure between real and sham exposures in EHS ($P=1.000$) and non-EHS ($P=1.000$) groups, indicating that the conditions experienced by subjects before real and sham exposures were the same. Similarly, Kruskal-Wallis tests showed that the percentages of those who believed they were being exposed among pre-, sham exposure, and post-exposure were not significantly different in the EHS ($P=0.263$) or non-EHS ($P=0.426$) groups, demonstrating that conditions were the same for subjects throughout sham-exposure sessions.

Figure 4 shows the percentages of subjects in the EHS and non-EHS groups for each inquiry number who believed they were being exposed in sham (Figure 4A) and real (Figure 4B) exposure sessions. Although there were significant differences between EHS and non-EHS groups during the real exposure period in Figure 4B, there were also significant differences during the sham exposure period (Figure 4A), suggesting that the significant differences between EHS and non-EHS groups during the real exposure period were not actually caused by exposure. The same reasoning applies to the significant differences during pre- and post-exposure in both sham and real exposure sessions. These higher

percentages in the EHS group during both the sham and real sessions probably resulted from a bias of EHS individuals, who believe they can feel EMF, as described in our previous reports^{24,25}. Therefore, there is no evidence that individuals in the EHS group perceived the radiation emitted by WCDMA mobile phones better than those in the non-EHS group.

Table 5. Percentages of those who believed they were being exposed during pre-exposure, exposure and post-exposure periods, and P-values for sham and real exposures in EHS and non-EHS groups¹³

Group	Session	Pre-exposure (%)	P-value	Exposure (%)	P-value	Post-exposure (%)	P-value
EHS (n = 17)	Real	47.1	1.000	65.9	0.572	62.8	1.000
	Sham	41.2		61.2		62.8	
Non-EHS (n = 20)	Real	0.0	1.000	5.0	0.375	6.7	1.000
	Sham	0.0		8.0		6.7	

IV. DISCUSSION

Neither the EHS nor the non-EHS group showed significant differences in heart or respiration rate between real and sham exposures or among stages. In the case of LFP/HFP, however, there were significant differences between some stages during both real and sham exposure sessions in both groups. One disadvantage of the LFP/ HFP analysis is that it is considerably influenced by stress, which can increase or decrease LFP/HFP²⁶. Hjortskov et al. reported that psychological stress could result in increased LFP/HFP²⁷. Nam et al. reported that LFP/HFP monotonically increased at each exposure stage in both EHS and non-EHS groups during 30 min of sham exposure²⁴. In a subsequent study, Nam et al. also confirmed that LFP/HFP significantly increased over time in the absence of exposure, an effect the authors attributed to acute sleep deprivation resulting from awakening subjects with a noise when they exhibited drowsiness²⁸. An additional potential source of stress was the requirement that subjects not move during a 64 min experiment. In fact, the “no-movement” requirement was the factor that drew the most complaints by subjects.

In the current study, neither the EHS nor non-EHS group showed significant differences in any of the four stages between real and sham sessions for any of the eight symptoms surveyed. Wilén et al., reported that exposure to RF-EMFs cannot explain perceived mobile phone attributed symptoms in EHS or non-EHS subjects²⁹. Koivisto et al. also reported that RF exposure did not produce any consistent subjective symptoms or sensations such as headache, dizziness, and fatigue in non-EHS subjects²³. Therefore, most likely, subjective symptoms resulted from a placebo effect, meaning adverse symptoms occurred due to negative expectations³⁰.

There were no significant differences in the percentages of perception in either group who believed they were being exposed during pre- or post-exposure periods between real and sham exposures. There were also no

significant differences in the perception percentages for either the EHS or non-EHS group during the sham exposure session (pre-exposure, sham exposure, post-exposure). Therefore, our experimental protocol seems minimally biased since we confirmed that there were no delayed effects, no differences in pre-exposure condition, and no difference in the percentage of those who believed they were being exposed among the pre-exposure, sham exposure, and post-exposure periods. With regard to the outliers, we included subjects who were outliers in the analyses and tested again to see whether their inclusion actually changed statistical tests for the physiological variables, symptoms, and perception. These results including the outliers were not significantly different from those excluding the outliers.

In this study, the subjects had only two choices, “yes” or “no”, to the perception inquiry of RF-EMFs. However, it could have been biased against subjects who were not sure. For future study, it is recommended to give subjects another choice, “unsure”, and to exclude these answers in calculating the perception accuracy.

We used the EHS screening tool developed by Eltiti et al. to identify individuals who were sensitivity to RF-EMFs¹². There is no objective diagnostic criterion for classifying someone as EHS at present. In the future, the statistical weighing of people’s self-reported hypersensitivity should substantiate their EHS claim.

V. CONCLUSION

In both the EHS and non-EHS groups, there were no significant differences in heart rate, respiration rate, or LFP/HFP between sham and real exposure to a WCDMA module (average power, 24 dBm at 1950 MHz; specific absorption rate, 1.57 W/kg) attached inside a dummy phone for 32 min. There was no association between eight subjective symptoms and RF-EMFs exposure in either group. There was also no indication that EHS subjects could detect exposure. Therefore, considering the physiological data analyzed, the subjective symptoms surveyed, and the percentages of those who believed they were being exposed, no effects were observed in EHS or non-EHS subjects as a result from 32 min of RF radiation emitted by WCDMA mobile phones.

REFERENCES

1. Heinrich S, Thomas S, Heumann C, vonKries R, Radon K. Association between exposure to radiofrequency electromagnetic fields assessed by dosimetry and acute symptoms in children and adolescents: a population based cross-sectional study. *Environ Health* 2010; 9:1-9.
2. Baan R, Grosse Y, Lauby-Secretan B, El Ghissassi F, Bouvard V, Benbrahim-Tallaa L, et al. Carcinogenicity of radiofrequency electromagnetic fields. *Lancet Oncol* 2011; 12:624–6.
3. Hillert L, Berglind N, Arnetz BB, Bellander T. Prevalence of self-reported hypersensitivity to electric or magnetic fields in a population-based questionnaire survey. *Scand J Work Environ Health* 2002; 28:33–41.
4. Levallois P, Neutra R, Lee G, Hristova L. Study of self-reported hypersensitivity to electromagnetic fields in California. *Environ Health Perspect* 2002; 110:619–23.
5. Schröttner J, Leitgeb N, Hillert L. Investigation of electric current perception thresholds of different EHS groups. *Bioelectromagnetics* 2007; 28:208–13.
6. Proceedings of an International Workshop on EMF Hypersensitivity. 25-7 October 2004. In Edited by Mild KH, Repacholi M, van Deventer E, Ravazzani P. World Health Organization; 2006.
7. Furubayashi T, Ushiyama A, Terao Y, Mizuno Y, Shirasawa K, Pongpaibool P, et al. Effects of short-term W-CDMA mobile phone base station exposure on women with or without mobile phone related symptoms. *Bioelectromagnetics* 2009; 30:100–13.
8. Leitgeb N, Schrottner J. Electrosensibility and electromagnetic hypersensitivity. *Bioelectromagnetics* 2003; 24:387–94.
9. Mueller CH, Krueger H, Schierz C. Project NEMESIS: perception of a 50 Hz electric and magnetic field at low intensities (laboratory experiment). *Bioelectromagnetics* 2002; 23:26–36.

10. Hietanen M, Hamalainen AM, Husman T. Hypersensitivity symptoms associated with exposure to cellular telephones: no causal link. *Bioelectromagnetics* 2002; 23:264–70.
11. Kwon MS, Koivisto M, Laine M, Hamalainen H. Perception of the electromagnetic field emitted by a mobile phone. *Bioelectromagnetics* 2008; 29:154–9.
12. Eltiti S, Wallace D, Zougkou K, Russo R, Joseph S, Rasor P, et al. Development and evaluation of the electromagnetic hypersensitivity questionnaire. *Bioelectromagnetics* 2007; 28:137–51.
13. Kwon MK, Choi JY, Kim SK, Yoo TK, Kim DW. Effects of radiation emitted by WCDMA mobile phones on electromagnetic hypersensitive subjects. *Environ Health* 2012; 11(69):1-8.
14. Croft RJ, Leung S, McKenzie RJ, Loughran SP, Iskra S, Hamblin DL. Effects of 2 G and 3 G mobile phones on human alpha rhythms: Resting EEG in adolescents, young adults, and the elderly. *Bioelectromagnetics* 2010; 31:434–4.
15. IEEE Standard 1528–2003. Recommended Practice for Determining the Peak Spatial-Average Specific Absorption Rate (SAR) in the Human Head From Wireless Communications Devices: Measurement Techniques. IEEE Standard Coordinating Committee 2003.
16. Beard BB, Kainz W, Onishi T, Iyama T, Watanabe S, Fujiwara O. Comparisons of computed mobile phone induced SAR in the SAM phantom to that in anatomically correct models of the human head. *IEEE Trans Electromagn Compat* 2006; 48:397–407.
17. Haarala C, Takio F, Rintee T, Laine M, Koivisto M, Revonsuo A, et al. Pulsed and continuous wave mobile phone exposure over left versus right hemisphere: effects on human cognitive function. *Bioelectromagnetics* 2007; 28:289–95.
18. Unterlechner M, Sauter C, Schmid G, Zeitlhofer J. No effect of an UMTS mobile phone-like electromagnetic field of 1.97 GHz on human attention and

reaction time. *Bioelectromagnetics* 2008; 29:145–53.

19. Marek MJ, Thomas BA, John C, Bobert EK, Alberto M, Arthu JM. Heart rate variability: standards of measurement, physiological interpretation, and clinical use. *Eur Heart J* 1996; 17:354–81.

20. Lyskov E, Sandström M, Mild KH. Neurophysiological study of patients with perceived ‘electrical hypersensitivity’. *Int J Psychophysiol* 2001; 42:233–41.

21. Sandström M, Lyskov E, Hörnsten R, Mild KH, Wiklund U, Rask P. Holter ECG monitoring in patients with perceived electrical hypersensitivity. *Int J Psychophysiol* 2003; 49:227–35.

22. Parazzini M, Ravazzani P, Tognola G, Thuroczy G, Molnar FB, Sacchettini A, et al. Electromagnetic fields produced by GSM cellular phones and heart rate variability. *Bioelectromagnetics* 2007; 28:122–9.

23. Koivisto M, Haarala C, Krause CM, Revonsuo A, Laine M, Hamalainen H. GSM phone signal does not produce subjective symptoms. *Bioelectromagnetics* 2001; 22:212–5.

24. Nam KC, Lee JH, Noh HW, Cha EJ, Kim NH, Kim DW. Hypersensitivity to RF fields emitted from CDMA cellular phones: a provocation study. *Bioelectromagnetics* 2009; 30:641–50.

25. Kim DW, Choi JL, Nam KC, Yang DI, Kwon MK. Origins of electromagnetic hypersensitivity to 60 Hz magnetic fields: A provocation study. *Bioelectromagnetics* 2012; 33:326–33.

26. Akselrod S, Gordon D, Ubel FA, Shannon DC, Berger AC, Cohen RJ. Power spectrum analysis of heart rate fluctuation: a quantitative probe of beat-to-beat cardiovascular control. *Science* 1981; 213:220–2.

27. Hjortskov N, Rissen D, Blangsted AK, Fallentin N, Lundberg U, Sjøgaard K. The effect of mental stress on heart rate variability and blood pressure during computer work. *Eur J Appl Physiol* 2004; 92:84–9.

28. Nam KC, Kwon MK, Kim DW. Effects of posture and acute sleep

deprivation on heart rate variability. *Yonsei Med J* 2011; 52:569–73.

29. Wilén J, Johansson A, Kalezic N, Lyskov E, Sandström M. Psychophysiological tests and provocation of subjects with mobile phone related symptoms. *Bioelectromagnetics* 2006; 27:204–14.

30. Oftedal G, Straume A, Johnsson A, Stovner LJ. Mobile phone headache: a double blind, sham-controlled provocation study. *Cephalalgia* 2007; 27:447–55.

< ABSTRACT (IN KOREAN)>

WCDMA 휴대폰 전자파 과민 증후군의 호소증상 원인 규명

<지도교수 김 덕 원>

연세대학교 대학원 의과학과

권 민 경

스마트 폰 사용량이 기하급수학적으로 증가함에 따라 WCDMA 휴대폰 전자파가 인체에 미치는 영향에 대한 관심이 고조되고 있다. 이와 더불어 휴대폰 전자파에 의해 두통, 어지러움, 피로 등의 주관적인 증상을 호소하는 전자파 과민 증후군 (EHS: electromagnetic hypersensitivity)이 증가하고 있다. 그러나 이러한 증상이 스마트 폰 전자파 노출에 의한 것인지 심리적 요인 때문인지 확실한 원인 규명이 되어 있지 않다.

이에 본 연구는 이중 맹검법을 이용하여 17명의 EHS 군과 20명의 일반인 군을 대상으로 전자파 노출여부에 따른 생리학적 변화(심박수, 심박 변이도, 호흡수)와 자각 증상(가려움, 욕신거림, 뜨거운 느낌, 피로, 두통, 어지러움, 메스꺼움, 가슴 두근거림), 전자파 인지 여부를 동시에 측정하여 자각 증상 원인을 복합적으로 분석하였다. 실험을 위해서 WCDMA 모듈(과워, 24 dBm; 주파수, 1950 MHz; SAR_{1g}, 1.57 W/kg)을 dummy 폰에 삽입한 후, 헤드 셋을 이용하여 32분간 두부에 전자파를 노출시켰다.

연구 결과, EHS 군과 일반인 군 모두 WCDMA 전자파 노출에 따른 생리학적 변화, 주관적 자각증상에 영향이 없었다. 또한 EHS군이 일반인 군보다 전자파 인지를 더 잘한다고 볼 수도 없었다. 따라서 생리학적 변화, 주관적 자각증상, 전자파 인지 여부를 다각적으로 종합한 결과, WCDMA 휴대폰 전자파는 EHS 군 뿐만 아니라 일반인군에서도 영향이 없었다고 사료된다.

핵심되는 말 : 역학, 생리학적 변화, 심박 변이도, 자각증상, 인지여부

PUBLICATION LIST

International Journal

1. **Kwon MK**, Choi JY, Kim SK, Yoo TK, Kim DW. Effects of radiation emitted by WCDMA mobile phones on electromagnetic hypersensitive subjects. *Environ Health* 2012; 11(69):1-8.

International Conference

2. **Kwon MK**, Kim SK, Koo JM, Choi JY, Kim DW. EHS subjects do not perceive RF EMF emitted from smart phones better than non-EHS subjects. *Conf Proc 34th IEEE Eng Med Biol Soc, San Diego, USA 2012 (Aug 28 - Sep 1)*.

3. **Kwon MK**, Choi JY, Kim SK, Yoo TK, Kim DW. Perception of RF Fields Emitted from Smart Phones. 34th Bioelectromagnetics Society, Brisbane, Australia 2012 (June 17 - 22) (Student Travel Fund Award).

4. **Kwon MK**, Nam KC, Lee DS, Jang KH, Kim DW. Effects of RF Fields Emitted from Smart Phones on Cardio-Respiratory Parameters: A Preliminary Provocation Study. *Conf Proc 33th IEEE Eng Med Biol Soc, Boston, USA 2011 (Aug 30 - Sep 3)*: 1961-4.

5. **Kwon MK**, Lee DS, Choi JY, Nam KC, Kim DW. Effects of RF fields emitted from smart phones on physiological changes: A preliminary provocation study. 33th Bioelectromagnetics Society, Halifax, Canada 2011 (June 12-17).

Domestic Journal

6. 권민경, 최재림, 최준열, 장경환, 김성권, 김덕원. WCDMA 휴대전화 전자파에 의한 심호흡계 영향. *의공학회지* 2012; 33(2): 59-64.