

**Acetyl-CoA carboxylase β expression
mediated by MyoD and MRF4 is
differentially affected by RAR and RXR**

Joo-Yeon Kim

Department of Medical Science
The Graduate School, Yonsei University

**Acetyl-CoA carboxylase β expressions
mediated by MyoD and MRF4 is
differentially affected by RAR and RXR**

Directed by Professor Kyung-Sup Kim

A Master's Thesis Submitted to the
Department of Medical Science, the Graduate School of
Yonsei University in partial fulfillment of the requirements
for the degree of Master of Science

Joo-Yeon Kim

December, 2002

**This certifies that the master's thesis
of Joo-Yeon Kim is approved.**

Thesis Supervisor: Kyung-Sup Kim

Yong-Ho Ahn

Bong-Soo Cha

The Graduate School
Yonsei University

December, 2002

Table of Contents

Abstract	1
I. Introduction	3
II. Materials and Methods	6
1. Materials	6
2. Cell culture	6
3. Transient transfection	7
4. Luciferase and β -galactosidase assay	8
5. Western immunoblot	9
6. Statistical analysis	9
III. Results	
ACCB induction by retinoids in H9C2 myotubes	10
Suppression of ACCB promoter by retinoid receptors	10
Effect of retinoids on the induction of ACCB promoter by MRFs	10
Retinoid effects on heterologous promoter containing E-box or noble MRF-binding elements	15
IV. Discussion	17
V. Conclusion	20
Reference	21
Abstract (in Korean)	26

List of Figures

Figure 1. Increased expression of ACC β by retinoids in H9C2 myotubes.....	11
Figure 2. RXR α or RAR α effects on ACC β promoter in H9C2 cells	12
Figure 3. Interaction of retinoic acid receptors and muscle specific regulatory factors on ACC β promoter	14
Figure 4. RXR α enhanced MRF4 effect on E-box regions of ACC β and RAR α suppressed MyoD effect on noble <i>cis</i> -element.....	16

Abstract

Acetyl-CoA carboxylase β expression mediated by MyoD and MRF4 is differentially affected by RAR and RXR

Joo-Yeon Kim

Department of medical science

The Graduate School, Yonsei University

(Directed by professor Kyung-Sup Kim)

Mammals have two major isoforms of acetyl-CoA carboxylase (ACC). The 275kDa β -form (ACC β) is predominantly expressed in heart and skeletal muscle while the 265kDa α -form (ACC α) is the major isoform in lipogenic tissues such as liver and adipose tissue. ACC β is thought to control fatty acid oxidation by means of the ability of malonyl-CoA to inhibit carnitine palmitoyl-CoA transferase-1 (CPT-1), which is a rate-limiting enzyme of fatty acid oxidation in mitochondria. Previously, it reported that MyoD and other muscle regulating factors (MRFs) up-regulate the expression of ACC β by interactions between these factors and several *cis*-elements of ACC β promoter. It described here that ACC β

expression mediated by MRFs is regulated by retinoic acids. Endogenous expression of ACC β in differentiated H9C2 myotube was significantly increased by retinoic acid treatment. However, upon transient transfection assay in H9C2 myoblast, ACC β promoter activity was suppressed by RXR α and more severely by RAR α . These effects on ACC β expression in myoblasts and myotubes by RXR α and RAR α seem to be mediated by their interactions with MRFs because no consensus sequence for RXR α and RAR α has been found in ACC β promoter and retinoic acid receptors did not affect this promoter activities by itself. In transient transfection in NIH3T3 fibroblast, the activation of ACC β promoter by MyoD, main MRF in myoblast, was significantly suppressed by RAR α and to a less extent by RXR α while the RXR α dramatically augmented the activation by MRF4, major MRF in myotube. These results explained that retinoic acids differentially affected the action of MRFs according to their types and RXR α specially elevates the expression of muscle specific genes by stimulating the action of MRF4.

Key Words : acetyl-CoA carboxylase β , MyoD, MRF4, RAR α , RXR α

Acetyl-CoA carboxylase β expression mediated by MyoD and MRF4 is differentially affected by RAR and RXR

(Directed by professor Kyung-Sup Kim)

Department of medical science

The Graduate School, Yonsei University

Joo-Yeon Kim

I. Introduction

Acetyl-CoA carboxylase catalyzed the carboxylation of acetyl-CoA to form malonyl-CoA. Mammals have two major isoforms of acetyl-CoA carboxylase. Acetyl-CoA carboxylase α (ACC α) is expressed in lipogenic tissues such as adipose tissue and liver¹. Its product malonyl-CoA is two carbon donor for fatty acid synthesis and its availability is rate limiting factor in fatty acid synthesis. In contrast, acetyl-CoA carboxylase β (ACC β) is predominantly expressed in skeletal muscle and heart^{2,3}. ACC β is known to control fatty acid oxidation by regulating malonyl-CoA levels, around mitochondria. Compared with α -isoform, ACC β has additional hydrophobic N-terminus which contributes to the interaction with outer membrane of mitochondria. Moreover, there are many

specific serine residues to be phosphorylated in response to various hormonal signals. Malonyl-CoA plays a critical role in controlling the activities of carnitine palmitoyl-CoA transferase 1(CPT-1), the rate-limiting step in fatty acid uptake and oxidation by mitochondria⁴⁻⁶. This function of ACC β in muscle contributes to maintaining adult muscle phenotype and controlling energy homeostasis.

Myogenesis is regulated by a family of four muscle regulatory transcription factors such as MyoD⁷, myogenin⁸, Myf-5⁹, and MRF4^{10,11} that share a common dimerization motif, DNA-binding domain and the basic helix-loop-helix motif¹². Muscle regulatory transcription factors bind to a consensus sequence, E-box, located in target gene promoter and activate the transcription of the target genes. Therefore, they affect muscle gene transcriptions in forms of either homodimer or heterodimer with E-protein^{13,14}. Although these factors contain the common motif, each of these factors activates the specific target genes inducing normal muscle development and maintaining normal muscle phenotype. Therefore, the ordered expression of these factors during myogenesis controls the temporal and spatial expression pattern of muscle specific genes, properly. MyoD and Myf5 can be considered as determinant factor of myogenic lineages. Cultured myoblast cell lines express MyoD mRNA, Myf-5 mRNA¹⁵ whereas myogenin mRNA is expressed upon myotube fusion and MRF4 plays major role in high expression of muscle gene in fully differentiated myotube¹⁶⁻¹⁸. Previously, it have been shown that MyoD and MRF4 are able to enhance promoter activity of ACC β ¹⁹. It was also previously reported that induction of ACC β accompanies the differentiation

of H9C2 cells into myocytes and the anti-sense RNA to ACC β mRNA inhibits muscle differentiation²⁰.

Retinoic acids have been known as potential inducer of cell growth arrest and differentiation²³. These receptors act as transcriptional activators by binding as heterodimer or homodimer to specific nucleotide sequence of target genes. It was reported to interact physically muscle b-HLH protein and retinoic acid receptors²¹. The interaction of retinoic acid receptors and myogenic b-HLH proteins is important for the execution of retinoid induced myogenic differentiation²².

In this study, it introduced that the transcriptional activities of MyoD or MRF4 which regulates ACC β promoter activity might be regulated by retinoic acids during muscle differentiation.

. Materials and Methods

1. Materials

9-cis retinoic acid and *all-trans* retinoic acid was purchased from Sigma Aldrich Co (ST, Louis, MO, USA). The concentrations of their stock solutions were adjusted to 1mM in DMSO. H9C2 and NIH3T3 cell lines were obtained from the American Type Culture Collection (ATCC, Rockville, MD, USA). Dulbecco's modified eagle's medium (DMEM), DMEM-F12 media, fetal bovine serum (FBS), horse serum, antibiotics, and LipofectAMINE Plus kit were purchased from Invitrogen (Int. Antwoordnummer, VT Groningen, Nederland). Polyvinylidene fluoride (PVDF) membrane and enhanced chemiluminescence (ECL plus) system was purchased from Amersham (Amersham-Pharmacia, Uppsala, Sweden). Luciferase assay kit and reporter lysis buffer were purchased from Promega (Promega, Medison, WI, USA).

2. Cell Culture

H9C2 Cells were maintained in DMEM-F12 medium supplemented with 10 %(v/v) fetal bovine serum and 100 unit/ml penicillin G sodium, 100mg/ml streptomycin sulfate and 250 ng/ml amphotericin B in 5 % CO₂ at 37 °C. The differentiation medium for H9C2 cell differentiation was DMEM-F12 supplemented with 2 %(v/v) horse serum and 100 unit/ml penicillin G sodium, 100 mg/ml streptomycin sulfate and 250 ng/ml amphotericin B. H9C2 cells at confluent state were differentiated into myotubes by cultivation in differentiation

medium for 6 days. NIH3T3 cells were maintained in DMEM medium with 10 % (v/v) fetal bovine serum and 100 unit/ml penicillin G sodium, 100 mg/ml streptomycin sulfate and 250 ng/ml amphotericin B.

3. Transient transfection

Cells used in this experiment were maintained as monolayer culture and grown in appropriate media. Plasmid DNAs were prepared using QIAGEN Plasmid Midi Kits (Qiagen, Hilden, Germany). Concentration of all plasmids was adjusted to 100 ng/ μ l and confirmed in 1 % agarose gel before transfection. Cells were plated in six-well tissue culture plate at a density of 2×10^5 cells/well in 2ml of media. After 24 hr attachment period of plated cells, transfections were performed with LipofectAMINE and Plus Reagent. According to the manufacturer's protocol, briefly, 500 ng of each construct of pPII β -1317/+17-Luc, pE-box-tk-Luc, and p6X(+8/+27)-tk-Luc, 100ng pCMV β -gal, 200ng of pcDNA3-MyoD or pcDNA3-MRF4, and 100ng of pCMX-RXR α or pCMX-RAR α were mixed. 4 μ l of plus reagent diluted in 100 μ l of media lacking serum and antibiotics was mixed, and then these Plus reagent mixtures were added to DNA mixture. After 15 minutes, LipofectAMINE mixtures which was prepared by mixing 2 μ l of LipofectAMINE and 100 μ l of media lacking serum and antibiotics were added to Plus reagent mixture following further incubation for 15 minutes. Also, cells plated in six-well tissue culture plate were washed by phosphate buffed saline (PBS) and replaced by 800 μ l media lacking serum and

antibiotics. After 15 minutes, the mixtures of DNA, LipofectAMINE and Plus reagent were added in each well. After 3 hr, the medium containing the DNA, LipofectAMINE and Plus reagent complex was removed and replaced by appropriated media containing serum and antibiotics. Cells were then further cultured for 24 hr and treated by *9-cis* retinoic acid or *all-trans* retinoic acid of concentration of 1 μ M for 48 hr.

4. Luciferase and β -galactosidase assay

Transfected cells were harvested in 200 μ l of reporter lysis buffer. The lysed cells were centrifuged to remove cell debris and supernatant was collected. Luciferase activities in lysates were measured, using Luciferase assay kit.

For β -galactosidase assay, 20 μ l of cell lysates were mixed with 180 μ l of assay reagent composed of 3.2 mg of *o*-nitrophenol- β -D-galactopyranoside, 4 ml of 0.1M sodium phosphate buffer adjusting PH7.5, 40 μ l of 100X Mg solution (0.1M MgCl₂, 4.5M β -mercaptoethanol) and OD change 420 nm for a minute at 37 °C was measured in Microplate Spectrometer.

Luciferase activities were expressed as the values normalized by β -galactosidase activity. Each experiment was performed in triplicate.

5. Western immunoblot

Cells in 10-cm plates were washed twice with cold PBS. 400 μ l of digitonin buffer [1 M Tris-HCl (pH 7.5), 0.5 M EDTA (pH8.0), 0.25 % sucrose, 1.5 mM

PMSF, 100 mg/ml digitonin, 10 mM sodium fluoride] were directly added to culture dish on ice. The diffusate was collected and cell debris was removed by centrifugation. The proteins in diffusate were separated on a 5 % SDS-PAGE and transferred to a PVDF membrane. ACC β protein band was detected using streptavidine-HRP conjugate and by ECL plus.

6. Statistical analysis

All transfection studies were performed in three to five separate experiments, where triplicate dishes were used. The data were represented as means \pm standard deviation. Statistical analysis was carried out using Microsoft Excel[®] (Microsoft, Redmond, WA, USA).

. Results

ACC β induction by retinoids in H9C2 myotubes

H9C2 cells are rat heart myoblast cells. H9C2 cells do not express ACC β protein in state of myoblast. But when H9C2 myoblast is fully differentiated into myotube, ACC β is actively expressed (Fig.1, lane1). When we treated *all-trans* or *9-cis* retinoic acid to completely differentiated H9C2 myotubes, ACC β expressions were remarkably increased (Fig.1, lane2 and 3). These results imply that retinoic acids enhance ACC β expression in myotubes.

Suppression of ACC β promoter by retinoid receptors

Even if ACC β is not expressed, ACC β promoter activity is very high in H9C2 myoblast when compared with those in fibroblast or hepatoma cell lines. This is probably due to the presence of MRFs in H9C2 myoblasts. We tested how the retinoid receptors affect the promoter activity in H9C2 myoblasts. Unexpectedly, transient transfection assay shows that ACC β promoter activities were suppressed 38 % by RXR α and more severely 65 % by RAR α (fig. 2).

These results displayed the disagreement of retinoic acid effects on ACC β promoter in myoblast and myotube.

Effects of retinoids on the induction of ACC β promoter by MRFs

Sequence analysis could not reveal any retinoid responsive element in ACC β




Figure 1. Increased expression of ACC β by retinoids in H9C2 myotubes.

H9C2 cells were almost completely differentiated into myotubes by incubation in differentiation medium for 6 days as described in Materials and Methods. Differentiated myotubes were treated with DMSO (lane 1), 1 μ M *all-trans* retinoic acid (lane 2), or 1 μ M *9 cis*-retinoic acid (lane 3) for 24 hours. Thirty μ g of protein of cellular diffusate were subjected to western immunoblot analysis using streptavidin-HRP conjugate.




Figure 2. RXR α or RAR α effects on ACC β promoter in H9C2 cells.

ACC β promoter, pP11 β -1317/+17-Luc, was transfected into H9C2 myoblast, accompanying with RXR α or RAR α -overexpressing construct. The amount of constructs transfected per each well were as following : 0.5 μ g of pP11 β -1317/+17-Luc, 0.1 μ g of pCMV β -galactosidase and 0.1 μ g of overexpression vectors. 24 hr after transfection, the media were replaced, containing 1 μ M of the respective ligand. The luciferase activities were assayed 24 hrs after ligand treatment and expressed as the values standardized with β -galactosidase activities.

promoter. Overexpression of RAR α or RXR α with treatment of their ligands in NIH3T3 fibroblasts did not significantly change the ACC β promoter activities (Fig. 3, A), suggesting that retinoid receptors affect the ACC β promoter not by their direct DNA-binding but probably by the interactions with MRFs existing in myoblasts or myotubes.

The fact that The retinoid and their receptors differentially affected the ACC β promoter activities in myoblasts and myotubes which express different kinds of MRFs led to test the retinoid actions on transcriptional activities of MyoD and MRF4 which are primary MRF in myoblasts and myotubes, respectively. The activation of ACC β promoter by MyoD in NIH3T3 cells was significantly suppressed 56 % by RAR α but was not affected by RXR α . The overexpression of both RAR α and RXR α also suppressed the MyoD activities to a similar extent as RAR α alone. However, the effects of retinoid receptors were completely different on MRF4-mediated activation of ACC β promoter. The activation of ACC β promoter by MRF4 was significantly increased 332 % by RXR α , 95 % by RAR α , and 268 % by both RAR α and RXR α . The RXR α activated MRF4 activities most effectively. These data conclude that the activities of myogenic determinant MyoD was inhibited by RAR α whereas those RAR α and RXR α markedly increase the activities of MRF4 being expressed in terminally differentiated myotubes.




Figure 3. Interaction of retinoic acid receptors and muscle specific regulatory factors on ACC β promoter. ACC β promoter construct, pII β -1317/+17-Luc, was transfected with the constructs expressing β -galactosidase, retinoid receptors, MyoD (A), or MRF4 (B) into NIH3T3 fibroblasts as denoted in above figure. Total DNA amounts for transfection were adjusted to 0.9 μ g per each well with pcDNA3.0, and the amounts of each plasmid are as followings : 0.5 μ g of pII β -1317/+17-Luc construct, 0.1 μ g of pCMV β -galactosidase, 0.1 μ g of pCMX-RAR α or pCMX-RXR α and 0.2 μ g of pcDNA3-MyoD or pcDNA3-MyoD. At 24 hrs after transfection, the media were replaced, containing 1 μ M of the respective ligand. At 48 hrs after transfection, the luciferase activities were measured and shown as the values.

Retinoids effects on heterologous promoters containing E-boxes or noble

MRF-binding elements

It was previously reported that the myogenic regulatory factor-mediated activation of ACC β promoter depends on E-box region from -493 to -403 and one novel *cis*-element from +17 to +24. MRFs are generally known to bind to E-boxes in a form of heterodimer with E-protein whereas the proximal novel sequence of ACC β promoter was bound by MRF homodimer. To test how the retinoid receptors affect the activation by different quaternary forms of MRF, we used the heterologous promoter containing E-box region or six copy of novel *cis*-element of ACC β promoter at upstream of tk minimal promoter. Our data showed that the activation of heterologous promoter through noble *cis*-element by MyoD is suppressed by RAR α and activation of heterologous promoter containing E-box region by MRF4 is enhanced by RXR α (Fig 4). However, RAR α can not inhibit activation of E-box containing promoter by MyoD and RXR α can not increase transcriptional activation by MRF4 through noble *cis*-element.




Figure 4. RXR α enhanced MRF4 effect on E-box regions of ACC β and RAR α suppressed MyoD effect on noble *cis*-element. Reporter constructs, such as pE-box-tk-Luc containing E-box regions(-493/-403) and p6X(+8/+27)-tk-Luc containing six copies of noble *cis*-element, were transfected with overexpression vectors as denoted in above figure into NIH3T3 fibroblasts. Total amounts of DNA for transfection were adjusted to 0.9 μ g with pcDNA3.0 and the amounts of each plasmid used in transfection are as following : 0.5 μ g of the pE-box-tk-Luc or the p6X(+8/+27)-tk-Luc, 0.1 μ g of the pCMV β -galactosidase, 0.2 μ g of the pcDNA3-MRF4 or the pcDNA3-MyoD and 0.1 μ g of the pCMX-RXR α or the pCMX-RAR α . Next day transfection, the cells were treated with 1 μ M of the respective ligand for 24 hrs. The luciferase activities were standardized with β -galactosidase activities.

. Discussion

Myogenesis is regulated by muscle specific regulating factors such as MyoD, myogenin, Myf5 and MRF4. These regulatory proteins are capable of directly activating expression of skeletal muscle specific genes such as those encoding the contractile proteins. Expression of either Myf5 or MyoD is required for skeletal muscle lineage determination and MRF4 is expressed at high level in adult muscle. MyoD is primary MRF expressed in myoblast stage³¹⁻³³, while MRF4 plays a major role in maintaining the high levels of skeletal muscle gene expressions³⁵. Retinoic acid is known as inducer of cell growth arrest and differentiation²³. Their actions are mediated by two families of ligand-inducible transcriptional factors, the retinoic acid receptor (RAR α , β and γ) and retinoid X receptor (RXR α , β and γ), which are members of the nuclear receptor superfamily²⁴⁻²⁶. RARs are bound and activated by *all-trans* retinoic acid^{27,28}, whereas RXRs by *9-cis* retinoic acid^{29,30}. These retinoic acid receptors also show the different expressional patterns according to muscle differentiation stages. RAR α is plentiful and RXR α is scarce in myoblast stage while, in contrast, RXR α is abundant and RAR α is rare in myotube stage³⁴. In this study, it was introduced that ACC β transcriptional induction by muscle specific regulating factors is differentially regulated by retinoic acids. When completely differentiated H9C2 myotubes were treated with retinoids, the levels of ACC β were significantly increased (Fig. 1). However, transient transfection assays in myoblasts, where the ACC β promoter is active probably due to the presence of

MyoD, showed that ACC β promoter activities were suppressed by RXR α and more severely by RAR α (Fig. 2). These data suggested the possibility that retinoid receptors might differentially affect each MRF expressed in stage-specific manner during myogenesis. To check this possibility, the transient transfection assay was performed using NIH3T3 fibroblasts which do not express any MRF. As expected, the data in NIH3T3 showed that RAR α suppressed MyoD-mediated activation of ACC β promoter, but in contrast RXR α rather significantly enhanced MRF4 activities (Fig. 3). These effects of retinoid receptors on ACC β promoter were not mediated by their direct interaction with ACC β promoter sequence, because any conserved responsive element for retinoid receptor can not be found in ACC β promoter and retinoic acid receptors by themselves did not activate ACC β promoter at all. Although transient transfection assays show that overexpression of MRFs markedly induce the ACC β promoter activities, the endogenous ACC β expression is not observed in myoblast cells endogenously expressing high levels of MyoD, until myoblasts differentiate into myotubes. These facts indicate the presence of the accessory factors involved in regulating MRF-mediated activations of muscle genes and suggest that retinoid receptors might probably have the roles, in part, as one of these accessory factors. Moreover, it has been reported that retinoic acid receptors and basic helix-loop-helix proteins directly interact each other and synergistically affect muscle target genes²². In general, the regulations of eukaryotic transcription occur by forming the complexes of several transcription activators and repressors on the promoter.

MRF-mediated activation of ACC β promoter is controlled by two important *cis*-elements, E-box region (-493/-403) and noble sequence (+17/+24). MRFs bound to these elements have different quaternary structures. Heterodimers of MRF and E-protein is known to generally bind to E-box region, whereas the novel *cis*-element of ACC β promoter has strong affinity to homodimers of MRFs and its MRF-binding is suppressed by heterodimer formation with E-protein¹⁹. To exclude the effects of other several transcription factors bound to ACC β promoter and to check the effects of retinoid receptors on the activities of MRFs containing the different quaternary structures, the transient transfection assays were performed using the heterologous promoters containing E-box region of ACC β or six copies of novel *cis*-element at upstream of tk minimal promoter. These attempts showed that RXR α enhances MRF4 action on E-box regions but not on noble *cis*-element. RAR α suppresses the activities of MyoD acting on noble *cis*-element but not on E-box regions (Fig. 4). These data suggest that RXR α preferentially enhances the activities of heterodimer of MRF4 and E-protein, and RAR α is likely to inhibit the activities of MyoD homodimer.

In summary, the data in present study conclude that RAR α suppressed the transcriptional activities of MyoD homodimer in myoblast, and RXR α elevated the actions of MRF4-E-protein heterodimer in terminally differentiated myotubes.

V. Conclusion

We have shown that retinoic acids differentially regulated ACC β expression mediated by MRFs during myogenesis. RAR α suppressed MyoD activities in myoblasts and RXR α significantly enhanced ACC β promoter activity induced by MRF4 in myotubes. From the above results, retinoic acids control proper ACC β expression through the interaction with MRFs during myogenesis.

Reference

1. Lopez-casillas F, Bai DH, Luo X, Kong IS, Hermodson MA, Kim KH. Structure of the coding sequence and primary amino acid sequence of acetyl-coenzyme A carboxylase. Proc. Natl. Acad. Sci. USA 1988;85: 5784~5788.
2. Thampy KG. Formation of malonyl Coenzyme A in rat heart. J. Biol. Chem 1989;264:17631~17634.
3. Bianchi A, Evans JL, Iverson AJ, Nordlund AC, Watts TD, Witters LA. Identification of an isozymic form of acetyl-CoA carboxylase. J. Biol. Chem 1990;265:1502~1509.
4. Cook GA. Differences in the sensitivity of carnitine palmitoyltransferase to inhibition by malonyl-CoA are due to difference in K_i values. J. Biol. Chem 1984;259:12030~12033.
5. Winz R, Hess D, Aebersold R, brownsey RW. Unique structural features and differential phosphorylation of the 280-kDa component (Isozyme) of rat liver acetyl-CoA carboxylase. J. Biol. Chem 1994; 269:14438~14445.
6. McGarry JD, Woeltje KF, Kuwajima M, Foster DW. Regulation of ketogenesis and the renaissance of carnitine palmitoyltransferase. Diabetes. Metab. Rev 1989;5:271~284.
7. Davis RL, Weintraub H, Lassar AB. Expression of a single transfected cDNA converts fibroblasts to myoblasts. Cell 1987;51:987~1000.
8. Wright WE, Sassoon DA, Lin VK. Myogenin, a factor regulating

- myogenesis, has a domain homologous to MyoD. Cell 1989;56: 607~617.
9. Braun T, Buschhausen-Denker G, Bober E, Tannich E, Arnold HH. A novel human muscle factor related to but distinct from MyoD1 induces myogenic conversion in 10T1/2 fibroblasts. EMBO. J 1989;8:701~709.
 10. Braun T, Bober E, Winter B, Rosenthal N, Arnold HH. Myf-6, a new member of the human gene family of myogenic determination factors: evidence for a gene cluster on chromosome 12. EMBO.J 1990;9:821~831.
 11. Jeffrey H, Wold M, Wold B. herculin, a fourth member of the MyoD family of myogenic regulatory genes. Proc. Natl. Acad. Sci. USA 1990; 87:1089~1093.
 12. Lassar AB, Buskin JN, Lockshon D, Davis RL, Apone S, Hauschka SD et al. MyoD is a sequence-specific DNA binding protein requiring a region of *myc* homology to bind to the muscle creatine kinase enhancer. Cell 1989;58:823~831.
 13. Murre C, McCaw PS, Vaessin H, Caudy M, Jan LY, Jan YN et al. Interactions between heterologous Helix-Loop-Helix proteins generate complexes that bind specifically to a common DNA sequence. Cell 1989;58:537~544.
 14. Lassar AB, Davis RL, Wright WE, Kadesch T, Murre C, Voronova A et al. Functional activity of myogenic HLH proteins requires hetero-oligomerization with E12/E47-like proteins in vitro. Cell 1991;66: 305~315.
 15. Ott MO, Bober E, Lyons G, Arnold H, Buckingham M. Early expression of

- the myogenic regulatory gene, myf-5, in precursor cells of skeletal muscle in the mouse embryo. *Development* 1991;111:1097~1107.
16. Smith TH, Kachinsky AM, Miller JB. Somite subdomains, muscle cell origins, and the four muscle regulatory factor proteins. *J. Cell. Biol* 1994;127:95~105.
 17. Megency LA, Rudnicki MA. Determination versus differentiation and the MyoD family of transcription factors. *Biochem. Cell. Biol* 1995;73: 723~732.
 18. Bober E, Lyons GE, Braun T, Cossu G, Buckingham M, Arnold HH. The muscle regulatory gene, Myf-6, has a biphasic pattern of expression during early mouse development. *J. Cell. Biol* 1991;113: 1255~1265.
 19. Lee JJ, Moon YA, Ha JH, Yoon DJ, Ahn YH, Kim KS. Cloning of human acetyl-CoA carboxylase β promoter and its regulation by muscle regulatory factor. *J. Biol. Chem* 2001;276: 2576~2585.
 20. Lee JK, Kim KH. Roles of acetyl-CoA carboxylase β in muscle cell differentiation and in mitochondrial fatty acid oxidation. *Biochemical and Biophysical Research Communication* 1999;254: 657~660.
 21. Solanes G, Pedraza N, Iglesias R, Giralt M, Villarroya F. The human uncoupling protein-3 gene promoter requires MyoD and is induced by retinoic acid in muscle cells. *FASEB. J* 2000;14:2141~2143.
 22. Froeschle A, Alric S, Kitzmann M, Carnac G, Aurade F, Rochette-Egly C et al. Retinoic acid receptors and muscle b-HLH proteins: partners in

- retinoid-induced myogenesis. *Oncogene* 1998;16:3369~3378.
23. De Luca LM. Retinoids and their receptors in differentiation, embryogenesis and neoplasia. *FASEB. J* 1991;5:2924~2933.
 24. Giguere V. Retinoic acid receptors and cellular retinoid binding proteins: complex interplay in retinoid signaling. *Endocrine review* 1994;15:61~79.
 25. Mangelsdorf D, Evans RM. The RXR heterodimers and orphan receptors. *Cell* 1995;83:841~850.
 26. Leid M, Kastner P, Lyons R, Nakshatri H, Saunders M, Zacharewski T et al. Purification, cloning and RXR identity of the HeLa cell factor with which RAR or TR heterodimerizes to bind target sequences efficiently. *Cell* 1992;68:377~395.
 27. Giguere V, Ong ES, Segui P, Evans RM. Identification of a receptor for the morphogen retinoic acid. *Nature* 1987;330:624~629.
 28. Petkovich M, Brand NJ, Krust A, Chambon P. A human retinoic acid receptor which belongs to the family of nuclear receptors. *Nature* 1987; 330:444~450.
 29. Heyman RA, Mangelsdorf DJ, Dyck JA, Stein RB, Eichele G, Evans RM et al. *9-cis* retinoic acid is a high affinity ligand for the retinoid X receptor. *Cell* 1992;68:397~406.
 30. Levin AA, Sturzenbecker LJ, Kazmer S, Bosakowski T, Huselton C, Allenby G et al. *9-cis* retinoic acid stereoisomer binds and activates the nuclear receptor RXR α . *Nature* 1992;355:359~361.

31. Rudnicki MA, Schnegelsberg PNJ, Stead RH, Braun T, Arnold HH, Jaenisch R. MyoD or Myf-5 is required for formation of skeletal muscle. *Cell* 1993;75:1351~1359.
32. Choi J, Costa ML, Mermelstein CS, Chagas C, Holtzer S, Holtzer H. MyoD converts primary dermal fibroblasts, chondroblasts, smooth muscle, and retinal pigmented epithelial cells into striated mononucleated myoblasts and multinucleated myotubes. *Proc. Natl. Acad. Sci. USA* 1990;87:7988~7992.
33. Atchley WR, Fitch WM, Bronner-Fraser M. Molecular evolution of the MyoD family of transcription factors. *Proc. Natl. Acad. Sci. USA* 1994; 91:11522~11526.
34. Downes M, Mynett-Johnson L, Muscat GEO. The retinoic acid and retinoid X receptors are differentially expressed during myoblast differentiation. *Endocrinology* 1994;134:2658~2661.
35. Hinterberger TJ, Sassoon DA, Rhodes S, Konieczny S. Expression of the muscle regulatory factor MRF4 during somite and skeletal myofiber development. *Developmental biology* 1991;147:144~156.

MyoD, MRF4

acetyl-CoA carboxylase

RAR RXR

< >

가 acetyl-CoA caboxylase (ACC) 가
 . 265kDa α
 , 275kDa β
 . ACC β malonyl-CoA가
 rate-limiting carnitine palmitoyl-CoA transferase-1(CPT-1)
 . MyoD
 가 ACC β promoter *cis*-element ACC β
 .
 ACC β retinoic acid ACC β retinoic acid
 . H9C2 myotube
 ACC β promoter retinoic acid , H9C2 myoblast transfection , ACC β
 RXR α , RAR α
 . 가 NIH3T3 fibroblast
 transfection , retinoid receptor ACC β promoter
 , ACC β promoter retinoid receptor가
 consensus sequence가 , Myoblast
 myotube retinoid ACC β retinoid

