

Inhibition by clonidine of the carbachol-
induced tension development and
nonselective cationic current in guinea
pig ileal smooth muscle

Dissertation by
SeungSoo Chung

Brain Korea 21 Project for Medical Sciences
The Graduate School of Yonsei University

Inhibition by clonidine of the carbachol-
induced tension development and
nonselective cationic current in guinea pig
ileal smooth muscle

directed by Professor TaickSang Nam

A Dissertation submitted to the Faculty of
the Graduate School of Yonsei University

December, 2001

By

SeungSoo Chung

Brain Korea 21 Project for Medical Sciences
The Graduate School of Yonsei University

A Dissertation for the Degree of Dr. of Philosophy
in Medical Sciences by SeungSoo Chung
has been approved by

(Supervisory committee, chairman)

Supervisory committee

Supervisory committee

Supervisory committee

Supervisory committee

Brain Korea 21 Project for Medical Sciences
The Graduate School of Yonsei University

December, 2001

ACKNOWLEDGEMENTS

First and foremost, I would like to express my gratitude and appreciation to Professor TaickSang Nam, my advisor, for his support, guidance and encouragement from the beginning of my graduate life here to the writing of my dissertation at the very end. His kindness and counsel have been overwhelming.

I also would like to thank assistant Professor SeongChun Kwon for his hearted aid and confidence throughout the experiemnt. And many appreciative thanks to Professor DuckSun Ahn, who has never failed to help me, without ever complaining even when I had gotten into trouble.

I thank my parents, TaeMo Chung and KyungIm Ko for their love to attain my goals.

And finally, I thank my wife, JeongAh Yoon and son SeongEun Chung for providing me the inspiration to finish, with an endless love and understanding though we have been apart.

Contents

ABSTRACT	1
I. INTRODUCTION	2
II. MATERIALS AND METHODS	4
1. Measurement of muscle tension	4
2. Cell isolation	4
3. Whole-cell voltage clamp	6
4. Solutions	6
5. Solution application	7
6. Data analysis	7
7. Drugs	7
III. RESULTS	8
1. Changes in muscle force	8
2. Carbachol-activated current (I_{CCh}) in ileal myocytes	9
3. Effect of clonidine on I_{CCh}	10
4. Voltage dependence of the clonidine effect on I_{CCh}	11
5. Influence of adrenergic and imidazoline receptor antagonists on the inhibitory effect of clonidine	11
6. The effect of clonidine on GTP γ S-activated current	13
7. Effect of clonidine on voltage-dependent Ca^{2+} channels and K^{+} channels	15
IV. DISCUSSION	15
V. CONCLUSION	18
REFERENCE	19
ABSTRACT (IN KOREAN)	22

List of Figures

Fig. 1.	Chemical structure of clonidine hydrochloride	2
Fig. 2.	The schematic diagram showing the equipment of isometric tension measurement and whole cell voltage clamp system	5
Fig. 3.	Effects of cumulative application of clonidine on muscle force of guinea pig ileal longitudinal muscle	8
Fig. 4.	Effects of clonidine on carbachol-activated cationic current (I_{CCh}) in single guinea pig ileal smooth muscle cells	9
Fig. 5.	Effects of clonidine in the pipette solution on I_{CCh} in single guinea pig ileal smooth muscle cells	10
Fig. 6.	Voltage-dependency of the effect of clonidine on I_{CCh}	11
Fig. 7.	Yohimbine, KU14R and BU224 effects on the suppressive effect of clonidine on I_{CCh}	12
Fig. 8.	Effects of clonidine on GTP γ S-induced current	13
Fig. 9.	Effects of clonidine on voltage activated Ba^{2+} current through Ca^{2+} channel and K^{+} currents in single ileal smooth cells of guinea pig	14

Abstract

Inhibition by clonidine of the carbachol-induced tension development and nonselective cationic current in guinea pig ileal myocytes

SeungSoo Chung

*Brain Korea 21 Project for Medical Sciences
The Graduate School, Yonsei University*

(Directed by Professor TaickSang Nam)

Effects of clonidine, an imidazoline derivative as well as α_2 -adrenoceptor agonist, on carbachol (CCh)-evoked contraction in guinea pig ileal smooth muscle were studied using isometric tension recording.

To investigate the cellular mechanisms of the inhibitory effect of clonidine, its effects on CCh-evoked nonselective cationic current (I_{CCh}), voltage-dependent Ca^{2+} current (I_{Ca}) and voltage-dependent K^+ current (I_K) was also studied using patch-clamp recording techniques in single ileal cells.

Clonidine inhibited the contraction evoked by CCh ($1\ \mu M$) in a concentration-dependent manner with an IC_{50} value of $61.7 \pm 2.5\ \mu M$. High K^+ ($40\ mM$)-evoked contraction was only slightly inhibited even when clonidine was used at $300\ \mu M$. Externally applied clonidine inhibited I_{CCh} dose-dependently with an IC_{50} of $42.0 \pm 2.6\ \mu M$. When applied internally via patch pipettes, clonidine was without effect. An I_{CCh} like current induced by GTP γS was also inhibited by bath application of clonidine. None of KU14R and BU224, both, imidazoline receptor blockers, and yohimbine, an α_2 -adrenergic blocker, significantly affects the inhibitory effect of clonidine on I_{CCh} . Clonidine ($300\ \mu M$) only slightly decreased membrane currents flowing through voltage-gated Ca^{2+} channels or K^+ channels.

These data indicate that clonidine relaxes smooth muscle contraction produced by muscarinic receptor activation and suggest that the effect of clonidine seems due mainly to inhibition of I_{CCh} via acting directly on the involved cationic channel.

Key Words: gastrointestinal smooth muscle, carbachol, cationic current, clonidine, imidazoline

Inhibition by clonidine of the carbachol-induced tension development and nonselective cationic current in guinea pig ileal smooth muscle

SeungSoo Chung

Brain Korea 21 Project for Medical Sciences
The Graduate School, Yonsei University

<Directed by Professor TaickSang Nam>

I. INTRODUCTION

Clonidine has been known as a centrally acting α_2 -adrenergic receptor agonist that is used in clinical practice for its antihypertensive effect¹ (Fig 1). However, its practical use for treatment of hypertension was limited because of its unpleasant side effects such as sedation and dry mouth. Today there is resurgence of interest in centrally acting agents because clonidine have repeatedly been shown to bind to distinct nonadrenergic sites with nanomolar affinities^{2,3} and its anti-hypertensive effect was mediated by a novel receptor type specific for imidazolines and its side effect was mediated by α_2 -adrenergic receptor. These sites are classified into the imidazoline I₁ and the imidazoline I₂ sites; the I₁ sites are associated with control blood pressure and exhibit




Fig. 1. Chemical structure of clonidine hydrochloride.

high affinity for clonidine and its derivatives.^{1,4} Imidazoline I₂ receptors show a higher affinity for idazoxan and related imidazolines and are involved in eating behavior and psychiatric disorder such as major depression.^{5,6} Meanwhile, others reported that some atypical imidazoline receptors, whose pharmacologic properties do not correspond to any of the foregoing subtypes, are involved in insulin secretion and were recently called imidazoline I₃ receptors,⁷ and another type of not-yet-classified imidazoline binding site with quite different pharmacological characteristics is quite different are present in various tissues,⁸ but its physiological roles are far less known.

Binding experiments with [³H]clonidine and [³H]idazoxan in membranes from guinea pig,⁹ rabbit¹⁰ and rat gastric tissues¹¹ provided evidence suggesting that nonadrenergic binding sites for imidazolines are also present in the gastrointestinal tissue. Felsen *et al*¹² showed that the clonidine-displacing substance (CDS), an endogenous ligand acting at the imidazoline binding sites, induced a concentration-dependent contraction of rat gastric fundus strips, but another group reported that these sites are neither directly related to a postsynaptic contractile effect on rat gastric smooth muscle nor to acid release from isolated gastric glands.¹³ Thus, although the effect of imidazoline derivatives on gastrointestinal smooth muscle contractility is still controversial, it is possible that the inhibitory effect of clonidine on intestinal motility is mediated by imidazoline binding sites and not by α_2 -adrenoreceptors. Recent studies have shown that imidazoline derivatives block ATP-sensitive K⁺ channels to stimulate insulin release in pancreatic β -cells and rat insulinoma (RIN) cells.^{14,15} Imidazoline derivatives also inhibit the acetylcholine (ACh)-induced secretion of catecholamines in adrenal chromaffin cells¹⁶ by blocking nicotinic ACh receptors.¹⁷ Finally, these compounds also interact with 5-HT₃ receptors in NIE-115 cells, inhibiting the veratridine-induced influx of guanidinium.¹¹ Thus, the mechanism of the imidazoline effect may be mediated by modulating various ion channels. In mammalian gastrointestinal myocytes, stimulation of muscarinic receptors by ACh or carbachol (CCh) induces nonselective cation current (I_{CCh}).¹⁸ This current acts to depolarize the cell membrane, causing Ca²⁺ influx *via* voltage-gated Ca²⁺ channels. Thus I_{CCh} plays a role in the muscarinic receptor-mediated contraction of intestinal

smooth muscle. Under the circumstances, it is possible that clonidine, an imidazoline derivative, may influence the intestinal contractility by modifying the activity of various channels including the muscarinic receptor cationic channel, voltage-dependent calcium channel and voltage-dependent potassium channel.

The present study was therefore, designed (1) to assess the effect of clonidine, the prototypical imidazoline derivative, on high K^+ solution-induced and CCh-induced contractions, and (2) to examine its effect on I_{CCh} , voltage-dependent Ca^{2+} current (I_{Ca}) and voltage-dependent K^+ current (I_K).

II. MATERIALS AND METHODS

1. Measurement of muscle tension

Guinea-pigs of either sex, weighing 300~350 g, were exsanguinated after stunning. The ileum was isolated and cut into segments of 3- to 4-cm lengths and placed in physiological salt solution (PSS). As shown in Fig. 2A, the longitudinal muscle layer of intestinal segments was peeled from the underlying circular muscle and washed in PSS. Each ileal longitudinal muscle strip was attached to a glass holder and mounted vertically in a 20-ml organ bath filled with Krebs-Henseleit (K-H) solution. K-H solution was saturated with a mixture of 95% O_2 and 5% CO_2 at 37°C to maintain pH at 7.4. Muscle strips, loaded with a resting tension of 0.5 g, were equilibrated for more than 60 min before starting the experiment. Changes in contractile tension were measured isometrically with a force-displacement transducer and recorded on a pen-recorder. The muscle strip was repeatedly stimulated by isotonic 40 mM KCl until the response became stable and then the experiment was started.

2. Cell isolation

The longitudinal muscle layer of ileum was cut into small pieces and placed in Ca^{2+} -free PSS. The Ca^{2+} -free PSS was then replaced with PSS containing 30 μM Ca^{2+} (low Ca^{2+} PSS); and




Fig. 2. The schematic diagram showing the equipment of isometric tension measurement (A) and whole cell voltage clamp system (B).

30-min incubations at 37°C were carried out in fresh low Ca²⁺ PSS containing collagenase (0.3 mg/ml), papain (0.6 mg/ml), and bovine serum albumin (1 mg/ml). After this enzyme digestion, tissue fragments were suspended in a fresh 120 μM Ca²⁺ containing PSS and gently agitated. The resulting suspension was centrifuged at 600 xg for 2 min, and the cells were resuspended in a 0.5 mM Ca²⁺ containing PSS, aliquoted into 12 mm poly-L-lysine-coated cover glasses and stored in a humidified atmosphere at 4°C. Experiments were carried out within 12 hrs of harvesting (22 ~ 24°C).

3. Whole-cell voltage clamp

Whole-cell membrane currents were recorded at room temperature using standard patch-clamp techniques (Fig. 2B). The patch pipette had a resistance of 3~6 M Ω when filled with a pipette solution. Membrane currents were measured with an Axoclamp 200 A voltage-clamp amplifier (Axon Instrument, Foster City, CA, USA). Command pulses were applied using an IBM-compatible computer and pCLAMP (version 6.0) software. The data were filtered at 5 kHz and displayed on an oscilloscope, a computer monitor, and a pen recorder.

4. Solutions

The K-H solution was of the following composition: 136.9 mM NaCl, 5.4 mM KCl, 1.5 mM CaCl₂, 1 mM MgCl₂, 23.8 mM NaHCO₃, 0.01 mM ethylenediamine tetraacetic acid (EDTA) and 5.5 mM glucose. Substituting NaCl with equimolar KCl in K-H solution made isotonic high K⁺ (40 mM) solutions. The PSS used for cell isolation and the bathing solution for potassium current recording had the following composition: 126 mM NaCl, 6 mM KCl, 2 mM CaCl₂, 1.2 mM MgCl₂, 14 mM glucose, 10.5 mM N-[2-hydroxyethyl]piperazine-N-[2-ethansulphonic acid] (HEPES) (titrated to pH 7.4 with NaOH). The patch pipette solution for the K⁺ current recording had the following composition (mM): 134 mM KCl, 1.2 mM MgCl₂, 1 mM MgATP, 0.1 mM Na₂GTP, 0.05 mM ethylene glycol-bis (β -aminoethylether)-N,N,N',N'-tetraacetic acid (EGTA), 14 mM glucose and 10.5 mM HEPES (titrated to pH 7.2 with NaOH). The bathing solution for recording the Ba²⁺ current through Ca²⁺ channels had the following composition; 126 mM NaCl, 6 mM TEACl, 2 mM BaCl₂, 1.2 mM MgCl₂, 14 mM Glucose, 10.5 mM HEPES (titrated to pH 7.2 with NaOH). The pipette solution for recording the Ba²⁺ current through Ca²⁺ channels had the following composition: 134 mM CsCl, 1.2 mM MgCl₂, 1 mM MgATP, 0.1 mM Na₂GTP, 0.05 mM EGTA, 14 mM glucose and 10.5 mM HEPES (titrated to pH 7.2 with CsOH). The bath solution for recording I_{CC_h} had the following composition; 120 mM CsCl, 12 mM glucose, 10 mM HEPES (titrated to pH 7.4 with CsOH) (total Cs⁺ = 124 mM). The pipette solution for recording I_{CC_h} had

the following composition; 80 mM CsCl, 1 mM MgATP, 5 mM creatine, 1 mM Na₂GTP, 10 mM BAPTA, 20 mM glucose and 10 mM HEPES, 4.6 mM CaCl₂ (calculated [Ca²⁺]_i=100 nM (titrated to pH 7.2 with CsOH) (total Cs⁺ = 124 mM). The presence of 1 mM GTP in this solution reduced desensitization to a minimum.¹⁹ In experiments designed to activate cationic channels, without muscarinic receptor activation, GTP in the pipette solution was replaced with 200 μM GTP γS.^{19,20}

5. Solution application

Test solutions were applied to single ileal cells via a gravity-fed fused silica capillary tube connected to an array of six polyethylene tubes. The outlet of the perfusion system was located within 100 mm of the cell. Drug application was started by switching the control external solution to a drug solution. Complete solution exchange occurred within 1 sec.

6. Data analysis

Concentration-effect curves were fitted by a logistic function in the following form:

$$\frac{I_{\text{clonidine}}}{I_{\text{control}}} = \frac{1}{1 + ([\text{clonidine}]/IC_{50})^p}$$

where I_{control} and $I_{\text{clonidine}}$ are the cationic current amplitudes in the absence and presence of various clonidine concentrations, [clonidine], respectively; IC_{50} equals the clonidine concentration at which current amplitude was reduced by 50% and p is the slope factor of the inhibition curve. The data was analyzed and plotted using GraphPad Prism software version 3.0 (San Diego, CA, USA). Values are given as the means ± S.E.M.

7. Drugs

The following drugs and chemicals were used: collagenase (type 1A), adenosine 5'- triphosphate (ATP, magnesium salt), guanosine 5'-triphosphate (GTP, sodium salt), clonidine hydrochloride,

carbamylcholine chloride (carbachol), yohimbine, guanosine 5'-O-(3-thiotriphosphate) (GTP γ S, tetralithium salt), creatine, HEPES, 1,2-bis(2-aminophenoxy) ethane-N,N,N',N'-tetraacetic acid (BAPTA), EGTA were obtained from Sigma Chemical (St. Louis, MO, USA). 2-(2-Ethyl-2,3-dihydro-2-benzofuranyl)-1H-imidazole (KU14R) and 2-(4,5-Dihydroimidazol-2-yl)quinoline (BU224) were from Tocris (Bristol, UK).

III. RESULTS

1. Changes in muscle force

Fig. 3 shows the inhibitory effect of cumulative application of clonidine on the sustained




Fig. 3. Effects of cumulative application of clonidine on muscle force of guinea pig ileal longitudinal muscle. (A) Typical tracing of effects of clonidine on muscle force evoked with 1 μ M carbachol and 40 mM KCl. (B) Concentration-response curves of clonidine effect on muscle force of guinea pig ileum. 100% represents muscle force induced by 1 μ M carbachol and 40 mM KCl before the cumulative addition of clonidine.

contraction induced by high K^+ (40 mM) solution and CCh (1 μ M) in guinea-pig ileal segment.

The CCh-induced contraction was more sensitive to clonidine than the high K^+ -induced contraction. IC_{50} of clonidine on the CCh-induced contraction was $61.7 \pm 2.5 \mu$ M (n=8). Clonidine decreased the high K^+ -induced contraction only by 20.7% even when applied at 300 μ M.

2. Carbachol-activated current (I_{CCh}) in ileal myocytes

In cells voltage-clamped at -40 mV, inward cationic current was evoked by bath application of CCh after at least 3 min elapsed from break-through of the patch membrane. CCh was applied at 50 μ M, a concentration close to the maximally effective in these cells.²¹ The current-voltage (I-V) relationship was measured with a negative going ramp pulse from 80 mV to -120 mV over 6s (Fig. 4A, control trace). The I-V relationship of I_{CCh} showed inward rectification at around $+15$ mV and a negative slope conductance at negative potential range (<40 mV, Fig. 4A, control trace); that is, it has a typical U-shaped dependence on the membrane potential. All these properties are similar to those of the muscarinic receptor operated nonselective cationic channels.^{21,22}




Fig. 4. Effects of clonidine on carbachol-activated cationic current (I_{CCh}) in single guinea pig ileal smooth muscle cells. (A) Steady-state current-voltage relationships measured by 6s duration voltage ramps in the presence of 50 μ M carbachol and clonidine at the indicated concentration. (B) Concentration-response curve of clonidine on I_{CCh} .

3. Effect of clonidine on I_{CCh}

I_{CCh} evoked by $50 \mu\text{M}$ CCh was suppressed by clonidine externally and cumulatively applied at ascending concentrations of $10 \sim 300 \mu\text{M}$ (Fig. 4A). The current suppression increased with increasing clonidine concentration and reversed nearly completely within about 1 min when clonidine was washed away. The dose-dependency of the inhibitory effect of clonidine is shown in Fig. 4B. The degree of inhibition was estimated as the ratio of remaining current to the control current. A mean value of IC_{50} was $42.0 \pm 2.6 \mu\text{M}$ ($n=4$).

Fig. 5A shows a representative of experiments designed to see if clonidine can act from the inside of the cell to block the I_{CCh} . Clonidine ($300 \mu\text{M}$) was applied internally via patch pipettes. I_{CCh} amplitudes, expressed per the membrane capacitance, were $38.0 \pm 3.8 \text{ pA/pF}$ ($n=4$) in the absence and $37.8 \pm 3.5 \text{ pA/pF}$ ($n=4$) in the presence of intracellular clonidine. There was no significant difference between the mean values (Fig. 5B). When clonidine was externally applied during generation of I_{CCh} in cells internally treated with clonidine, the I_{CCh} was profoundly inhibited




Fig. 5. Effects of clonidine in the pipette solution on I_{CCh} in single guinea pig ileal smooth muscle cells. Clonidine was added at $300 \mu\text{M}$, the maximally suppressing dose when externally applied, in the pipette solution. A) Typical tracing of effects of clonidine on I_{CCh} in single guinea pig ileal smooth muscle cells in the presence of clonidine ($300 \mu\text{M}$) in the pipette solution. B) Comparison of amplitude of I_{CCh} between absence and presence of clonidine ($300 \mu\text{M}$) in the pipette solution. Inward current induced by $50 \mu\text{M}$ carbachol were normalized to the membrane capacitance measured (mean \pm S.E.M., $n=4$).

(Fig. 5A). Therefore, clonidine seems likely to act only from the outside of the cell to block I_{CCh} .

4. Voltage dependence of the clonidine effect on I_{CCh}

Steady-state I-V relationships in the presence of $50\ \mu\text{M}$ CCh and clonidine ($100\ \mu\text{M}$) were measured using the same ramp pulse as described in Fig. 6. I_{CCh} was partially suppressed by $100\ \mu\text{M}$ of clonidine. The ratio of I_{CCh} amplitude in the presence of clonidine to that in its absence was plotted against membrane potential ($n=3$) (Fig. 6). The degree of I_{CCh} inhibition by the clonidine decreased with membrane depolarization; the degree of I_{CCh} inhibition at $-120\ \text{mV}$ was 2.1 ± 0.3 ($n=3$) times greater than that at $-10\ \text{mV}$.

5. Influence of adrenergic and imidazoline receptor antagonists on the inhibitory effect of clonidine

Clonidine ($100\ \mu\text{M}$) suppressed I_{CCh} evoked at $-40\ \text{mV}$ to $68.5 \pm 1.7\%$ ($n=4$). A 15-min pre-treatment with yohimbine ($1\ \mu\text{M}$), an α_2 -adrenoceptor antagonist, did not influence the inhibition by clonidine ($66.3 \pm 1.9\%$ suppression, $n=4$) (Fig. 7A). Substantially similar result was obtained with




Fig. 6. Voltage-dependency of the effect of clonidine on I_{CCh} . Stimulation protocol is the same with Fig. 4. Steady-state current-voltage relationships measured by 6s duration voltage ramps in the presence of $50\ \mu\text{M}$ carbachol and clonidine ($100\ \mu\text{M}$). I_{CCh} was partially suppressed by $100\ \mu\text{M}$ of clonidine. The ratios of amplitude of I_{CCh} applied with clonidine to those of control currents were plotted against membrane potential ($n=3$).




Fig. 7. Yohimbine (A), KU14R (B) and BU224 (C) effects on the suppressive effect of clonidine on I_{CCh} . The preparations were pretreated with yohimbine, KU14R and BU224 for 15min and I_{CCh} was recorded at -40 mV holding potential. All of the antagonists did not block the clonidine effect on I_{CCh} .

KU14R (10 μ M), an imidazoline I_1 receptor antagonist ($68.2 \pm 2.4\%$ suppression, $n=4$) (Fig. 7B) or BU224 (10 μ M), an imidazoline I_2 receptor antagonist ($62.1 \pm 2.1\%$ suppression, $n=5$) (Fig. 7C).

These suggest that clonidine may cause I_{CCh} inhibition by acting on some sites other than imidazline I_1 receptors, I_2 receptors and α_2 -adrenoceptors.

6. The effect of clonidine on GTP γ S-activated current

It is known that muscarinic stimulation activates non-selective cation channels through the activation of GTP-binding protein-mediated signal transduction.²² As shown in Fig. 8B, in cells held at -40 mV, intracellular application of $200\text{ }\mu\text{M}$ GTP γ S *via* patch pipettes produced inward cationic current, which developed slowly to reach a steady-state level within $3\sim 5$ min and then sustained for several 10 min as previously reported.¹⁹ The GTP γ S-activated current showed




Fig. 8. Effects of clonidine on GTP γ S-induced current. A) GTP γ S (0.2 mM) was added in the pipette solution. Slowly increasing inward current was induced 2 min after break through (arrowhead). B) This current- voltage (I-V) relation curve was recorded with the same method as Fig. 4. This I-V curve are similar to those induced by carbachol (see Fig. 4, control trace).




Fig. 9. Effects of clonidine on voltage activated Ba^{2+} current through Ca^{2+} channel and K^{+} currents in single ileal smooth cells of guinea pig. A, 1) Ba^{2+} current was elicited by 200 ms depolarization from -60 mV to 0 mV at 0.1 Hz. Clonidine, $300 \mu\text{M}$, the maximally I_{CCl} -suppressing dose when applied externally, was applied in the bath solution. A, 2) Current-voltage relationships for the peak inward Ba^{2+} current activated from a holding potential of -60 mV. The peak amplitude of Ba^{2+} current evoked by stepping to different potential from holding potential is plotted against the potential. B, 1) In each panel the cell was held at -60 mV, and test depolarization with duration 1 s were applied from -40 mV to $+80$ mV in 10 mV increments. Currents are recorded using a pipette containing 0.05 mM EGTA. Clonidine at $300 \mu\text{M}$, the maximally I_{CCl} -suppressing dose when applied externally, was applied in the bath solution. B, 2) Current-voltage relationships for the outward K^{+} current activated from a holding potential of -60 mV. The sustained amplitude of K^{+} current evoked by stepping to different potential from holding potential is plotted against the potential.

time- and voltage-dependent characteristics similar to those of I_{CCh} (Fig. 8B), and it was also inhibited by 100 μ M clonidine ($72.1 \pm 1.1\%$ suppression, $n=4$) (Fig. 8A). Therefore, it seems likely that the binding site for clonidine is the cationic channel rather than the muscarinic receptor.

7. Effect of clonidine on voltage-dependent Ca^{2+} channels and K^+ channels

The effects of clonidine on barium current (I_{Ba}) through voltage-dependent Ca^{2+} channels were investigated. I_{Ba} was evoked by stepping to various depolarizing levels from a holding potential of -60 mV. As shown in Fig. 9A, clonidine (300 μ M) produced only a small decrease in the amplitude of I_{Ba} ; the percent inhibition of the current activated at 0 mV was $11.6 \pm 2.6\%$ ($n=4$) on average.

As shown in Fig. 9B, clonidine (300 μ M) also slightly suppressed K^+ current (I_K) evoked by stepping to various potentials (up to 80 mV) from -60 mV. Actually, the percent inhibition of I_K evoked at 30 mV was $14.9 \pm 4.4\%$ ($n=4$).

IV. DISCUSSION

The actions of the imidazoline derivative clonidine in the guinea pig ileal longitudinal smooth muscle can be summarized as follows; 1) it inhibited CCh-evoked contraction in a dose-dependent manner. 2) The inhibition seemed to be brought about by suppression of I_{CCh} .

As mentioned in the introduction, it is reported that imidazoline receptors are located in various tissues and play a role according to their physiological circumstances. For example, the imidazoline receptors have been identified in several kinds of smooth muscle cells^{23,24} and suggested to contribute to regulation of muscle tone. In the gastrointestinal tract, radioligand binding data provide evidence that the rat and human stomachs are endowed with I_2 -like binding sites and non- I_{ij} non- I_2 -[3H]clonidine binding sites that represent two different entities, suggesting that they participate in the regulation of intestinal smooth muscle motility.¹⁶

In the present study, clonidine suppressed CCh-evoked contraction with the IC_{50} value of 61.7 μ M

range. The K_d of clonidine for α_2 -adrenoceptors is estimated to be in the nanomolar range (7.3 nM) in vascular smooth muscle.²⁵ These, therefore, suggested that the action of clonidine is not mediated by α_2 -adrenoceptors.

As mentioned in the introduction, imidazolines have been suggested to block various ion channels including the nicotinic acetylcholine receptor channel,¹⁷ the 5HT₃ receptor channel,²⁶ ATP-sensitive K⁺ channel,¹⁴ and the NMDA channel²⁷ with IC₅₀ of 10 micromolar range possibly by binding to a phencyclidine recognition site within their ion-conducting pore. Inhibition of single-channel conductance in cell-free patches by I₁-receptor ligands is highly suggestive of direct blockade.²⁸ Furthermore, imidazolines displace [³H] phencyclidine and [³H] MK801 from nicotinic ACh receptors and NMDA receptors, respectively, consistent with direct activation at the pore.¹⁷ Phentolamine, an imidazoline derivative, can block currents flowing via ATP-sensitive K⁺ channel expressed in *Xenopus* oocytes,²⁹ also indicating the direct action of imidazolines at the pore of this channel. In the gastrointestinal tract, the signal transduction pathway between muscarinic receptor activation and ion channels has been studied in guinea pig ileal myocytes.²² The muscarinic agonist ACh causes membrane depolarization of smooth muscle and thus an increase in Ca²⁺ influx via voltage-gated Ca²⁺ channels. The depolarizing action is mediated primarily by M₂ muscarinic receptor activation^{21,30} which is linked to the voltage-dependent, [Ca²⁺]_i-sensitive cationic channels via G_i/G_o proteins.³⁰ In the present study, we found for the first time, that clonidine suppressed the CCh-induced I_{CCh} with the IC₅₀ value of 42 μ M, similar to the corresponding values reported for various other types of ion channels (see above). This IC₅₀ was also well matched with that for clonidine effect on CCh-induced contraction. Substances affecting neurotransmitter-operated channels like this CCh-activated channel may act at any of the steps between ligand-receptor binding and channel gating. However, indirect interference of clonidine with activation of the G-protein by the receptor is not likely because: 1) clonidine suppressed the GTP γ S-activated current (Fig. 8) and 2) it failed to inhibit I_{CCh} when applied intracellularly (Fig. 5). 3) neither yohimbine (α_2 -adrenoceptor antagonist) nor KUI4R (imidazoline I₁ receptor

antagonist), blocked the clonidine effect on I_{CCh} (Fig. 7). 4) furthermore, clonidine inhibited the I_{CCh} voltage dependent manner such as 4-AP, quinidine and quinidine. So clonidine seems to act directly on the cation channel mainly from the external side of the membrane. This is contradictory to the previous report that imidazolines do not inhibit non-selective cation channels whose opening is mediated by G_i -protein-coupled α_{2A} -adrenoceptors in human erythroleukemia (HEL) cells.³¹ Therefore, imidazoline effect on the cationic channel might be tissue- and/or activated receptor-specific.

Could the clonidine-binding site on the muscarinic receptor-operated cationic channel actually be the imidazoline receptor? At first sight, it would seem not, as clonidine action on I_{CCh} was blocked neither by KU14R, a putative imidazoline I_1 -antagonist nor by BU224, I_2 -receptor antagonist, and the apparent affinities for clonidine for the I_{CCh} are several orders of magnitude lower than the affinity of these drugs for I_1 sites. However, radioligand binding experiments carried out in cell membrane from guinea pig small intestine also revealed the existence of another non-adrenergic [3H]clonidine binding site, namely, non- I_1 /non- I_2 -[3H]clonidine binding sites that are the different entities.¹³ The characteristics of these binding sites were not clarified. So, whether the clonidine-binding site responsible for I_{CCh} inhibition is an imidazoline receptor cannot be determined at present. Naturally, a connection between the clonidine binding site for I_{CCh} inhibition and the imidazoline receptor is purely speculative at this stage; to demonstrate this connection, further study needed.

Meanwhile, clonidine slightly suppressed the I_K (11.6% decrease at 30 mV) or I_{Ca} (14.9% decrease at 0 mV) in guinea pig ileal longitudinal smooth muscle at 300 μM . Needless to say, clonidine may inhibit the I_{Ca} more significantly at more than 300 μM . However, clonidine can inhibit I_{CCh} at much lower concentration ($IC_{50}=42.0\pm2.6 \mu M$). Thus the inhibitory effect of clonidine on carbachol-induced contraction of guinea pig ileal longitudinal tract is mainly mediated by suppression of I_{CCh} through involved cation channels. These results are not inconsistent with the previous report that the voltage-operated calcium channel is only slightly affected by most

imidazolines that are ligands for I_1 receptors.³¹

The voltage-dependence of I_{CCh} inhibition by clonidine resembled that reported for quinidine, TEA, 4-AP, caffeine and quinine in ileal myocytes.³² The I_{CCh} inhibition by these agents was strongly attenuated by membrane depolarization. clonidine have positive charge in the water. So It is possible that clonidine may pass partway through the field of the membrane. Such blockade tends to voltage-dependent because the interference would be relieved by depolarization due to the reduction of driving force.

V. CONCLUSION

Effects of clonidine, an imidazoline derivative as well as α_2 -adrenoceptor agonist, on carbachol (CCh)-evoked contraction in guinea pig ileal smooth muscle were studied using isometric tension recording. To investigate the cellular mechanisms of the inhibitory effect of clonidine, its effects on CCh-evoked nonselective cationic current (I_{CCh}), voltage-dependent Ca^{2+} current (I_{Ca}) and voltage-dependent K^+ current (I_K) was also studied using patch-clamp recording techniques in single ileal cells. The results of the experiments were as followings;

1. Clonidine inhibited the contraction evoked by carbachol ($1 \mu M$) in a concentration-dependent manner with an IC_{50} value of $61.7 \pm 2.5 \text{ mM}$. High K^+ (40 mM)-evoked contraction was only slightly inhibited even when $300 \mu M$ clonidine, effective enough to abolish carbachol-evoked contraction, was used.

2. Externally applied clonidine inhibited I_{CCh} dose-dependently with IC_{50} of $42.0 \pm 2.6 \mu M$, and in a reversible manner. When applied internally via patch pipettes, clonidine was without effect.

3. A I_{CCh} like current induced by GTP γS , a direct G protein activation, was also inhibited by bath application of clonidine.

4. None of KU14R, and BU224, both imidazoline receptor blockers, and yohimbine, an α_2 -adrenergic blocker, significantly affects the inhibitory effect of clonidine on I_{CCh} .

5. Clonidine ($300 \mu M$) only slightly decreased the amplitude of I_{Ca} (11.6% decrease at 0 mV)

or that of I_K (14.9% decrease at 30 mV).

In conclusion, clonidine relaxes the ileal smooth muscle precontracted by CCh, possibly by the inhibiting I_{CCh} , a depolarizing current, *via* a direct blocking action on the muscarinic receptor-operated cationic channel. This action may underlie the regulating effect on intestinal motility by imidazolines observed so far.

REFERENCES

1. Bousquet P, Feldman J, Schwartz J. Central cardiovascular effects of a adrenergic drugs: differences between catecholamines and imidazolines. *J Pharmacol Exp Ther* 1984;230:232-6.
2. Atlas D. Clonidine displacing substance (CDS) and its putative imidazoline receptor. *Biochem Pharmacol* 1991;41:1541-9.
3. Konno F, Kusunoki M, Takayanagi. Receptor interactions of a series of imidazolines: comparison of the alpha-2-adrenoceptors between the rabbit vas deferens and guinea pig ileum. *Jpn J Pharamcol* 1987;44:171-8.
4. Ernsberger P, Graves ME, Graff LM, Zakieh N, Nguyen P, Collins LA, Westbrooks KL, Johnson GG. I_1 -imidazoline receptors: definition, characterization, distribution, and transmembrane signaling. *Ann NY Acad Sci* 1995;763:22-42.
5. Garcia-sevilla JA, Escriba PV, Sastre M, Walzer C, Busquets X, Jaquet G, Reis D, Guimon J. Immunodetection and quantitation of imidazoline receptor proteins in platelets of patients with major depression and in brain of suicide victims. *Arch Gen Psychiatry* 1996;53:803-10.
6. Menargues AM, Ced M, Artiga O, Obach R, Garcia-sevilla JA. Modulation of food intake by α_2 -adrenoceptor antagonists and I_2 -imidazoline drugs in rats. LSL60101 as a novel and selective ligand for I_2 -imidazoline site (abstr). *Br J Pharmacol* 1994;111:298.
7. Chan SLF, Brown CA, Scarpello KE, Morgan NG. The imidazoline site involved in control of insulin secretion: characteristics that distinguish it from I_1 and I_2 sites. *Br J Pharmacol* 1994;112:1065-70.
8. Muramatsu I, Kigoshi S. Tizandine may discriminate between imidazoline-receptors and alpha2-adrenoceptors. *Jpn J Pharmacol* 1992;59:457-9.
9. Houi N, Kamasaki Y, Itoh T. Effects of histamine H_2 receptor antagonists on acid secretion stimulated by imidazoline derivatives in isolated parietal cells. *Eur J Pharmacol* 1987;144:67-76.
10. Tesson F, Limon I, Parini A. Tissue-specific location of mitochondrial imidazoline guanidinium receptive sites. *Eur J Pharmacol* 1992;219:335-8.

11. Molderings GJ, Ruppert K, Bonisch H, Göthert M. No relationship of I₁- and I₂-imidazoline binding sites to inhibiting effects of imidazolines on ligand-gated ion channels. *Ann NY Acad Sci* 1995;763:420-32.
12. Felsen D, Ernsberger P, Meeley MP, Reis DJ. Clonidine displacing substance is biologically active on smooth muscle. *Eur J Pharmacol* 1987;42:453-5.
13. Molderings GJ, Burian M, Menzel S, Donecker K, Homann J, Nilius M, Göthert M. Imidazoline recognition sites and stomach function. *Ann NY Acad Sci* 1999;881:332-43.
14. Dunne MJ. Block of ATP-regulated potassium channels by phentolamine and other α -adrenergic antagonists. *Br J Pharmacol* 1991;103:1847-50.
15. Jonas JC, Pant TD, Henguin JC. Imidazoline antagonists of α_2 -adrenoceptors increase insulin release in vitro by inhibiting ATP-sensitive K⁺ channels in pancreatic β -cells. *Br J Pharmacol* 1992;107:8-14.
16. Oharaimaizumi M, Kumakura K. Effects of imidazole compounds on catecholamine release in adrenal chromaffin cells. *Cell Mol Neurobiol* 1992;12:273-83.
17. Musgrave IF, Krautwurst D, Hescheler J, Schultz G. Clonidine and cirazoline inhibit activation of nicotinic channels in PC12 cells. *Ann NY Acad Sci* 1995;763:272-82.
18. Inoue R, Kitamura K, Kuriyama H. Acetylcholine activates single sodium channels in smooth muscle cells. *Pfügers Arch* 1985;410:67-74.
19. Zholos AV, Bolton TB. A novel GTP-dependent mechanism of ileal muscarinic metabotropic channel desensitization. *Br J Pharmacol* 1996;119:997-1012.
20. Zholos AV, Bolton TB. G-protein control of voltage dependence as well as gating of muscarinic metabotropic channels in guinea-pig ileum. *J Physiol* 1994;486:195-202.
21. Bolton TB, Zholos AV. Activation of M₂ muscarinic receptors in guinea-pig ileum opens cationic channels modulated by M₃ muscarinic receptors. *Life Sci* 1997;60:1121-8.
22. Inoue R, Isenberg G. Acetylcholine activates nonselective cation channels in guinea pig ileum through G protein. *Am J Physiol* 1990;258:C1173-8.
23. Regunathan S, Youngson C, Wang H, Reis D. Imidazoline receptors in vascular smooth muscle and endothelial cells. *Ann NY Acad Sci* 1995;763:580-90.
24. Yablonsky F, Rausse J. Non-adrenergic binding sites for the α_2 -antagonist [³H]idazoxan in the rabbit urethral smooth muscle. *Biochem Pharmacol Toxicol (Copenh)* 1991;50:266-71.
25. Weis R, Webb R, Smith C. α_2 adrenoceptors on arterial smooth muscle: selective labelling by [³H] clonidine. *J Pharmacol Exp Ther* 1983;225:599-605.
26. Molderings GJ, Schmidt K, Bonisch H, Göthert M. Inhibition of 5-HT₃ receptor function by imidazolines in mouse neuroblastoma cells: potential involvement of α_2 binding sites. *N-S Arch Pharmacol* 1996;354:245-52.
27. Olmos G, Ribera J, Garcia-sevilla JA. Imidazoli(di)ne compounds interact with the phencyclidine site of NMDA receptors in the rat brain. *Eur J Pharmacol* 1996;310:273-6.

28. Cull Candy SG, Mathie A, Powis DA. Acetylcholine receptor channels and their block by clonidine in cultured bovine chromaffin cells. *J Physiol* 1988;402:255-78.
29. Proks P, Ashcroft FM. Phentolamine block of K_{ATP} channels is mediated by Kir6.2. *Proc Natl Acad Sci USA* 1997;94:11716-20.
30. Wang YX, Fleischmann BK, Kotlikoff MI. M_2 receptor activation of non selective cation channels in smooth muscle cells. Calcium and G_i/G_o requirements. *Am J Physiol* 1997; 273:C500-C8.
31. Musgrave IF, Seifert R. α_{2A} -adrenoceptors mediate activation of non-selective cation channels via G_i -proteins in human erythroleukaemia (HEL) cells. No evidence for functional role of imidazoline receptors in modulating calcium. *Biochem Pharmacol* 1995;49:187-96.
32. Chen S, Inoue R, Ito Y. Pharmacological characterization of muscarinic receptor-activated cation channels in guinea-pig ileum. *Br J Pharmacol* 1993;109:793-80.

국문요약

기니 픽 회장 평활근에서 carbachol 유발성 장력과 비선택성 양이온 전류의 clonidine에 의한 억제

본 논문은 기니 픽 회장 평활근에서 carbachol에 의해 유발된 장력에 대하여, α_2 아드레날린성 수용체 효현제이자 imidazoline 유도체인 clonidine의 효과를 등척성 수축장력을 측정함으로써 알아보고자 하였다. 또 clonidine 효과의 세포 내 기전을 알아보기 위하여 분리한 단일 회장 평활근 세포에서 막전압 고정법을 사용하여 carbachol 유발성 비선택성 양이온 전류(I_{CCh}), 막전압 의존성 Ca^{2+} 전류(I_{Ca}), 막전압 의존성 K^+ 전류(I_K)에 대한 clonidine의 효과를 연구하였다.

Clonidine은 carbachol $1 \mu M$ 에 의해 유발된 수축을 농도 의존적으로 억제하였고 이에 대한 IC_{50} 은 $61.7 \pm 2.5 \mu M$ 이었다. 이에 반하여 고농도 K^+ (40 mM)에 의해 유발된 수축은 carbachol에 의해 유발된 수축에 비하여 clonidine의 수축억제효과가 현저히 적었다. 막전압 고정법을 이용한 실험에서 관류액에 clonidine을 포함시켜 세포에 투여한 결과 Clonidine은 I_{CCh} 을 농도 의존적으로 억제하였다($42.0 \pm 2.6 \mu M$). Clonidine을 patch pipette 속에 투여하여 세포안쪽에 작용하도록 하였을 경우 clonidine은 I_{CCh} 의 크기에 거의 영향을 미치지 못하였다. 뿐만 아니라 $GTP \gamma S$ 에 의해 유발된 비선택성 양이온 전류도 I_{CCh} 와 마찬가지로 clonidine에 의해 억제되었다. Imidazoline 수용체 길항제인 KU14R, BU224 그리고 α_2 아드레날린성 수용체 길항제인 yohimbine 모두가 clonidine의 I_{CCh} 억제작용에 아무 효과도 나타내지 못하였다. 한편, clonidine은 막전압 의존성 Ca^{2+} 전류나 막전압 의존성 K^+ 전류에 대해서는 큰 영향을 나타내지 못하였다.

위의 결과를 종합하여 볼 때 clonidine은 주로 carbachol 유발성 비선택성 양이온 통로에 직접 작용하여 그를 통한 전류를 억제함으로써 carbachol에 의해 유발된 수축을 억제하는 것으로 생각된다.

핵심되는 말: 장관 평활근, 회장, carbachol, 양이온 전류, clonidine, imidazoline