

WJG 20th Anniversary Special Issues (18): Pancreatitis**Balloon dilation itself may not be a major determinant of post-endoscopic retrograde cholangiopancreatography pancreatitis**

Sung Ill Jang, Gak Won Yun, Dong Ki Lee

Sung Ill Jang, Department of Internal Medicine, Kangnam Sacred Heart Hospital, Hallym University College of Medicine, Seoul 150-950, South Korea

Gak Won Yun, Dong Ki Lee, Department of Internal Medicine, Gangnam Severance Hospital, Yonsei University College of Medicine, Seoul 135-720, South Korea

Author contributions: Jang SI reviewed the literature, collected data, wrote and prepared the first draft of this manuscript; Yun GW reviewed the literature, collected data and prepared the tables of this review; and Lee DK designed, reviewed the literature, wrote, revised the paper and performed significant editing.

Correspondence to: Dong Ki Lee, MD, PhD, Department of Internal Medicine, Gangnam Severance Hospital, Yonsei University College of Medicine, 211 Eonju-Ro, Gangnam-Gu, Seoul 135-720, South Korea. dklee@yuhs.ac

Telephone: +82-2-20193214 Fax: + 82-2-34633882

Received: February 27, 2014 Revised: August 2, 2014

Accepted: September 12, 2014

Published online: December 7, 2014

Abstract

Endoscopic retrograde cholangiopancreatography (ERCP) is the essential first modality for common bile duct (CBD) stone therapy. The conventional endoscopic treatment for CBD stones is stone removal after endoscopic sphincterotomy (EST). Stone removal after papillary stretching using balloon dilation instead of the conventional method has been widely adopted. There are many reports regarding endoscopic papillary balloon dilation (EPBD) utilizing a small balloon (< 10 mm) instead of EST for the removal of small CBD stones. In contrast, two cases of mortality due to post-ERCP pancreatitis (PEP) were reported after an EPBD clinical trial in the Western world, and the psychological barrier caused by these incidences hinders the use of this technique in Western countries. Endoscopic papillary large balloon dilation (EPLBD), which is used to treat large CBD stones, was not widely adopted when first

introduced due to concerns about perforation and severe pancreatitis from the use of a large balloon (12-20 mm). However, as experience with this procedure accumulates, the occurrence of PEP with EPLBD is confirmed to be much lower than with EPBD. This report reviews whether EPBD and EPLBD, two procedures that use balloon dilation but differ in terms of indications and concept, contribute to the occurrence of PEP.

© 2014 Baishideng Publishing Group Inc. All rights reserved.

Key words: Post-endoscopic retrograde cholangiopancreatography pancreatitis; Endoscopic papillary balloon dilation; Endoscopic papillary large balloon dilation; Common bile duct stone

Core tip: Endoscopic papillary balloon dilation (EPBD) and endoscopic papillary large balloon dilation (EPLBD) have been performed for removal of common bile duct stones. Although the rates of post-endoscopic retrograde cholangiopancreatography pancreatitis (PEP) after EPBD and EPLBD varied in many studies, the safety and feasibility of balloon dilation have been proven as results have accumulated. However, the exact mechanism of PEP after balloon dilation is unclear. The main determinant of severe PEP may be edema or spasm caused by irritation of the pancreatic orifice while performing difficult selective cannulation and struggling to remove the stone rather than balloon compression of the pancreatic flow.

Jang SI, Yun GW, Lee DK. Balloon dilation itself may not be a major determinant of post-endoscopic retrograde cholangiopancreatography pancreatitis. *World J Gastroenterol* 2014; 20(45): 16913-16924 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v20/i45/16913.htm> DOI: <http://dx.doi.org/10.3748/wjg.v20.i45.16913>

INTRODUCTION

Post-endoscopic retrograde cholangiopancreatography (ERCP) pancreatitis (PEP) is a complicating adverse event, for which doctors can do little to treat. Removing common bile duct (CBD) stones is the most frequently performed procedure carried out using ERCP. Conventional endoscopic biliary stone removal through endoscopic sphincterotomy (EST) has been replaced by the balloon dilation method.

PEP, the most common and potentially serious complication of ERCP-related procedures, occurs in 1%-9% of all procedures^[1]. Many studies of the risk and predictive factors to prevent PEP have been conducted^[1-16]. To discuss the risk factors for PEP, not only procedural and technical factors but also patient characteristics should be considered^[1,15]. Patient-related factors for PEP include a history of post-ERCP pancreatitis^[1,9,10,12,16,17], female sex^[1,10,15], young age^[3,4,6,12,15], suspected sphincter of Oddi dysfunction (SOD)^[1,2,4,10,12,14], and absence of chronic pancreatitis^[1]. Procedural-related factors include difficult cannulation^[1,9,15-17], precut sphincterotomy^[2,10,15,16], pancreatic deep wire pass^[15], pancreatic sphincterotomy^[1], pancreatic contrast injections^[1,4,10,12,17,18], and biliary balloon sphincter dilation^[1]. These reported risk factors vary among studies, and some contradict each other. Hence, the data for risk factors for PEP should be interpreted with caution. Such discrepancies may have arisen from heterogeneous patient populations or from differences in the level of endoscopic expertise, cannulation techniques, and definition of post-ERCP pancreatitis^[12].

The PEP mechanism is not well defined, but it is commonly accepted to be multifactorial, involving mechanical, chemical, hydrostatic, enzymatic, microbial, and thermal factors^[1,12]. A certain triggering event may prematurely activate proteolytic enzymes intracellularly within acinar cells, which may cause cellular injury and autodigestion of pancreatic tissue^[19,20]. Various PEP mechanisms have been suggested. PEP may occur from incidental injection of contrast medium into the pancreatic duct in cases where cannulation of the bile duct is difficult; in such cases, the type of contrast medium injected and the speed and pressure of injection leading to complete acinar filling of the pancreas can have an influence^[8,21-23]. Hydrostatic injury caused by pancreatic duct overfilling may be a major trigger factor for pancreatic reactions^[16]. Difficult cannulation may inflict trauma to the papilla and pancreatic sphincter, leading to pancreatic drainage disruption and causing PEP^[12,24,25]. Pancreatitis after pancreatic sphincterotomy, and precutting have been discussed based on the possibility of incidental temporary obstruction of the pancreatic duct, caused by direct thermal damage to the duct by the cutting wire or by edema induced by thermal tissue injury^[8,26]. PEP occurs more frequently in patients with SOD^[1,2,10,14,24,27-29], which may cause a flow disturbance in pancreatic drainage due to pancreatic sphincter hypertension.

Balloon inflation is also a possible cause of PEP. The pancreatic orifice is compressed during ballooning, and

pancreatic flow is transiently disrupted. However, it is unknown whether ballooning itself is a major determinant for the development of PEP due to endoscopic papillary balloon dilation (EPBD) and endoscopic papillary large balloon dilation (EPLBD). This report provides a systemic review of how the balloon itself affects PEP in balloon dilation during EPBD and EPLBD.

ENDOSCOPIC PAPILLARY BALLOON DILATION AND ENDOSCOPIC PAPILLARY LARGE BALLOON DILATION

Definition and concepts

The ampullary orifice and distal CBD are dilated temporarily by balloon dilation during EPBD and EPLBD procedures. However, EPBD and EPLBD are not the same procedure in either concept or indications. EPBD is performed in patients with a non- or minimally dilated CBD^[30]. In contrast, EPLBD should be performed in patients with dilated CBD and ampulla due to a large long-standing stone. Ballooning of the CBD and ampullary orifice stretches the tissue transiently to form a tubular shape to facilitate stone removal. In this process, successful dilation depends on the elasticity, the degree of dilation of the CBD tissue, and the absence of stricture. Moreover, the ease of selective bile duct cannulation and stone removal through the widened ampullary orifice might be associated with the occurrence of PEP.

EPBD has been performed to dilate the biliary sphincter without prior EST by using a small-diameter dilating balloon (≤ 10 mm). EPBD can avoid the short-term complications of bleeding and perforation, preserve the function of the biliary sphincter, and reduce long-term sequelae of EST. Additionally, EPLBD has been used to remove large CBD stones after dilating the biliary sphincter with EST using a large-diameter dilation (≥ 12 mm). EPLBD can reduce the use of mechanical lithotripsy (MLT), thereby avoiding complications related to the use of full EST for the removal of large or difficult CBD stones. The majority of EPLBDs involve limited EST (minimal or mid-EST rather than full EST) followed by large balloon inflation. If EST is performed toward the CBD, tearing advances in the same direction^[31]. This combined approach does not require full EST and can enlarge the biliary orifice to a greater extent than a standard full EST^[30].

Indications

EPBD is a possible alternative to EST in patients with impaired hemostasis^[11]. To maximize the effect of EPBD while minimizing complications, a technique with proper indication and in the proper manner is necessary^[32]. Recommended indications for balloon dilation include coagulopathy, periampullary diverticulum, Billroth II gastrojejunostomy, and prior EST status^[32]. Another study suggested that the ideal patients are those with a limited number of CBD stones (≤ 3), CBD stones of a maximum diameter ≤ 10 mm, and minimally dilated bile

duct^[33]. In addition, in cases with difficult cannulation, impractical cannulation should be avoided. The use of EPBD for removing CBD stones > 10 mm may necessitate a laborious and papilla-traumatic procedure, such as MLT, and may increase the risk of pancreatitis^[34]. EPBD for a large CBD stone requires multiple sessions and is time-consuming because the biliary orifice is not dilated sufficiently^[35]. Therefore, relatively large stones with a non-dilated CBD are not good candidates for EPBD. Further large-scale studies with a longer follow-up are imperative to identify more distinct indications and the optimum method.

Strict indications are needed for EPLBD to avoid serious adverse events. The patients suitable for this method are those who already have a dilated CBD due to a large stone^[36]. The tissue of ampulla and distal CBD in these patients are ready to be dilated and further gradual stretching of the tissue will not cause stress or sudden tearing of the ampullary roof. However, patients with the CBD of less than the balloon size or strictures of the distal CBD should be excluded because of the possibility of perforation. The velocity and duration of balloon inflation vary across studies, ranging from a few seconds to minutes. Although guidelines pertaining to the optimal velocity of balloon inflation have yet to be established, the following guidelines for safe EPLBD were proposed based on the current knowledge^[36,37]: (1) selection of suitable candidates; *i.e.*, EPLBD should be reserved for patients with a dilated CBD, but avoided in patients with distal CBD strictures; (2) avoidance of full-EST immediately before large balloon dilation to prevent perforation and bleeding; (3) gradual inflation of the dilating balloon to recognize a narrowed distal CBD indicated by lack of disappearance of the balloon waist; (4) discontinuation of inflation when resistance is encountered in the presence of a persistent balloon waist; (5) not inflating the dilation balloon beyond the maximal size of the upstream dilated CBD; and (6) conversion to alternative stone removal or drainage methods when difficulty in removal of a stone is encountered. When a tapered, distal CBD or occult stricture is identified, the operator should pay particular attention to avoiding fatal adverse events caused by large perforations occurring during balloon inflation.

BALLOONING AND PEP

Acute pancreatitis after endoscopic papillary balloon dilation

EPBD, suggested by Staritz *et al.*^[38], is an alternative method to EST for removal of CBD stones. EPBD was adopted to reduce the risk of bleeding^[39-42] and preserve the function of the sphincter of Oddi^[40,43-47]. Some studies have reported that there was no difference in efficacy and safety between EST and EPBD^[46,48], whereas others claimed that the incidence of pancreatitis was higher among patients who received EPBD compared with those who received EST^[33,49,50]. The results of randomized control studies comparing EPBD and

EST are inconsistent, particularly in terms of the incidence and severity of PEP^[46,48-50]. Severe morbidity in the EPBD group compared with the EST group caused early termination of one study^[51], and some clinicians assert that EPBD should not be performed based on the pancreatitis-induced mortality that occurred during some studies^[50,52]. In particular, another randomized, controlled multicenter trial was also terminated early during the first interim analysis, because two patients died from severe pancreatitis as a complication of EPBD^[50]. Thus, the investigation of risk factors for EPBD-related pancreatitis remains controversial.

In studies from Holland^[46] and the United Kingdom^[48], the incidence rates of pancreatitis with EPBD appeared to be similar to those with EST. In a Japanese study, the pancreatitis rate was slightly higher with EPBD than with EST^[49]. In contrast, an American study by Disario *et al.*^[50] reported that the post-EPBD pancreatitis rate was higher than that of EST, and mortality was due to pancreatitis. Although EPBD is used less frequently in Western countries due to these complications, it has been continuously adopted in some Japanese groups. There is no clear explanation for this discrepancy, but it is presumed that differences in patient populations and methods of balloon dilation may play a role^[42]. In addition, the US study may have included patients with SOD^[42]. Thus, pancreatitis-associated EPBD is a very controversial and serious topic^[53,54].

The incidence of acute pancreatitis after EPBD ranges from 5%-20%, although most figures fall in the range of 5%-7%^[43,46,51]. The frequency and severity of PEP following EPBD are summarized in Table 1. The precise mechanism underlying post-EPBD pancreatitis is not well defined, and appears to be multifactorial. Contrast medium injection into the pancreatic duct^[11] and a history of prior pancreatitis^[54] are suggested to be risk factors for post-EPBD pancreatitis. Younger age is a risk factor for post-ERCP pancreatitis^[3,4,6,12,15], but not for post-EPBD pancreatitis^[11,54]. However, in real practice, most cases of severe pancreatitis involve relatively young patients with unatrophied pancreatic tissue.

The mechanism of pancreatitis induced by superfluous injection of contrast medium is regarded as the same as that with EST^[11]. Some research suggests that papillary edema or spasm caused by balloon dilation can result in pancreatitis by obstructing pancreatic outflow^[11,42]. Balloon compression of the papilla or the pancreatic duct orifice may provoke peripapillary edema and/or spasm of the sphincter of Oddi^[51,53]. However, peripapillary trauma by cannulation can more definitely and frequently provoke spasm of the sphincter of Oddi and/or hemorrhagic edematous change^[55], and it is a potential risk factor for asymptomatic hyperamylasemia after EPBD^[54]. In addition, the biliary orifice may not fully dilate during stone removal with EPBD^[56,57]. In this situation, stone removal can be more technically challenging and time-consuming^[46,50,58], and subsequent papillary injury or edema during stone extraction can cause pancreatitis^[35].

Table 1 Frequency and severity of pancreatitis and complications after endoscopic papillary balloon dilation

Ref.	Study design	Study's location	Comparison groups (n)	Pancreatitis n (%)	Pancreatitis severity (n)			Other complications (n)			Overall AEs-related death (n)
					Mild/moderate	Severe	Death	Bleeding	Perforation	Cholangitis	
Minami <i>et al</i> ^[85] , (1995)	RCT	Japan	EPBD (n = 20) EST (n = 20)	2 (10) 2 (10)	2 2	0 0	0 0	0 0	0 0	0 0	0 0
Mathuna <i>et al</i> ^[86] , (1995)	R	Ireland	EPBD (n = 100)	5 (4.8)	5	0	0	0	0	0	0
Bergman <i>et al</i> ^[46] , (1997)	RCT	The Netherlands	EPBD (n = 101) EST (n = 101)	7 (6.9) 7 (6.9)	5 6	2 1	0 0	0 4	2 1	0 0	0 1
Yasuda <i>et al</i> ^[87] , (1998)	P	Japan	EPBD (n = 92)	8 (8.7)	8	0	0	0	0	0	0
Ueno <i>et al</i> ^[55] , (1999)	R	Japan	EPBD (n = 109)	21 (19.8)	21	0	0	NA	NA	NA	0
Ochi <i>et al</i> ^[88] , (1999)	RCT	Japan	EPBD (n = 55) EST (n = 55)	0 2 (3.6)	0 2	0 0	0 0	0 0	0 1	0 0	0 0
Arnold <i>et al</i> ^[51] , (2001)	RCT	Germany	EPBD (n = 30) EST (n = 30)	6 (20.0) 3 (10.0)	4 3	2 0	0 0	0 2	0 0	3 0	0 0
Bergman <i>et al</i> ^[53] , (2001)	RCT	The Netherlands	EPBD (n = 93) EST (n = 87)	7 (7.5) 7 (8.0)	5 6	2 1	0 0	0 2	2 0	0 0	1 0
Yasuda <i>et al</i> ^[43] , (2001)	RCT	Japan	EPBD (n = 35) EST (n = 35)	2 (5.7) 2 (5.7)	2 2	0 0	0 0	0 1	0 0	0 0	0 0
Natsui <i>et al</i> ^[89] , (2002)	RCT	Japan	EPBD (n = 70) EST (n = 70)	4 (5.7) 3 (4.3)	4 3	0 0	0 0	0 2	0 0	2 3	0 0
Fujita <i>et al</i> ^[49] , (2003)	RCT	Japan	EPBD (n = 138) EST (n = 144)	15 (10.9) 4 (2.7)	15 4	0 0	0 0	0 2	0 0	2 6	0 0
Vlavianos <i>et al</i> ^[48] , (2003)	RCT	United Kindom	EPBD (n = 103) EST (n = 99)	5 (4.9) 1 (1.0)	4 1	1 0	0 0	0 0	0 0	0 1	0 1
Sugiyama <i>et al</i> ^[54] , (2003)	R	Japan	EPBD (n = 118)	7 (6.0)	7	0	0	0	0	0	0
Lin <i>et al</i> ^[90] , (2004)	RCT	Taiwan	EPBD (n = 51) EST (n = 53)	0 0	0 0	0 0	0 0	1 14	0 0	0 0	0 0
Disario <i>et al</i> ^[90] , (2004)	RCT	United States	EPBD (n = 117) EST (n = 120)	18 (15.4) 1 (0.8)	12 1	6 0	2 0	11 32	0 1	1 1	2 0
Tanake <i>et al</i> ^[91] , (2004)	RCT	Japan	EPBD (n = 16) EST (n = 16)	3 (18.8) 3 (18.8)	1 2	2 1	0 0	0 0	0 0	0 2	0 0
Toda <i>et al</i> ^[92] , (2005)	RCT	Japan	EPBD (n = 94) EST (n = 102)	7 (6.4) 3 (3)	7 3	0 0	0 0	0 2	0 2	4 4	0 0
Tsujino <i>et al</i> ^[11] , (2005)	R	Japan	EPBD (n = 304)	15 (5.0)	15	0	0	0	1	6	0
Nakagawa <i>et al</i> ^[93] , (2006)	R	Japan	EPBD (n = 201)	2 (1.0)	0	2	0	0	0	3	0
Tsujino <i>et al</i> ^[42] , (2007)	P	Japan	EPBD (n = 1000)	48 (4.8)	47	1	0	2	2	27	2
Ito <i>et al</i> ^[35] , (2008)	R	Japan	EPBD (n = 406)	19 (5.7)	19	0	0	0	1	4	0
Liao <i>et al</i> ^[94] , (2008)	RCT	Taiwan	EPBD (n = 35) EST (n = 25)	2 (5.7) 3 (12)	2 3	0 0	0 0	0 2	0 0	1 2	0 0
Natsui <i>et al</i> ^[95] , (2011)	RCT	Japan	EPBD (n = 41) EST (n = 42)	2 (4.8) 1 (2.3)	2 1	0 0	0 0	0 0	0 0	1 1	0 0
Kuo <i>et al</i> ^[96] , (2012)	R	Taiwan	EPBD (n = 273)	30 (10.1)	22	8	0	1	1	9	1
Seo <i>et al</i> ^[97] , (2014)	RCT	South Korea	EPBD (n = 62) EST (n = 70)	5 (8.1) 5 (7.1)	5 5	0 0	0 0	0 2	0 1	0 0	0 0

EST: Endoscopic sphincterotomy; AEs: Adverse events; EPBD: Endoscopic papillary balloon dilation; R: Retrospective; P: Prospective; RCT: Randomized controlled trial; NA: Not available.

In particular, stone removal by EPBD becomes more difficult when the stone is large and the use of MLT is more frequent^[33,50]. In such cases, the biliary orifice is more likely to be damaged, and the risk of pancreatitis can be greater.

Acute pancreatitis after endoscopic papillary large balloon dilation with EST

EPLBD with limited EST is gradually being recognized as an important modality for the removal of large CBD stones^[36,59-67]. Pancreatitis occurs in 2.4% (0%-13.2%) of

patients; almost all cases have been of mild to moderate severity (98.4%)^[37,68,69]. The frequency and severity of PEP after EPLBD with EST are summarized in Table 2.

Standard procedural guidelines have not been established; yet, most procedures involve limited EST followed by large balloon inflation. If EST is performed toward the CBD, the direction of tearing advances toward the CBD, and less pressure is applied on the pancreatic duct^[31]. It has been suggested that the radial force generated by the dilating balloon is exerted toward the CBD and moves away from the pancreatic duct, which lessens the likelihood of pancreatitis by reducing periampullary injury^[31,36,37,62,70]. Moreover, and in contrast to EPBD, EPLBD dilates the ampullary orifice sufficiently to allow for straightforward removal of a large CBD stone, using a Dormia basket or retrieval balloon, and so that it is wide enough to reduce the need for MLT^[71]. Additionally, because of the patulous papillary orifice caused by a large stone, endoscopists feel comfortable with selective cannulation of the bile duct in most patients. These are all reasons for decreased occurrence of pancreatitis by reduced ampulla injury, which can cause periampullary trauma or edema^[36].

Surprisingly, according to a multicenter retrospective study^[37], there was no severe PEP after EPLBD among 946 patients. This provides strong evidence that ballooning itself is not a major determinant of PEP in EPLBD. Moreover, according to a systemic review of EPLBD, even though the inclusion criteria and procedure type were heterogeneous^[69], PEP following EPLBD was not problematic. The duration of ballooning in EPLBD is usually 30-60 s in real practice. However, the timing of balloon inflation is not related to PEP^[30,69], so prolonged balloon inflation does not increase PEP in EPLBD.

Acute pancreatitis after EPLBD without EST

EPLBD without EST is preferred in patients with bleeding tendencies, altered anatomy, and, in some cases, periampullary diverticulum^[34]. If EST is not performed prior to balloon application, theoretically, the PEP rate may increase because pancreatic outflow could be more completely obstructed by the balloon. In addition, the balloon could press the pancreatic orifice from a more acute angle than when the papillary roof incision is made, because the biliary and pancreatic orifices are not separated. However, according to a retrospective analysis^[30] and systematic review^[69], the PEP rate is not high and does not differ between EPLBD with and without EST. Moreover, the incidence of PEP did not change with ballooning time^[68]. Therefore, ballooning during EPLBD is not a major factor for PEP, regardless of whether EST is performed.

Some recent studies have reported that EPLBD without EST is safe and effective in patients with large CBD stones^[34,72,73]. Pancreatitis and bleeding occurred at a rate of 0.8%-6.5%, and all cases were of mild to moderate severity. The frequency and severity of PEP after EPLBD without EST are summarized in Table 3. This is sup-

ported by a large-scale study reporting less frequent pancreatitis resulting from a larger balloon^[37]. In other words, the extent of biliary orifice dilation is relevant to the incidence of pancreatitis, rather than the size of the balloon, EST performance, or balloon dilation time^[65,68,73].

DISCUSSION

Although EPBD involves a high incidence of pancreatitis, the reports are inconsistent, and it remains controversial. In studies with high rates of pancreatitis, a discrepancy in patient selection should have been made before suggesting balloon dilation as the primary risk factor for pancreatitis. The reason for the high incidence of pancreatitis in EPBD is that enables removal of only small-to-medium sized stones, and patients with such a stone size tend to possess the known risk factors for pancreatitis: young age, non-dilated CBD, normal pancreas parenchyma, obesity, and SOD dysfunction. In other words, careful patient selection can lessen the risk of post-ERCP pancreatitis. EPBD for larger CBD stones, rather than a non-dilated CBD, requires multiple sessions and is more time consuming for stone removal than EST, because EPBD cannot dilate the biliary orifice sufficiently^[35]. Baron *et al.*^[33] recommended extreme caution when performing EPBD in patients with severe acute cholangitis, a history of previous or ongoing acute pancreatitis, age ≤ 50 years, and difficult biliary cannulation. To prevent post-ERCP pancreatitis, pancreatic duct stent insertion is also recommended when EPBD is performed in young patients^[33].

The frequency or severity of PEP after EPBD did not vary with ballooning time. This implies that balloon compression of the pancreatic orifice for < 1 or 2 min without stimulated pancreatic secretion does not cause significant pancreatitis. In one randomized prospective study^[74], 5-min EPBD reduced the risk of pancreatitis compared with conventional 1-min EPBD. Rather than ballooning itself, we believe that pancreatic edema or spasm caused by papillary irritation due to difficult selective cannulation and forcible stone extraction might be the major determinant of PEP after EPBD.

During the early period of EPLBD, PEP is the main concern, because the pancreatic orifice is compressed with a balloon larger than that used in EPBD. However, accumulated data inform clinicians that the larger balloon does not result in PEP, although in practice, one case of severe pancreatitis with mortality has been reported^[75]. Although the major etiological factors of pancreatitis and its mechanism remain unclear, the mechanism of pancreatitis may differ between EPLBD and EPBD. EPLBD and EPBD are different procedures clinically. The major difference is that EPLBD cannot be applied to a non-dilated bile duct, which can be a risk factor for PEP^[54]. If the orifice is sufficiently dilated by EPLBD, papillary edema or spasm is less likely to occur due to use of a basket or retrieval balloon catheter, unlike EPBD, and the incidence of pancreatitis may decline due to the less frequent

Table 2 Frequency and severity of pancreatitis and complications after endoscopic papillary large balloon dilation with endoscopic sphincterotomy

Ref.	Study design	Study's location	Comparison groups (n)	Balloon diameter (mm)	Pancreatitis n (%)	Pancreatitis severity (n)			Other complications (n)			Overall AEs-related death (n)
						Mild/moderate	Severe	death	Bleeding	Perforation	Cholangitis	
Ersoz <i>et al</i> ^[31] , (2003)	R	Turkey	EPLBD (n = 58)	12-20	2 (3.4)	2	0	0	5	0	2	0
Maydeo <i>et al</i> ^[60] , (2007)	P	India	EPLBD (n = 60)	12-20	0 (0)	0	0	0	5	0	0	0
Minami <i>et al</i> ^[62] , (2007)	R	Japan	EPLBD (n = 88)	20	1 (1.1)	1	0	0	1	0	1	0
Heo <i>et al</i> ^[61] , (2007)	RCT	South Korea	EPLBD (n = 100)	12-20	4 (4.0)	4	0	0	0	0	0	0
			EST (n = 100)	-	4 (4.0)	4	0	0	2	0	0	0
Lee <i>et al</i> ^[98] , (2007)	R	South Korea	EPLBD (n = 55)	15-20	0 (0)	0	0	0	2	0	0	0
Kim <i>et al</i> ^[99] , (2007)	R	South Korea	EPLBD (n = 35)	12-20	0 (0)	0	0	0	0	1	0	0
Lee <i>et al</i> ^[100] , (2007)	R	South Korea	EPLBD (n = 41)	13-20	2 (4.8)	0	0	0	1	0	0	0
Misra <i>et al</i> ^[101] , (2008)	R	India	EPLBD (n = 50)	15-20	4 (8.0)	4	0	0	3	0	0	0
Attasaranya <i>et al</i> ^[63] , (2008)	R	United States	EPLBD (n = 103)	12-18	0 (0)	0	0	0	2	1	0	0
Espinel <i>et al</i> ^[102] , (2008)	P	Spain	EPLBD (n = 93)	12-20	1 (1.1)	1	0	0	1	0	0	0
Itoi <i>et al</i> ^[103] , (2009)	R	Japan	EPLBD (n = 53)	15-20	1 (1.9)	1	0	0	0	0	1	0
			EST (n = 48)	-	2 (4.1)	2	0	0	0	0	1	0
Kim <i>et al</i> ^[104] , (2009)	RCT	South Korea	EPLBD (n = 27)	15-18	0 (0)	0	0	0	4	0	0	0
			EST (n = 28)	-	0 (0)	0	0	0	2	0	0	0
Itoi <i>et al</i> ^[105] , (2010)	R	Japan	EPLBD (n = 18)	15-18	0 (0)	0	0	0	0	0	0	0
Kurita <i>et al</i> ^[106] , (2010)	R	Japan	EPLBD (n = 24)	15-20	0 (0)	0	0	0	0	0	0	0
Ghazanfar <i>et al</i> ^[107] , (2010)	P	Pakistan	EPLBD (n = 84)	15-18	3 (3.6)	3	0	0	3	0	0	1
Kim <i>et al</i> ^[108] , (2010)	R	South Korea	EPLBD (n = 70)	12-18	1 (2.3)	1	0	0	0	0	0	0
Youn <i>et al</i> ^[65] , (2011)	R	South Korea	EPLBD (n = 101)	15-20	2 (2.0)	2	0	0	2	1	0	0
Kim <i>et al</i> ^[109] , (2011)	R	South Korea	EPLBD (n = 72)	12-20	5 (6.9)	5	0	0	0	0	1	0
			EST (n = 77)	-	9 (11.7)	9	0	0	0	1	0	0
Stefanidis <i>et al</i> ^[75] , (2011)	RCT	Greece	EPLBD (n = 45)	15-20	1 (2.2)	1	0	0	1	0	0	0
			EST (n = 45)	-	1 (2.2)	1	0	0	1	1	6	0
Rebelo <i>et al</i> ^[67] , (2012)	R	Portugal	EPLBD (n = 30)	12-18	1 (3.3)	1	0	0	0	0	0	0
Sakai <i>et al</i> ^[110] , (2013)	R	Japan	EPLBD (n = 59)	12-20	0 (0)	0	0	0	1	1	0	0
Park <i>et al</i> ^[37] , (2013)	R	South Korea, Japan	EPLBD (n = 946)	12-20	24 (25.3)	24	0	0	56	9	6	4
Poincloux <i>et al</i> ^[64] , (2013)	R	France	EPLBD (n = 64)	15-20	2 (3.1)	2	0	0	5	0	0	0
Hwang <i>et al</i> ^[73] , (2013)	R	South Korea	EPLBD (n = 69)	12-20	3 (4.3)	3	0	0	0	1	0	0
Paspatis <i>et al</i> ^[68] , (2013)	RCT	Greece	60 s dilation ¹ (n = 60)	15-20	2 (1.6)	2	0	0	4	1	3	0
			30 s dilation ¹ (n = 64)	15-20	2 (1.6)	1	1	1	2	1	2	1
Rosa <i>et al</i> ^[66] , (2013)	R	Portugal	EPLBD (n = 68)	12-18	9 (13.2)	9	0	0	0	0	1	0
			EST (n = 45)	-	2 (4.7)	2	0	0	0	0	1	0

Yang <i>et al.</i> ^[111] , (2013)	R	China	EPLBD (n = 171)	12-18	2 (1.2)	2	0	0	4	1	1	0
Yoon <i>et al.</i> ^[112] , (2013)	P	South Korea	EPLBD (n = 52)	12-20	0 (0)	0	0	0	0	0	0	0
Harada <i>et al.</i> ^[113] , (2013)	R	Japan	EPLBD (n = 30)	15-20	0 (0)	0	0	0	0	0	0	0

¹Sixty and thirty seconds mean the duration of balloon dilation (s) in endoscopic papillary large balloon dilation (EPLBD). AEs: Adverse events; EST: Endoscopic sphincterotomy; R: Retrospective; P: Prospective; RCT: Randomized controlled trial.

Table 3 Frequency and severity of pancreatitis and complications after endoscopic papillary large balloon dilation without endoscopic sphincterotomy

Ref.	Years	Study design	Patients (n)	Pancreatitis n (%)	Pancreatitis severity (n)				Other complications (n)			Overall AEs-related death (n)
					Mild	Moderate	Severe	Death	Bleeding	Perforation	Cholangitis	
Jeong <i>et al.</i> ^[34]	2009	R	38	1 (2.6)	1	0	0	0	0	0	0	0
Chan <i>et al.</i> ^[72]	2011	R	247	2 (0.8)	2	0	0	0	0	0	1	0
Hwang <i>et al.</i> ^[73]	2013	R	62	4 (6.4)	4	0	0	0	0	0	0	0

AEs: Adverse events; R: Retrospective.

use of MLT^[34,72]. There is no significant difference in the frequency of the requirement for MLT between EPLBD and the conventional method. However, adequate fragmentation of the CBD stone by MLT after EPLBD can reduce the frequency of the requirement for MLT in EPLBD compared with the conventional method^[69]. This could be one reason for the lower incidence of PEP in EPLBD. Patients who receive EPLBD are relatively older individuals in whom pancreatic exocrine function has declined, and pancreatitis is less likely^[72]. Further, easy selective cannulation into the bile duct can reduce the incidence of pancreatitis. Additionally, contrary to our concern, EPLBD without a preceding papillary incision did not cause severe pancreatitis^[34,72,73]. Therefore, ballooning itself may not be the culprit. And the cause of fatal pancreatitis during EPBD should be reconsidered.

The incidence of pancreatitis, using percutaneous papillary balloon dilation (PTPBD) for CBD stone removal, is extremely low (0%-1.4%)^[76-82]. A retrospective study reported that pancreatitis occurred only in the EPBD group; in another study, comparing PTPBD with EPBD, the only significant predictor was the use of MLT^[76]. The size of the balloon used in PTPBD varies (4-23 mm) among studies and can also vary within the same study, due to the presence of differently sized CBD stones (5-20 mm)^[77-82]. In one study^[76], patients with a CBD stone < 12 mm in diameter were enrolled homogeneously, and the balloon dilation diameter was 8-10 mm, which was compatible with EPBD. The largest balloons used for papillary dilation were of diameters 22 mm^[80] and 23 mm^[82] in other studies; this balloon inflation size is compatible with EPLBD. Although the balloon dilation diameter was different in each study, no severe pancreatitis occurred. These studies confirmed that ballooning does not increase the incidence of PEP. Moreover, the rates of post-procedural pancreatitis and hyperamylasemia were significantly higher following retrograde dilation using EPBD, compared with antegrade dilation using

PTPBD, during the removal of bile duct stones^[76].

The reason for the lower rate of PEP with antegrade application of balloon inflation compared with a retrograde fashion is the lack of difficulty in selective cannulation and lower chance of difficult procedure for forcible removal unless the stone descends. Compared with EPBD, PTPBD inflicts less mechanical trauma to papilla during stone removal, and it is nearly equivalent to the effect of ballooning^[83,84]. In addition, MLT application does not involve lithotripsy moving back and forth through the ampullary orifice, in which there is no chance of pancreatic orifice damage. Such a result demonstrates that ballooning may not be a risk factor for pancreatitis. Moreover, the rates of post-procedural pancreatitis and hyperamylasemia were significantly higher after retrograde dilation with EPBD than after antegrade dilation with PTPBD for the removal of bile duct stones. This reveals that pancreatitis can be induced by other factors, such as repeated cannulation or pancreatic duct injection, during retrograde dilation with EPBD.

CONCLUSION

Although the mechanism of PEP is unclear, the occurrence of pancreatitis is more associated with the catheter, basket, or MLT causing ampullary injury. Instead of balloon compression of the pancreatic flow, the main determinants of severe pancreatitis during endoscopic stone removal with balloon dilation may involve edema or spasm caused by irritation of the pancreatic orifice while performing difficult selective cannulation and struggling to remove the stone. Therefore, ballooning itself may not be the culprit for PEP in either EPBD or EPLBD.

REFERENCES

- 1 Freeman ML, DiSario JA, Nelson DB, Fennerty MB, Lee JG, Bjorkman DJ, Overby CS, Aas J, Ryan ME, Bochna GS, Shaw

- MJ, Snady HW, Erickson RV, Moore JP, Roel JP. Risk factors for post-ERCP pancreatitis: a prospective, multicenter study. *Gastrointest Endosc* 2001; **54**: 425-434 [PMID: 11577302]
- 2 **Freeman ML**, Nelson DB, Sherman S, Haber GB, Herman ME, Dorsher PJ, Moore JP, Fennerty MB, Ryan ME, Shaw MJ, Lande JD, Pheley AM. Complications of endoscopic biliary sphincterotomy. *N Engl J Med* 1996; **335**: 909-918 [PMID: 8782497 DOI: 10.1056/NEJM199609263351301]
 - 3 **Loperfido S**, Angelini G, Benedetti G, Chilovi F, Costan F, De Berardinis F, De Bernardin M, Ederle A, Fina P, Fratton A. Major early complications from diagnostic and therapeutic ERCP: a prospective multicenter study. *Gastrointest Endosc* 1998; **48**: 1-10 [PMID: 9684657]
 - 4 **Mehta SN**, Pavone E, Barkun JS, Bouchard S, Barkun AN. Predictors of post-ERCP complications in patients with suspected choledocholithiasis. *Endoscopy* 1998; **30**: 457-463 [PMID: 9693893 DOI: 10.1055/s-2007-1001308]
 - 5 **Rabenstein T**, Schneider HT, Bulling D, Nicklas M, Katalinic A, Hahn EG, Martus P, Ell C. Analysis of the risk factors associated with endoscopic sphincterotomy techniques: preliminary results of a prospective study, with emphasis on the reduced risk of acute pancreatitis with low-dose anticoagulation treatment. *Endoscopy* 2000; **32**: 10-19 [PMID: 10691266 DOI: 10.1055/s-2000-138]
 - 6 **Masci E**, Toti G, Mariani A, Curioni S, Lomazzi A, Dinelli M, Minoli G, Crosta C, Comin U, Fertitta A, Prada A, Passoni GR, Testoni PA. Complications of diagnostic and therapeutic ERCP: a prospective multicenter study. *Am J Gastroenterol* 2001; **96**: 417-423 [PMID: 11232684 DOI: 10.1111/j.1572-0241.2001.03594.x]
 - 7 **Aronson N**, Flamm CR, Bohn RL, Mark DH, Speroff T. Evidence-based assessment: patient, procedure, or operator factors associated with ERCP complications. *Gastrointest Endosc* 2002; **56**: S294-S302 [PMID: 12447284 DOI: 10.1067/mge.2002.129021]
 - 8 **Christoforidis E**, Goulimaris I, Kanellos I, Tsalis K, Demetriades C, Betsis D. Post-ERCP pancreatitis and hyperamylasemia: patient-related and operative risk factors. *Endoscopy* 2002; **34**: 286-292 [PMID: 11932783 DOI: 10.1055/s-2002-23630]
 - 9 **Vandervoort J**, Soetikno RM, Tham TC, Wong RC, Ferrari AP, Montes H, Roston AD, Slivka A, Lichtenstein DR, Ruyman FW, Van Dam J, Hughes M, Carr-Locke DL. Risk factors for complications after performance of ERCP. *Gastrointest Endosc* 2002; **56**: 652-656 [PMID: 12397271 DOI: 10.1067/mge.2002.129086]
 - 10 **Masci E**, Mariani A, Curioni S, Testoni PA. Risk factors for pancreatitis following endoscopic retrograde cholangiopancreatography: a meta-analysis. *Endoscopy* 2003; **35**: 830-834 [PMID: 14551860 DOI: 10.1055/s-2003-42614]
 - 11 **Tsujino T**, Isayama H, Komatsu Y, Ito Y, Tada M, Minagawa N, Nakata R, Kawabe T, Omata M. Risk factors for pancreatitis in patients with common bile duct stones managed by endoscopic papillary balloon dilation. *Am J Gastroenterol* 2005; **100**: 38-42 [PMID: 15654778 DOI: 10.1111/j.1572-0241.2005.40638.x]
 - 12 **Cheng CL**, Sherman S, Watkins JL, Barnett J, Freeman M, Geenen J, Ryan M, Parker H, Frakes JT, Fogel EL, Silverman WB, Dua KS, Aliperti G, Yakshe P, Uzer M, Jones W, Goff J, Lazzell-Pannell L, Rashdan A, Temkit M, Lehman GA. Risk factors for post-ERCP pancreatitis: a prospective multicenter study. *Am J Gastroenterol* 2006; **101**: 139-147 [PMID: 16405547 DOI: 10.1111/j.1572-0241.2006.00380.x]
 - 13 **Williams EJ**, Taylor S, Fairclough P, Hamlyn A, Logan RF, Martin D, Riley SA, Veitch P, Wilkinson ML, Williamson PR, Lombard M. Risk factors for complication following ERCP; results of a large-scale, prospective multicenter study. *Endoscopy* 2007; **39**: 793-801 [PMID: 17703388 DOI: 10.1055/s-2007-966723]
 - 14 **Cotton PB**, Garrow DA, Gallagher J, Romagnuolo J. Risk factors for complications after ERCP: a multivariate analysis of 11,497 procedures over 12 years. *Gastrointest Endosc* 2009; **70**: 80-88 [PMID: 19286178 DOI: 10.1016/j.gie.2008.10.039]
 - 15 **Wang P**, Li ZS, Liu F, Ren X, Lu NH, Fan ZN, Huang Q, Zhang X, He LP, Sun WS, Zhao Q, Shi RH, Tian ZB, Li YQ, Li W, Zhi FC. Risk factors for ERCP-related complications: a prospective multicenter study. *Am J Gastroenterol* 2009; **104**: 31-40 [PMID: 19098846 DOI: 10.1038/ajg.2008.5]
 - 16 **Testoni PA**, Mariani A, Giussani A, Vailati C, Masci E, Maccarri G, Ghezzi L, Familiari L, Giardullo N, Mutignani M, Lombardi G, Talamini G, Spadaccini A, Briglia R, Piazzi L. Risk factors for post-ERCP pancreatitis in high- and low-volume centers and among expert and non-expert operators: a prospective multicenter study. *Am J Gastroenterol* 2010; **105**: 1753-1761 [PMID: 20372116 DOI: 10.1038/ajg.2010.136]
 - 17 **Friedland S**, Soetikno RM, Vandervoort J, Montes H, Tham T, Carr-Locke DL. Bedside scoring system to predict the risk of developing pancreatitis following ERCP. *Endoscopy* 2002; **34**: 483-488 [PMID: 12048633 DOI: 10.1055/s-2002-32004]
 - 18 **Andriulli A**, Clemente R, Solmi L, Terruzzi V, Suriani R, Sigillito A, Leandro G, Leo P, De Maio G, Perri F. Gabexate or somatostatin administration before ERCP in patients at high risk for post-ERCP pancreatitis: a multicenter, placebo-controlled, randomized clinical trial. *Gastrointest Endosc* 2002; **56**: 488-495 [PMID: 12297762 DOI: 10.1067/mge.2002.128130]
 - 19 **Gottlieb K**, Sherman S. ERCP and biliary endoscopic sphincterotomy-induced pancreatitis. *Gastrointest Endosc Clin N Am* 1998; **8**: 87-114 [PMID: 9405753]
 - 20 **Hofbauer B**, Saluja AK, Lerch MM, Bhagat L, Bhatia M, Lee HS, Frossard JL, Adler G, Steer ML. Intra-acinar cell activation of trypsinogen during caerulein-induced pancreatitis in rats. *Am J Physiol* 1998; **275**: G352-G362 [PMID: 9688663]
 - 21 **Barkin JS**, Casal GL, Reiner DK, Goldberg RI, Phillips RS, Kaplan S. A comparative study of contrast agents for endoscopic retrograde pancreatography. *Am J Gastroenterol* 1991; **86**: 1437-1441 [PMID: 1928034]
 - 22 **Lüthen R**, Niederau C, Niederau M, Ferrell LD, Grendell JH. Influence of ductal pressure and infusates on activity and subcellular distribution of lysosomal enzymes in the rat pancreas. *Gastroenterology* 1995; **109**: 573-581 [PMID: 7615208]
 - 23 **Hacıahmetoglu T**, Ertekin C, Dolay K, Yanar F, Yanar H, Kapran Y. The effects of contrast agent and intraductal pressure changes on the development of pancreatitis in an ERCP model in rats. *Langenbecks Arch Surg* 2008; **393**: 367-372 [PMID: 17674029 DOI: 10.1007/s00423-007-0214-1]
 - 24 **Sherman S**, Lehman GA. ERCP- and endoscopic sphincterotomy-induced pancreatitis. *Pancreas* 1991; **6**: 350-367 [PMID: 1713676]
 - 25 **Saari A**, Kivisaari L, Standertskjöld-Nordenstam CG, Brackett K, Schröder T. Experimental pancreatography: a comparison of three contrast media. *Scand J Gastroenterol* 1988; **23**: 53-58 [PMID: 3344398]
 - 26 **Sivak MV**. Endoscopic management of bile duct stones. *Am J Surg* 1989; **158**: 228-240 [PMID: 2672845]
 - 27 **Chen YK**, Foliente RL, Santoro MJ, Walter MH, Collen MJ. Endoscopic sphincterotomy-induced pancreatitis: increased risk associated with nondilated bile ducts and sphincter of Oddi dysfunction. *Am J Gastroenterol* 1994; **89**: 327-333 [PMID: 8122639]
 - 28 **Sherman S**, Ruffolo TA, Hawes RH, Lehman GA. Complications of endoscopic sphincterotomy. A prospective series with emphasis on the increased risk associated with sphincter of Oddi dysfunction and nondilated bile ducts. *Gastroenterology* 1991; **101**: 1068-1075 [PMID: 1889699]
 - 29 **Fogel EL**, Eversman D, Jamidar P, Sherman S, Lehman GA. Sphincter of Oddi dysfunction: pancreaticobiliary sphincterotomy with pancreatic stent placement has a lower rate of pancreatitis than biliary sphincterotomy alone. *Endoscopy*

- 2002; **34**: 280-285 [PMID: 1889699]
- 30 **Jeong SU**, Moon SH, Kim MH. Endoscopic papillary balloon dilation: revival of the old technique. *World J Gastroenterol* 2013; **19**: 8258-8268 [PMID: 24363517 DOI: 10.3748/wjg.v19.i45.8258]
- 31 **Ersoz G**, Tekesin O, Ozutemiz AO, Gunsar F. Biliary sphincterotomy plus dilation with a large balloon for bile duct stones that are difficult to extract. *Gastrointest Endosc* 2003; **57**: 156-159 [PMID: 12556775 DOI: 10.1067/Mge.2003.52]
- 32 **Chung JW**, Chung JB. Endoscopic papillary balloon dilation for removal of choledocholithiasis: indications, advantages, complications, and long-term follow-up results. *Gut Liver* 2011; **5**: 1-14 [PMID: 21461066 DOI: 10.5009/gnl.2011.5.1.1]
- 33 **Baron TH**, Harewood GC. Endoscopic balloon dilation of the biliary sphincter compared to endoscopic biliary sphincterotomy for removal of common bile duct stones during ERCP: a metaanalysis of randomized, controlled trials. *Am J Gastroenterol* 2004; **99**: 1455-1460 [PMID: 15307859 DOI: 10.1111/j.1572-0241.2004.30151.x]
- 34 **Jeong S**, Ki SH, Lee DH, Lee JI, Lee JW, Kwon KS, Kim HG, Shin YW, Kim YS. Endoscopic large-balloon sphincteroplasty without preceding sphincterotomy for the removal of large bile duct stones: a preliminary study. *Gastrointest Endosc* 2009; **70**: 915-922 [PMID: 19647241 DOI: 10.1016/j.gie.2009.04.042]
- 35 **Ito Y**, Tsujino T, Togawa O, Yamamoto N, Isayama H, Nakata R, Kawabe T, Omata M. Endoscopic papillary balloon dilation for the management of bile duct stones in patients 85 years of age and older. *Gastrointest Endosc* 2008; **68**: 477-482 [PMID: 18760175 DOI: 10.1016/j.gie.2007.10.066]
- 36 **Lee DK**, Han JW. Endoscopic papillary large balloon dilation: guidelines for pursuing zero mortality. *Clin Endosc* 2012; **45**: 299-304 [PMID: 22977823 DOI: 10.5946/ce.2012.45.3.299]
- 37 **Park SJ**, Kim JH, Hwang JC, Kim HG, Lee DH, Jeong S, Cha SW, Cho YD, Kim HJ, Kim JH, Moon JH, Park SH, Itoi T, Isayama H, Kogure H, Lee SJ, Jung KT, Lee HS, Baron TH, Lee DK. Factors predictive of adverse events following endoscopic papillary large balloon dilation: results from a multicenter series. *Dig Dis Sci* 2013; **58**: 1100-1109 [PMID: 23225136 DOI: 10.1007/s10620-012-2494-8]
- 38 **Staritz M**, Ewe K, Meyer zum Büschenfelde KH. Endoscopic papillary dilation (EPD) for the treatment of common bile duct stones and papillary stenosis. *Endoscopy* 1983; **15** Suppl 1: 197-198 [PMID: 6872989 DOI: 10.1055/s-2007-1021507]
- 39 **Kawabe T**, Komatsu Y, Tada M, Toda N, Ohashi M, Shiratori Y, Omata M. Endoscopic papillary balloon dilation in cirrhotic patients: removal of common bile duct stones without sphincterotomy. *Endoscopy* 1996; **28**: 694-698 [PMID: 8934088 DOI: 10.1055/s-2007-1005579]
- 40 **Komatsu Y**, Kawabe T, Toda N, Ohashi M, Isayama M, Tateishi K, Sato S, Koike Y, Yamagata M, Tada M, Shiratori Y, Yamada H, Ithori M, Kawase T, Omata M. Endoscopic papillary balloon dilation for the management of common bile duct stones: experience of 226 cases. *Endoscopy* 1998; **30**: 12-17 [PMID: 9548037 DOI: 10.1055/s-2007-993721]
- 41 **Park DH**, Kim MH, Lee SK, Lee SS, Choi JS, Song MH, Seo DW, Min YI. Endoscopic sphincterotomy vs. endoscopic papillary balloon dilation for choledocholithiasis in patients with liver cirrhosis and coagulopathy. *Gastrointest Endosc* 2004; **60**: 180-185 [PMID: 15278041]
- 42 **Tsujino T**, Kawabe T, Komatsu Y, Yoshida H, Isayama H, Sasaki T, Kogure H, Togawa O, Arizumi T, Matsubara S, Ito Y, Nakai Y, Yamamoto N, Sasahira N, Hirano K, Toda N, Tada M, Omata M. Endoscopic papillary balloon dilation for bile duct stone: immediate and long-term outcomes in 1000 patients. *Clin Gastroenterol Hepatol* 2007; **5**: 130-137 [PMID: 17234559 DOI: 10.1016/j.cgh.2006.10.013]
- 43 **Yasuda I**, Tomita E, Enya M, Kato T, Moriwaki H. Can endoscopic papillary balloon dilation really preserve sphincter of Oddi function? *Gut* 2001; **49**: 686-691 [PMID: 11600473]
- 44 **Kawabe T**, Komatsu Y, Isayama H, Takemura T, Toda N, Tada M, Imai Y, Shiratori Y, Omata M. Histological analysis of the papilla after endoscopic papillary balloon dilation. *Hepatogastroenterology* 2003; **50**: 919-923 [PMID: 12845950]
- 45 **Isayama H**, Komatsu Y, Inoue Y, Toda N, Shiratori Y, Tsujino T, Yamada H, Saitou K, Kawabe T, Omata M. Preserved function of the Oddi sphincter after endoscopic papillary balloon dilation. *Hepatogastroenterology* 2003; **50**: 1787-1791 [PMID: 14696405]
- 46 **Bergman JJ**, Rauws EA, Fockens P, van Berkel AM, Bossuyt PM, Tijssen JG, Tytgat GN, Huibregtse K. Randomised trial of endoscopic balloon dilation versus endoscopic sphincterotomy for removal of bileduct stones. *Lancet* 1997; **349**: 1124-1129 [PMID: 9113010 DOI: 10.1016/S0140-6736(96)11026-6]
- 47 **Sato H**, Kodama T, Takaaki J, Tatsumi Y, Maeda T, Fujita S, Fukui Y, Ogasawara H, Mitsufuji S. Endoscopic papillary balloon dilatation may preserve sphincter of Oddi function after common bile duct stone management: evaluation from the viewpoint of endoscopic manometry. *Gut* 1997; **41**: 541-544 [PMID: 9391256]
- 48 **Vlavianos P**, Chopra K, Mandalia S, Anderson M, Thompson J, Westaby D. Endoscopic balloon dilatation versus endoscopic sphincterotomy for the removal of bile duct stones: a prospective randomised trial. *Gut* 2003; **52**: 1165-1169 [PMID: 12865276]
- 49 **Fujita N**, Maguchi H, Komatsu Y, Yasuda I, Hasebe O, Igarashi Y, Murakami A, Mukai H, Fujii T, Yamao K, Maeshiro K. Endoscopic sphincterotomy and endoscopic papillary balloon dilatation for bile duct stones: A prospective randomized controlled multicenter trial. *Gastrointest Endosc* 2003; **57**: 151-155 [PMID: 12556774 DOI: 10.1067/mge.2003.56]
- 50 **Disario JA**, Freeman ML, Bjorkman DJ, Macmathuna P, Petersen BT, Jaffe PE, Morales TG, Hixson LJ, Sherman S, Lehman GA, Jamal MM, Al-Kawas FH, Khandelwal M, Moore JP, Derfus GA, Jamidar PA, Ramirez FC, Ryan ME, Woods KL, Carr-Locke DL, Alder SC. Endoscopic balloon dilation compared with sphincterotomy for extraction of bile duct stones. *Gastroenterology* 2004; **127**: 1291-1299 [PMID: 15520997]
- 51 **Arnold JC**, Benz C, Martin WR, Adamek HE, Riemann JF. Endoscopic papillary balloon dilation vs. sphincterotomy for removal of common bile duct stones: a prospective randomized pilot study. *Endoscopy* 2001; **33**: 563-567 [PMID: 11473325 DOI: 10.1055/s-2001-15307]
- 52 **Kozarek RA**. Balloon dilation of the sphincter of Oddi. *Endoscopy* 1988; **20** Suppl 1: 207-210 [PMID: 3168949 DOI: 10.1055/s-2007-1018177]
- 53 **Bergman JJ**, van Berkel AM, Bruno MJ, Fockens P, Rauws EA, Tijssen JG, Tytgat GN, Huibregtse K. Is endoscopic balloon dilation for removal of bile duct stones associated with an increased risk for pancreatitis or a higher rate of hyperamylasemia? *Endoscopy* 2001; **33**: 416-420 [PMID: 11396759 DOI: 10.1055/s-2001-14424]
- 54 **Sugiyama M**, Izumisato Y, Abe N, Masaki T, Mori T, Atomi Y. Predictive factors for acute pancreatitis and hyperamylasemia after endoscopic papillary balloon dilation. *Gastrointest Endosc* 2003; **57**: 531-535 [PMID: 12665764 DOI: 10.1067/mge.2003.143]
- 55 **Ueno N**, Ozawa Y. Pancreatitis induced by endoscopic balloon sphincter dilation and changes in serum amylase levels after the procedure. *Gastrointest Endosc* 1999; **49**: 472-476 [PMID: 10202061]
- 56 **Weinberg BM**, Shindy W, Lo S. Endoscopic balloon sphincter dilation (sphincteroplasty) versus sphincterotomy for common bile duct stones. *Cochrane Database Syst Rev* 2006; **(4)**: CD004890 [PMID: 17054222 DOI: 10.1002/14651858.CD004890.pub2]

- 57 **Zhao HC**, He L, Zhou DC, Geng XP, Pan FM. Meta-analysis comparison of endoscopic papillary balloon dilatation and endoscopic sphincteroplasty. *World J Gastroenterol* 2013; **19**: 3883-3891 [PMID: 23840129 DOI: 10.3748/wjg.v19.i24.3883]
- 58 **Minami A**, Maeta T, Kohi F, Nakatsu T, Morshed SA, Nishioka M. Endoscopic papillary dilation by balloon and isosorbide dinitrate drip infusion for removing bile duct stone. *Scand J Gastroenterol* 1998; **33**: 765-768 [PMID: 9712243]
- 59 **Attam R**, Freeman ML. Endoscopic papillary large balloon dilation for large common bile duct stones. *J Hepatobiliary Pancreat Surg* 2009; **16**: 618-623 [PMID: 19551331 DOI: 10.1007/s00534-009-0134-2]
- 60 **Maydeo A**, Bhandari S. Balloon sphincteroplasty for removing difficult bile duct stones. *Endoscopy* 2007; **39**: 958-961 [PMID: 17701853 DOI: 10.1055/s-2007-966784]
- 61 **Heo JH**, Kang DH, Jung HJ, Kwon DS, An JK, Kim BS, Suh KD, Lee SY, Lee JH, Kim GH, Kim TO, Heo J, Song GA, Cho M. Endoscopic sphincterotomy plus large-balloon dilation versus endoscopic sphincterotomy for removal of bile-duct stones. *Gastrointest Endosc* 2007; **66**: 720-726; quiz 768, 771 [PMID: 17905013 DOI: 10.1016/j.gie.2007.02.033]
- 62 **Minami A**, Hirose S, Nomoto T, Hayakawa S. Small sphincterotomy combined with papillary dilation with large balloon permits retrieval of large stones without mechanical lithotripsy. *World J Gastroenterol* 2007; **13**: 2179-2182 [PMID: 17465497]
- 63 **Attasaranya S**, Cheon YK, Vittal H, Howell DA, Wakelin DE, Cunningham JT, Ajmere N, Ste Marie RW, Bhattacharya K, Gupta K, Freeman ML, Sherman S, McHenry L, Watkins JL, Fogel EL, Schmidt S, Lehman GA. Large-diameter biliary orifice balloon dilation to aid in endoscopic bile duct stone removal: a multicenter series. *Gastrointest Endosc* 2008; **67**: 1046-1052 [PMID: 18178208 DOI: 10.1016/j.gie.2007.08.047]
- 64 **Poincloux L**, Rouquette O, Privat J, Gorce D, Abergel A, Dapoigny M, Bommelaer G. Large-balloon dilation of the sphincter of Oddi after sphincterotomy or infundibulotomy to extract large calculi or multiple common bile duct stones without using mechanical lithotripsy. *Scand J Gastroenterol* 2013; **48**: 246-251 [PMID: 22229762 DOI: 10.3109/00365521.2011.647064]
- 65 **Youn YH**, Lim HC, Jahng JH, Jang SI, You JH, Park JS, Lee SJ, Lee DK. The increase in balloon size to over 15 mm does not affect the development of pancreatitis after endoscopic papillary large balloon dilatation for bile duct stone removal. *Dig Dis Sci* 2011; **56**: 1572-1577 [PMID: 20945093 DOI: 10.1007/s10620-010-1438-4]
- 66 **Rosa B**, Moutinho Ribeiro P, Rebelo A, Pinto Correia A, Cotter J. Endoscopic papillary balloon dilation after sphincterotomy for difficult choledocholithiasis: A case-controlled study. *World J Gastrointest Endosc* 2013; **5**: 211-218 [PMID: 23678373 DOI: 10.4253/wjge.v5.i5.211]
- 67 **Rebelo A**, Ribeiro PM, Correia AP, Cotter J. Endoscopic papillary large balloon dilation after limited sphincterotomy for difficult biliary stones. *World J Gastrointest Endosc* 2012; **4**: 180-184 [PMID: 22624069 DOI: 10.4253/wjge.v4.i5.180]
- 68 **Paspatis GA**, Konstantinidis K, Tribonias G, Voudoukis E, Tavernaraki A, Theodoropoulou A, Chainaki I, Manolaraki M, Chlouverakis G, Vardas E, Paraskeva K. Sixty- versus thirty-seconds papillary balloon dilation after sphincterotomy for the treatment of large bile duct stones: a randomized controlled trial. *Dig Liver Dis* 2013; **45**: 301-304 [PMID: 23195665 DOI: 10.1016/j.dld.2012.10.015]
- 69 **Kim JH**, Yang MJ, Hwang JC, Yoo BM. Endoscopic papillary large balloon dilation for the removal of bile duct stones. *World J Gastroenterol* 2013; **19**: 8580-8594 [PMID: 24379575 DOI: 10.3748/wjg.v19.i46.8580]
- 70 **Akashi R**, Kiyozumi T, Tanaka T, Sakurai K, Oda Y, Sagara K. Mechanism of pancreatitis caused by ERCP. *Gastrointest Endosc* 2002; **55**: 50-54 [PMID: 11756914 DOI: 10.1067/mge.2002.118964]
- 71 **Feng Y**, Zhu H, Chen X, Xu S, Cheng W, Ni J, Shi R. Comparison of endoscopic papillary large balloon dilation and endoscopic sphincterotomy for retrieval of choledocholithiasis: a meta-analysis of randomized controlled trials. *J Gastroenterol* 2012; **47**: 655-663 [PMID: 22361862 DOI: 10.1007/s00535-012-0528-9]
- 72 **Chan HH**, Lai KH, Lin CK, Tsai WL, Wang EM, Hsu PI, Chen WC, Yu HC, Wang HM, Tsay FW, Tsai CC, Chen IS, Chen YC, Liang HL, Pan HB. Endoscopic papillary large balloon dilation alone without sphincterotomy for the treatment of large common bile duct stones. *BMC Gastroenterol* 2011; **11**: 69 [PMID: 21668994 DOI: 10.1186/1471-230X-11-69]
- 73 **Hwang JC**, Kim JH, Lim SG, Kim SS, Shin SJ, Lee KM, Yoo BM. Endoscopic large-balloon dilation alone versus endoscopic sphincterotomy plus large-balloon dilation for the treatment of large bile duct stones. *BMC Gastroenterol* 2013; **13**: 15 [PMID: 23324454 DOI: 10.1186/1471-230X-13-15]
- 74 **Liao WC**, Lee CT, Chang CY, Leung JW, Chen JH, Tsai MC, Lin JT, Wu MS, Wang HP. Randomized trial of 1-minute versus 5-minute endoscopic balloon dilation for extraction of bile duct stones. *Gastrointest Endosc* 2010; **72**: 1154-1162 [PMID: 20869710 DOI: 10.1016/j.gie.2010.07.009]
- 75 **Stefanidis G**, Viazis N, Pleskow D, Manolakopoulos S, Theocharis L, Christodoulou C, Kotsikoros N, Giannousis J, Sgourou S, Rodias M, Katsikani A, Chuttani R. Large balloon dilation vs. mechanical lithotripsy for the management of large bile duct stones: a prospective randomized study. *Am J Gastroenterol* 2011; **106**: 278-285 [PMID: 21045816 DOI: 10.1038/ajg.2010.421]
- 76 **Seo YR**, Moon JH, Choi HJ, Kim DC, Lee TH, Cha SW, Cho YD, Park SH, Kim SJ. Papillary balloon dilation is not itself a cause of post-endoscopic retrograde cholangiopancreatography pancreatitis; results of anterograde and retrograde papillary balloon dilation. *J Gastroenterol Hepatol* 2013; **28**: 1416-1421 [PMID: 23701518 DOI: 10.1111/jgh.12277]
- 77 **Graziani L**, Fabrizzi G, Manfrini E, Galeazzi R, Freddara U. Percutaneous transhepatic Oddi-sphincter dilatation for bile duct stone removal. *AJR Am J Roentgenol* 1989; **152**: 73-75 [PMID: 2783292 DOI: 10.2214/ajr.152.1.73]
- 78 **Gil S**, de la Iglesia P, Verdú JF, de España F, Arenas J, Irurzun J. Effectiveness and safety of balloon dilation of the papilla and the use of an occlusion balloon for clearance of bile duct calculi. *AJR Am J Roentgenol* 2000; **174**: 1455-1460 [PMID: 10789811 DOI: 10.2214/ajr.174.5.1741455]
- 79 **Park YS**, Kim JH, Choi YW, Lee TH, Hwang CM, Cho YJ, Kim KW. Percutaneous treatment of extrahepatic bile duct stones assisted by balloon sphincteroplasty and occlusion balloon. *Korean J Radiol* 2005; **6**: 235-240 [PMID: 16374081]
- 80 **Szulman C**, Giménez M, Sierre S. Antegrade papillary balloon dilation for extrahepatic bile duct stone clearance: lessons learned from treating 300 patients. *J Vasc Interv Radiol* 2011; **22**: 346-353 [PMID: 21277793 DOI: 10.1016/j.jvir.2010.11.015]
- 81 **García-García L**, Lanciego C. Percutaneous treatment of biliary stones: sphincteroplasty and occlusion balloon for the clearance of bile duct calculi. *AJR Am J Roentgenol* 2004; **182**: 663-670 [PMID: 14975967 DOI: 10.2214/ajr.182.3.1820663]
- 82 **García-Vila JH**, Redondo-Ibáñez M, Díaz-Ramón C. Balloon sphincteroplasty and transpapillary elimination of bile duct stones: 10 years' experience. *AJR Am J Roentgenol* 2004; **182**: 1451-1458 [PMID: 15149989 DOI: 10.2214/ajr.182.6.1821451]
- 83 **Moon JH**, Cho YD, Ryu CB, Kim JO, Cho JY, Kim YS, Lee JS, Lee MS, Shim CS. The role of percutaneous transhepatic papillary balloon dilation in percutaneous choledochoscopic lithotomy. *Gastrointest Endosc* 2001; **54**: 232-236 [PMID: 11474400]
- 84 **Nagashima I**, Takada T, Shiratori M, Inaba T, Okinaga K.

- Percutaneous transhepatic papillary balloon dilation as a therapeutic option for choledocholithiasis. *J Hepatobiliary Pancreat Surg* 2004; **11**: 252-254 [PMID: 15368109 DOI: 10.1007/s00534-003-0851-x]
- 85 **Minami A**, Nakatsu T, Uchida N, Hirabayashi S, Fukuma H, Morshed SA, Nishioka M. Papillary dilation vs sphincterotomy in endoscopic removal of bile duct stones. A randomized trial with manometric function. *Dig Dis Sci* 1995; **40**: 2550-2554 [PMID: 8536511]
- 86 **Mathuna PM**, White P, Clarke E, Merriman R, Lennon JR, Crowe J. Endoscopic balloon sphincteroplasty (papillary dilation) for bile duct stones: efficacy, safety, and follow-up in 100 patients. *Gastrointest Endosc* 1995; **42**: 468-474 [PMID: 8566640]
- 87 **Yasuda I**, Tomita E, Moriwaki H, Kato T, Wakahara T, Sugihara J, Nagura K, Nishigaki Y, Sugiyama A, Enya M. Endoscopic papillary balloon dilatation for common bile duct stones: efficacy of combination with extracorporeal shock-wave lithotripsy for large stones. *Eur J Gastroenterol Hepatol* 1998; **10**: 1045-1050 [PMID: 9895052]
- 88 **Ochi Y**, Mukawa K, Kiyosawa K, Akamatsu T. Comparing the treatment outcomes of endoscopic papillary dilation and endoscopic sphincterotomy for removal of bile duct stones. *J Gastroenterol Hepatol* 1999; **14**: 90-96 [PMID: 10029284]
- 89 **Natsui M**, Narisawa R, Motoyama H, Hayashi S, Seki K, Wakabayashi H, Itoh S, Asakura H. What is an appropriate indication for endoscopic papillary balloon dilation? *Eur J Gastroenterol Hepatol* 2002; **14**: 635-640 [PMID: 12072597]
- 90 **Lin CK**, Lai KH, Chan HH, Tsai WL, Wang EM, Wei MC, Fu MT, Lo CC, Hsu PI, Lo GH. Endoscopic balloon dilatation is a safe method in the management of common bile duct stones. *Dig Liver Dis* 2004; **36**: 68-72 [PMID: 14971818 DOI: 10.1016/j.dld.2003.09.014]
- 91 **Tanaka S**, Sawayama T, Yoshioka T. Endoscopic papillary balloon dilation and endoscopic sphincterotomy for bile duct stones: long-term outcomes in a prospective randomized controlled trial. *Gastrointest Endosc* 2004; **59**: 614-618 [PMID: 15114302]
- 92 **Toda N**, Saito K, Wada R, Kawabe T, Shiratori Y, Mitsushima T, Omata M. Endoscopic sphincterotomy and papillary balloon dilation for bile duct stones. *Hepatogastroenterology* 2005; **52**: 700-704 [PMID: 15966186]
- 93 **Nakagawa H**, Ohara K. Safeguards against acute pancreatitis associated with endoscopic papillary balloon dilatation. *J Hepatobiliary Pancreat Surg* 2006; **13**: 75-79 [PMID: 16547665 DOI: 10.1007/s00534-005-1061-5]
- 94 **Liao WC**, Huang SP, Wu MS, Lin JT, Wang HP. Comparison of endoscopic papillary balloon dilatation and sphincterotomy for lithotripsy in difficult sphincterotomy. *J Clin Gastroenterol* 2008; **42**: 295-299 [PMID: 18223494 DOI: 10.1097/MCG.0b013e31802c3458]
- 95 **Natsui M**, Honma T, Genda T, Nakadaira H. Effects of endoscopic papillary balloon dilation and endoscopic sphincterotomy on bacterial contamination of the biliary tract. *Eur J Gastroenterol Hepatol* 2011; **23**: 818-824 [PMID: 21730870 DOI: 10.1097/MEG.0b013e328348c0bf]
- 96 **Kuo CM**, Chiu YC, Changchien CS, Tai WC, Chuah SK, Hu TH, Kuo YH, Kuo CH. Endoscopic papillary balloon dilation for removal of bile duct stones: evaluation of outcomes and complications in 298 patients. *J Clin Gastroenterol* 2012; **46**: 860-864 [PMID: 23060218 DOI: 10.1097/MCG.0b013e3182617a42]
- 97 **Seo YR**, Moon JH, Choi HJ, Kim DC, Ha JS, Lee TH, Cha SW, Cho YD, Park SH, Kim SJ. Comparison of endoscopic papillary balloon dilation and sphincterotomy in young patients with CBD stones and gallstones. *Dig Dis Sci* 2014; **59**: 1042-1047 [PMID: 24287639 DOI: 10.1007/s10620-013-2949-6]
- 98 **Lee D**, Lee B, Hwhang S, Baik Y, Lee S. Endoscopic papillary large balloon dilation after endoscopic sphincterotomy for treatment of large common bile duct stone. *Digest Endosc* 2007; **19**: S52-S56
- 99 **Kim MK**, Kim MH, Lee TY, Oh HC, Kwon SH, Han JH, Choi HO, Park SJ, Kim TH, Lee SS, Seo DW, Lee SK. Combined endoscopic sphincterotomy and large balloon sphincteroplasty for bile duct stones. *Korean J Med* 2007; **73**: 474-480
- 100 **Lee SH**, Hong SW, Cho YD, Cheon YK, Kim SG, Jang JY, Kim YS, Moon JH, Lee JS, Lee MS, Shim CS, Kim BS. The Safety and Effectiveness of Medium Endoscopic Sphincterotomy with Endoscopic Papillary Large Balloon Dilation for Removing Difficult Common Bile Duct Stones. *Korean J Gastrointest Endosc* 2007; **35**: 80-86
- 101 **Misra SP**, Dwivedi M. Large-diameter balloon dilation after endoscopic sphincterotomy for removal of difficult bile duct stones. *Endoscopy* 2008; **40**: 209-213 [PMID: 18264886 DOI: 10.1055/s-2007-967040]
- 102 **Espinel J**, Pinedo E. [Large balloon dilation for removal of bile duct stones]. *Rev Esp Enferm Dig* 2008; **100**: 632-636 [PMID: 19119789]
- 103 **Itoi T**, Itokawa F, Sofuni A, Kurihara T, Tsuchiya T, Ishii K, Tsuji S, Ikeuchi N, Moriyasu F. Endoscopic sphincterotomy combined with large balloon dilation can reduce the procedure time and fluoroscopy time for removal of large bile duct stones. *Am J Gastroenterol* 2009; **104**: 560-565 [PMID: 19174779 DOI: 10.1038/ajg.2008.67]
- 104 **Kim HG**, Cheon YK, Cho YD, Moon JH, Park do H, Lee TH, Choi HJ, Park SH, Lee JS, Lee MS. Small sphincterotomy combined with endoscopic papillary large balloon dilation versus sphincterotomy. *World J Gastroenterol* 2009; **15**: 4298-4304 [PMID: 19750573]
- 105 **Itoi T**, Sofuni A, Itokawa F, Kurihara T, Tsuchiya T, Ishii K, Tsuji S, Ikeuchi N, Umeda J, Moriyasu F. New large-diameter balloon-equipped sphincterotomy for removal of large bile duct stones (with videos). *Gastrointest Endosc* 2010; **72**: 825-830 [PMID: 20883862 DOI: 10.1016/j.gie.2010.06.018]
- 106 **Kurita A**, Maguchi H, Takahashi K, Katanuma A, Osanai M. Large balloon dilation for the treatment of recurrent bile duct stones in patients with previous endoscopic sphincterotomy: preliminary results. *Scand J Gastroenterol* 2010; **45**: 1242-1247 [PMID: 20521873 DOI: 10.3109/00365521.2010.495420]
- 107 **Ghazanfar S**, Qureshi S, Leghari A, Taj MA, Niaz SK, Quraishy MS. Endoscopic balloon sphincteroplasty as an adjunct to endoscopic sphincterotomy in removing large and difficult bile duct stones. *J Pak Med Assoc* 2010; **60**: 1039-1042 [PMID: 21381560]
- 108 **Kim KO**, Kim TN, Lee SH. Endoscopic papillary large balloon dilation for the treatment of recurrent bile duct stones in patients with prior sphincterotomy. *J Gastroenterol* 2010; **45**: 1283-1288 [PMID: 20635102 DOI: 10.1007/s00535-010-0284-7]
- 109 **Kim TH**, Oh HJ, Lee JY, Sohn YW. Can a small endoscopic sphincterotomy plus a large-balloon dilation reduce the use of mechanical lithotripsy in patients with large bile duct stones? *Surg Endosc* 2011; **25**: 3330-3337 [PMID: 21533521 DOI: 10.1007/s00464-011-1720-3]
- 110 **Sakai Y**, Tsuyuguchi T, Sugiyama H, Nishikawa T, Kurosawa J, Saito M, Tawada K, Mikata R, Tada M, Ishihara T, Yokosuka O. Endoscopic sphincterotomy combined with large balloon dilation for removal of large bile duct stones. *Hepatogastroenterology* 2013; **60**: 58-64 [PMID: 22641109 DOI: 10.5754/hge12351]
- 111 **Yang XM**, Hu B, Pan YM, Gao DJ, Wang TT, Wu J, Ye X. Endoscopic papillary large-balloon dilation following limited sphincterotomy for the removal of refractory bile duct stones: experience of 169 cases in a single Chinese center. *J Dig Dis* 2013; **14**: 125-131 [PMID: 23167553 DOI: 10.1111/1751-2980.12013]
- 112 **Yoon HG**, Moon JH, Choi HJ, Kim DC, Kang MS, Lee TH, Cha SW, Cho YD, Park SH, Kim SJ. Endoscopic papillary large balloon dilation for the management of recurrent dif-

ficult bile duct stones after previous endoscopic sphincterotomy. *Dig Endosc* 2014; **26**: 259-263 [PMID: 23581623 DOI: 10.1111/den.12102]

- 113 **Harada R**, Maguchi H, Takahashi K, Katanuma A, Osanai M, Yane K, Hashigo S, Kaneko M, Katoh R, Katoh S. Large

balloon dilation for the treatment of recurrent bile duct stones prevents short-term recurrence in patients with previous endoscopic sphincterotomy. *J Hepatobiliary Pancreat Sci* 2013; **20**: 498-503 [PMID: 23361432 DOI: 10.1007/s00534-012-0579-6]

P- Reviewer: Fujii L, Kawakami H, Kawakubo K
S- Editor: Gou SX **L- Editor:** A **E- Editor:** Wang CH





Published by **Baishideng Publishing Group Inc**
8226 Regency Drive, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgooffice@wjgnet.com
Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>
<http://www.wjgnet.com>



ISSN 1007-9327

