



## OPEN

SUBJECT AREAS:  
GENETICS RESEARCH  
RISK FACTORSReceived  
2 June 2014Accepted  
15 July 2014Published  
7 August 2014Correspondence and  
requests for materials  
should be addressed to  
M.U. (mueta@koto.  
kpu-m.ac.jp)

# Trans-ethnic study confirmed independent associations of *HLA-A\*02:06* and *HLA-B\*44:03* with cold medicine-related Stevens-Johnson syndrome with severe ocular surface complications

Mayumi Ueta<sup>1,2</sup>, Chitra Kannabiran<sup>3</sup>, Tais Hitomi Wakamatsu<sup>4</sup>, Mee Kum Kim<sup>5</sup>, Kyung-Chul Yoon<sup>6</sup>, Kyoung Yul Seo<sup>7</sup>, Choun-Ki Joo<sup>8</sup>, Virender Sangwan<sup>9</sup>, Varsha Rathi<sup>9</sup>, Sayan Basu<sup>9</sup>, Almas Shamaila<sup>3</sup>, Hyo Seok Lee<sup>6</sup>, Sangchul Yoon<sup>7</sup>, Chie Sotozono<sup>1</sup>, José Álvaro Pereira Gomes<sup>4</sup>, Katsushi Tokunaga<sup>10</sup> & Shigeru Kinoshita<sup>1</sup>

<sup>1</sup>Department of Ophthalmology, Kyoto Prefectural University of Medicine, Kyoto, Japan, <sup>2</sup>Research Center for Inflammation and Regenerative Medicine, Faculty of Life and Medical Sciences, Doshisha University, Kyoto, Japan, <sup>3</sup>Prof Brien Holden Eye Research Centre, L V Prasad Eye Institute, Hyderabad, India, <sup>4</sup>Department of Ophthalmology, Federal University of São Paulo, São Paulo, Brazil, <sup>5</sup>Department of Ophthalmology, Seoul National University College of Medicine, Seoul, Korea, <sup>6</sup>Department of Ophthalmology, Chonnam National University, Gwangju, Korea, <sup>7</sup>Department of Ophthalmology, Severance Hospital, Institute of Vision Research, Yonsei University College of Medicine, Seoul, Korea, <sup>8</sup>Department of Ophthalmology & Visual Science, Seoul St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Seoul, Korea, <sup>9</sup>Cornea and Anterior Segment Services, L V Prasad Eye Institute, Hyderabad, India, <sup>10</sup>Department of Human Genetics, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan.

Stevens-Johnson syndrome (SJS) and its severe variant, toxic epidermal necrolysis (TEN), are acute inflammatory vesiculobullous reactions of the skin and mucous membranes. Cold medicines including non-steroidal anti-inflammatory drugs and multi-ingredient cold medications are reported to be important inciting drugs. Recently, we reported that cold medicine related SJS/TEN (CM-SJS/TEN) with severe mucosal involvement including severe ocular surface complications (SOC) is associated with *HLA-A\*02:06* and *HLA-B\*44:03* in the Japanese. In this study, to determine whether *HLA-B\*44:03* is a common risk factor for CM-SJS/TEN with SOC in different ethnic groups we used samples from Indian, Brazilian, and Korean patients with CM-SJS/TEN with SOC, and investigated the association between CM-SJS/TEN with SOC and *HLA-B\*44:03* and/or *HLA-A\*02:06*. We found that *HLA-B\*44:03* was significantly associated with CM-SJS/TEN with SOC in the Indian and Brazilian but not the Korean population, and that *HLA-A\*02:06* might be weakly associated in the Korean- but not the Indian and Brazilian population.

Stevens-Johnson syndrome (SJS) and its severe variant, toxic epidermal necrolysis (TEN) with spots, are acute inflammatory vesiculobullous reactions of the skin and mucous membranes such as the ocular surface, oral cavity, and genitals. They are rare but often associated with inciting drugs and/or infectious agents<sup>1-3</sup>.

The association between human leukocyte antigen (HLA) genotypes and drug-induced severe cutaneous adverse reactions (SCARs) including SJS/TEN has been reported. There was a strong association between *HLA-B\*58:01* and SCARs, including SJS/TEN and the drug-induced hypersensitivity syndrome (DIHS), induced by the uric acid lowering drug allopurinol. This association was observed in Han Chinese<sup>4</sup>, Caucasian<sup>5</sup>, and Japanese patients<sup>6</sup>, suggesting that different ethnic groups share the same risk factor(s) for allopurinol-induced SCARs. *HLA-B\*15:02* exhibited a very strong association with carbamazepine-



Table 1 | Results of association analyses in patients with CM-SJS/TEN with SOC

ethnic group	HLA genotype	Carrier frequency (%)				Dominant model analysis				Gene frequency (%)				Dominant model analysis							
		CM-SJS/TEN with SOC		Control		P	Pc	Odds ratio (95% CI)	CM-SJS/TEN with SOC	Control		P	Pc	Odds ratio (95% CI)	CM-SJS/TEN with SOC	Control		P	Pc	Odds ratio (95% CI)	
		CM-SJS/TEN with SOC	Control	CM-SJS/TEN with SOC	Control				CM-SJS/TEN with SOC	Control				CM-SJS/TEN with SOC	Control						
Indian	A*02:06	1/20 (5.0%)	3/55 (5.5%)	0.938	-	0.91 (0.09–9.31)	1/39 (2.5%)	3/110 (2.7%)	0.939	-	0.91 (0.09–9.06)	1/39 (2.5%)	3/110 (2.7%)	0.939	-	0.91 (0.09–9.06)	1/39 (2.5%)	3/110 (2.7%)	0.939	-	0.91 (0.09–9.06)
	B*44:03	12/20 (60.0%)	6/55 (10.9%)	<b>1.07.E-05</b>	<b>2.14.E-05</b>	<b>12.25 (3.57–42.01)</b>	17/40 (42.5%)	7/110 (6.4%)	<b>9.37.E-08</b>	<b>1.87.E-07</b>	<b>10.88 (4.04–29.3)</b>	17/40 (42.5%)	7/110 (6.4%)	<b>9.37.E-08</b>	<b>1.87.E-07</b>	<b>10.88 (4.04–29.3)</b>	17/40 (42.5%)	7/110 (6.4%)	<b>9.37.E-08</b>	<b>1.87.E-07</b>	<b>10.88 (4.04–29.3)</b>
Brazilian	A*02:06	0/39 (0.00%)	0/134 (0.00%)	-	-	-	0/78 (0.00%)	0/268 (0.00%)	-	-	-	0/78 (0.00%)	0/268 (0.00%)	-	-	-	0/78 (0.00%)	0/268 (0.00%)	-	-	-
	B*44:03	10/39 (25.6%)	15/134 (11.2%)	<b>0.0239</b>	<b>0.0478</b>	<b>2.74 (1.12–6.71)</b>	11/78 (14.1%)	15/268 (5.60%)	<b>0.0121</b>	<b>0.0242</b>	<b>2.77 (1.22–6.31)</b>	11/78 (14.1%)	15/268 (5.60%)	<b>0.0121</b>	<b>0.0242</b>	<b>2.77 (1.22–6.31)</b>	11/78 (14.1%)	15/268 (5.60%)	<b>0.0121</b>	<b>0.0242</b>	<b>2.77 (1.22–6.31)</b>
Korean	A*02:06	11/31 (35.5%)	14/90 (15.6%)	<b>0.0181</b>	<b>0.0362</b>	<b>3.00 (1.18–7.57)</b>	12/62 (19.4%)	16/180 (8.9%)	<b>0.0263</b>	<b>0.0526</b>	<b>2.46 (1.09–5.54)</b>	12/62 (19.4%)	16/180 (8.9%)	<b>0.0263</b>	<b>0.0526</b>	<b>2.46 (1.09–5.54)</b>	12/62 (19.4%)	16/180 (8.9%)	<b>0.0263</b>	<b>0.0526</b>	<b>2.46 (1.09–5.54)</b>
	B*44:03	6/31 (19.4%)	18/90 (20.0%)	0.938	-	0.96 (0.34–2.69)	7/62 (11.3%)	19/180 (10.6%)	0.872	-	1.07 (0.43–2.70)	7/62 (11.3%)	19/180 (10.6%)	0.872	-	1.07 (0.43–2.70)	7/62 (11.3%)	19/180 (10.6%)	0.872	-	1.07 (0.43–2.70)

P: P values obtained with the  $\chi^2$ -test (Pearson), CI: Confidence interval.

Pc: P values corrected for the multiplicity of testing by the number of comparisons 2 (HLA-A\*02:06 + HLA-B\*44:03).

CM-SJS/TEN: cold medicine-related SJS/TEN, SOC: severe ocular surface complications.

induced SJS/TEN in Taiwanese Han Chinese patients<sup>7</sup> and *HLA-A\*31:01* was strongly associated with carbamazepine-induced SCARs including SJS/TEN in Japanese<sup>8</sup> and European patients<sup>9</sup>. We recently reported that cold medicine-related SJS/TEN with severe mucosal involvement including severe ocular surface complications (SOC) is associated with *HLA-A\*02:06* and *HLA-B\*44:03* in Japanese patients<sup>10</sup>.

The ophthalmologists Mondino et al.<sup>11</sup> and the dermatologists Roujeau et al.<sup>12,13</sup> reported that *HLA-B12* (*HLA-Bw44*) was significantly increased in Caucasian SJS patients many of whom developed SJS/TEN after taking non-steroidal anti-inflammatory drugs (NSAIDs). *HLA-B12* is primarily coded by *HLA-B\*44:02* or *HLA-B\*44:03* (<http://www.allele frequencies.net/>). The significant association between *HLA-B12* and SJS/TEN in Caucasian patients may be attributable to their genetic background.

To determine whether *HLA-B\*44:03* is a common risk factor for CM-SJS/TEN with SOC in different ethnic groups we used samples from Indian, Brazilian, and Korean patients with CM-SJS/TEN with SOC, and investigated the association between CM-SJS/TEN with SOC and *HLA-B\*44:03* and/or *HLA-A\*02:06*.

## Methods

Our study was approved by the institutional review boards of the participating institutions. All experimental procedures were conducted in accordance with the principles of the Helsinki Declaration. The purpose of the research and the experimental protocols were explained to all participants, and their prior written informed consent was obtained.

**Patients and controls.** Ophthalmologists diagnosed SJS/TEN based on a confirmed history of acute-onset high fever, serious mucocutaneous illness with skin eruptions, and involvement of at least two mucosal sites including the ocular surface<sup>14,15</sup>. They defined patients with SOC as those who manifested a pseudomembrane and an epithelial defect on the ocular surface in the acute stage, and as patients with ocular sequelae such as dry eye, trichiasis, symblepharon, and conjunctival invasion into the cornea in the chronic stage.

As in our previous study, we focused on SJS/TEN with SOC suspected of having been induced by cold medicines such as multi-ingredient cold medications and NSAIDs. As we found earlier that the genetic predisposition might be different between SJS/TEN with and without severe mucosal involvement including SOC<sup>10</sup> we focused on patients from different ethnic groups who presented with SJS/TEN with SOC.

Samples from Indian patients with CM-SJS/TEN were collected at the LV Prasad Eye Institute (n = 20; 12 males, 8 females; age range 7 to 63 years; median age 27.1 ± 13.4 (SD) years). Their age at onset ranged from 3 to 42 years (median age at onset, 19.2 ± 12.2 (SD) years; in 8 patients the age at onset was unknown). The drugs administered to these patients and the HLA type (A and B) of patients with CM-SJS/TEN with SOC are shown in Supplemental Table 1. The specific drug(s) were not known in all patients. Healthy volunteers (n = 55; 29 males, 26 females; median age 36.0 ± 11.6 years) served as the Indian controls.

Samples from Brazilian patients with CM-SJS/TEN were collected at the Federal University of Sao Paulo (n = 39, 15 males, 24 females; age range 13 to 69 years; median age, 37.1 ± 15.9 years; age range at onset, 3 to 69 years; median age at onset, 24.0 ± 17.2 years). The drugs administered, the ethnicity, and the HLA type (A and B) of these CM-SJS/TEN patients with SOC are shown in Supplemental Table 2. Healthy volunteers (n = 134; 55 males, 79 females; median age 41.2 ± 12.8 years) were the Brazilian controls (ethnicity: pardo, n = 66; white, n = 62; black, n = 4, Indian plus white, n = 2).

Samples from Korean patients with CM-SJS/TEN were collected at the Seoul National University College of Medicine, Chonnam National University, Yonsei University, and the Catholic University of Korea. There were 31 patients (12 males, 19 females) ranging in age from 4 to 71 years (median age 33.7 ± 19.0 years). Their age at SJS/TEN onset ranged from 3 to 63 years (median age at onset, 23.0 ± 16.1 years). The drugs used and the HLA type (A and B) of these patients with SOC are presented in Supplemental Table 3. The specific drug(s) were not known in all patients. Healthy volunteers (n = 90; 35 males, 55 females; median age 31.7 ± 7.9 years) were the Korean controls.

Samples from Indian subjects were obtained by extracting DNA from whole peripheral blood with the phenol chloroform method. For Brazilian samples, DNA was extracted from whole peripheral blood using the PAX gene blood DNA kit (Qiagen, Hilden, Germany) or from saliva using Oragene DNA (Kyodou International, Kanagawa, Japan). To obtain the samples from Korean subjects, DNA was extracted from whole peripheral blood using the PAXgene Blood DNA kit (Qiagen).

**HLA genotyping.** For the analysis of *HLA-A* and *HLA-B* we performed polymerase chain reaction (PCR) assays followed by hybridization with sequence-specific oligonucleotide probes using commercial bead-based typing kits (Wakunaga,



Hiroshima, Japan). Briefly, the target DNA was PCR-amplified with biotinylated primers specifically designed for amplified exons 2 and 3 of HLA-A, and -B genes. Then the PCR amplicon was denatured and hybridized to complementary oligonucleotide probes (72 probes for HLA-A, 93 probes for HLA-B) immobilized on fluorescent-coded microsphere beads. At the same time, the biotinylated PCR product was labeled with phycoerythrin-conjugated streptavidin and immediately examined with Luminex 100 (Luminex, Austin, TX, USA). Genotype determination and data analysis were performed automatically using the WAKFLOW typing software (Wakunaga, Hiroshima, Japan) according to the manufacturer's instructions.

**Statistical analysis.** We compared the carrier frequency and gene frequency of individual HLA alleles in the patients and controls with the  $\chi^2$ -test (Pearson) (JMP version 11 software; SAS Institute Japan Ltd., Tokyo, Japan).

## Results

**Strong association between HLA-B\*44:03 and CM-SJS/TEN with SOC in Indian patients.** We genotyped HLA-A and HLA-B in samples from Indian subjects (20 CM-SJS/TEN with SOC patients and 55 controls). Although the number of Indian subjects was small, we found a strong and significant association between their CM-SJS/TEN with SOC and HLA-B\*44:03 (carrier frequency:  $p = 1.07 \times 10^{-5}$ , odds ratio (OR) = 12.25, gene frequency:  $p = 9.37 \times 10^{-8}$ , OR = 10.88) but not HLA-A\*02:06 (Table 1).

**Significant association between HLA-B\*44:03 and CM-SJS/TEN with SOC in Brazilian patients.** Next we genotyped HLA-A and HLA-B in samples from Brazilian subjects (39 CM-SJS/TEN with SOC patients and 134 controls). Although the number of Brazilian subjects was small we found a significant association between Brazilian patients with CM-SJS/TEN with SOC and HLA-B\*44:03 (carrier frequency:  $p = 0.0239$ , OR = 2.74, gene frequency:  $p = 0.0121$ , OR = 2.77) but not HLA-A\*02:06 which is absent in the Brazilian population (Table 1). Interestingly, in Caucasians in the Brazilian samples (Brazilian Caucasian CM-SJS/TEN with SOC patients:  $n = 15$ , Brazilian Caucasian controls:  $n = 62$ ), the association with HLA-B\*44:03 was stronger (carrier frequency:  $p = 0.0037$ , OR = 6.22, gene frequency:  $p = 0.0011$ , OR = 5.99).

**Association between HLA-A\*02:06 and Korean patients with CM-SJS/TEN with SOC.** We also genotyped HLA-A and HLA-B in samples from Koreans (31 patients with CM-SJS/TEN with SOC and 90 controls). Although the number of Korean patients was small we found a significant association between patients with CM-SJS/TEN with SOC and HLA-A\*02:06 (carrier frequency:  $p = 0.0362$ , OR = 3.00, gene frequency:  $p = 0.0263$ , OR = 2.46) but not HLA-B\*44:03 (Table 1).

## Discussion

We previously reported that in the Japanese, CM-SJS/TEN with severe mucosal involvement including SOC was associated with HLA-A\*02:06 and HLA-B\*44:03<sup>10</sup>. In the present study we investigated whether the association with these alleles is shared by other ethnic groups. We found that HLA-B\*44:03 was strongly associated with CM-SJS/TEN with SOC in the Indian population which is genetically close to European populations<sup>16</sup> and significantly associated in the Brazilian population which is comprised of individuals with different ethnic backgrounds. There was no association between HLA-B\*44:03 and CM-SJS/TEN with SOC in the Korean population. HLA-A\*02:06 was weakly associated in the Korean population which is genetically close to the Japanese, but not in the Indian and Brazilian population.

HLA-B12 (HLA-Bw44) was significantly increased in Caucasian SJS patients many of whom developed SJS/TEN after taking NSAIDs<sup>11–13</sup>. Because HLA-B12 is primarily coded by HLA-B\*44:02 or HLA-B\*44:03 (<http://www.allelefrequencies.net/>), the significant association of HLA-B12 with SJS/TEN in Caucasian patients may be attributable to the association with the HLA-B\*44:03 genotype.

We also found that in Brazilian Caucasian patients with CM-SJS/TEN with SOC, the significant association with HLA-B\*44:03 was stronger than in the entire study population of Brazilians with CM-SJS/TEN with SOC. To determine whether HLA-B\*44:03 is a common marker for CM-SJS/TEN with SOC in Caucasian, HLA analysis of European patients with CM-SJS/TEN with SOC is needed.

Although HLA-A\*02:06 was strongly associated with the Japanese CM-SJS/TEN with SOC, and the Korean and Japanese population is genetically close<sup>16</sup>, in Korean patients CM-SJS/TEN with SOC was not strongly associated with HLA-A\*02:06. To determine whether HLA-A\*02:06 is a common marker for CM-SJS/TEN with SOC in East Asian populations further investigations using a larger number of samples are needed.

We also performed a meta-analysis by adding our previously-reported samples<sup>10</sup>. We used Cochran-Mantel-Haenszel statistics and found that both HLA-A\*02:06 and HLA-B\*44:03 are significantly associated with CM-SJS/TEN with SOC (Supplemental Table 4).

SCARs including SJS/TEN and DIHS induced by allopurinol were commonly and strongly associated with HLA-B\*58:01 in patients of different ethnic backgrounds including Han Chinese<sup>4</sup>, Caucasian<sup>5</sup>, and Japanese patients<sup>6</sup>. This observation suggests that different ethnic groups share the same risk factor(s) for allopurinol-induced SCARs.

With respect to carbamazepine-induced SJS/TEN, different HLA alleles are associated. HLA-B\*15:02 is associated in Taiwanese Han Chinese patients<sup>7</sup> and HLA-A\*31:01 in Japanese<sup>8</sup> and European patients<sup>9</sup>.

In CM-SJS/TEN with SOC, the associated alleles we identified are HLA-A\*02:06 in Japanese and Korean patients and HLA-B\*44:03 in Indian-, Brazilian-, and Japanese patients. Studies are underway to determine whether other HLA alleles are associated with CM-SJS/TEN with SOC in other populations.

1. Ueta, M. *et al.* Toll-like receptor 3 gene polymorphisms in Japanese patients with Stevens-Johnson syndrome. *Br J Ophthalmol* **91**, 962–965 (2007).
2. Yamane, Y., Aihara, M. & Ikezawa, Z. Analysis of Stevens-Johnson syndrome and toxic epidermal necrolysis in Japan from 2000 to 2006. *Allergol Int* **56**, 419–425 (2007).
3. Yetiv, J. Z., Bianchine, J. R. & Owen, J. A. Jr. Etiologic factors of the Stevens-Johnson syndrome. *South Med J* **73**, 599–602 (1980).
4. Hung, S. I. *et al.* HLA-B\*5801 allele as a genetic marker for severe cutaneous adverse reactions caused by allopurinol. *Proc Natl Acad Sci U S A* **102**, 4134–4139 (2005).
5. Lonjou, C. *et al.* A European study of HLA-B in Stevens-Johnson syndrome and toxic epidermal necrolysis related to five high-risk drugs. *Pharmacogenet Genomics* **18**, 99–107 (2008).
6. Tohkin, M. *et al.* A whole-genome association study of major determinants for allopurinol-related Stevens-Johnson syndrome and toxic epidermal necrolysis in Japanese patients. *Pharmacogenomics J* **13**, 60–69 (2013).
7. Chung, W. H. *et al.* Medical genetics: A marker for Stevens-Johnson syndrome. *Nature* **428**, 486 (2004).
8. Ozeki, T. *et al.* Genome-wide association study identifies HLA-A\*3101 allele as a genetic risk factor for carbamazepine-induced cutaneous adverse drug reactions in Japanese population. *Hum Molec Genet* **20**, 1034–1041 (2011).
9. McCormack, M. *et al.* HLA-A\*3101 and carbamazepine-induced hypersensitivity reactions in Europeans. *N Engl J Med* **364**, 1134–1143 (2011).
10. Ueta, M. *et al.* Independent strong association of HLA-A\*02:06 and HLA-B\*44:03 with cold medicine-related Stevens-Johnson syndrome with severe mucosal involvement. *Sci Rep* **4**, 4862 (2014).
11. Mondino, B. J., Brown, S. I. & Biglan, A. W. HLA antigens in Stevens-Johnson syndrome with ocular involvement. *Arch Ophthalmol* **100**, 1453–1454 (1982).
12. Roujeau, J. C. *et al.* HLA phenotypes and bullous cutaneous reactions to drugs. *Tissue Antigens* **28**, 251–254 (1986).
13. Roujeau, J. C. *et al.* Genetic susceptibility to toxic epidermal necrolysis. *Arch Dermatol* **123**, 1171–1173 (1987).
14. Ueta, M. *et al.* Association between prostaglandin E receptor 3 polymorphisms and Stevens-Johnson syndrome identified by means of a genome-wide association study. *J Allergy Clin Immunol* **126**, 1218–1225 e1210 (2010).
15. Ueta, M., Sotozono, C., Tokunaga, K., Yabe, T. & Kinoshita, S. Strong association between HLA-A\*0206 and Stevens-Johnson Syndrome in the Japanese. *Am J Ophthalmol* **143**, 367–368 (2007).



16. Abdulla, M. A. *et al.* Mapping human genetic diversity in Asia. *Science* **326**, 1541–1545 (2009).

## Acknowledgments

This work was conducted as part of the BioBank Japan Project supported by the Ministry of Education, Culture, Sports, Science and Technology of the Japanese government, and as part of the Promotion Project of Knowledge-Based Industrial Clustering of Okinawa Prefecture. It was supported in part by grants-in-aid for scientific research from the Japanese Ministry of Health, Labour and Welfare, a research grant from the Kyoto Foundation for the Promotion of Medical Science, and the Intramural Research Fund of Kyoto Prefectural University of Medicine. The funding agencies had no role in the study design, data collection and analysis, the decision to publish, or the preparation of the manuscript.

## Author contributions

M.U. wrote the main manuscript text and prepared the tables. M.U., C.K., T.W., M.K., K.Y., K.S., C.J., V.S., V.R., S.B., A.S., H.L., S.Y., C.S., J.G., K.T. and S.K. contributed to material of the research and reviewed the manuscript.

## Additional information

**Supplementary information** accompanies this paper at <http://www.nature.com/scientificreports>

**Competing financial interests:** The authors declare no competing financial interests.

**How to cite this article:** Ueta, M. *et al.* Trans-ethnic study confirmed independent associations of *HLA-A\*02:06* and *HLA-B\*44:03* with cold medicine-related Stevens-Johnson syndrome with severe ocular surface complications. *Sci. Rep.* **4**, 5981; DOI:10.1038/srep05981 (2014).



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>