

# The Effect of Low-Dose $\beta$ -Blocker on Heart Rate and Heart Rate Variability in Healthy Subjects with a Resting Heart Rate of Less than 65 Beats per Minute: Effect on the Image Quality of Prospective Electrocardiography-Gated Coronary CT Angiography<sup>1</sup>

## 저용량 베타차단제가 낮은 심박수 무증상 환자의 심박수, 심박변화, 그리고 전향적 동조화 심장 CT에 미치는 영향<sup>1</sup>

Chul Hwan Park, MD<sup>1</sup>, Sang Min Lee, MD<sup>2</sup>, Yoo Jin Hong, MD<sup>3</sup>, Tae Hoon Kim, MD<sup>1</sup>

<sup>1</sup>Department of Radiology and Research Institute of Radiological Science, Gangnam Severance Hospital, Yonsei University College of Medicine, Seoul, Korea

<sup>2</sup>Department of Radiology and Research Institute of Radiology, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea

<sup>3</sup>Department of Radiology and Research Institute of Radiological Science, Severance Hospital, Yonsei University College of Medicine, Seoul, Korea

**Purpose:** We assessed the effect of a low-dose  $\beta$ -blocker on heart rate (HR), HR variability (HRV) and image quality of prospective electrocardiography-gated coronary CT angiography (CCTA) in healthy subjects with low HR.

**Materials and Methods:** CCTA was performed with a 64-slice CT in 75 subjects with a HR of less than 65 beats per minute (bpm). Subjects were divided into 2 groups: Group 1 (G1), 35 with a low dose  $\beta$ -blocker; and Group 2 (G2), 40 without pre-medication. The image quality (IQ) of the CCTA was assessed on a 4-point grading scale (1, poor; 4, excellent).

**Results:** Initial HR (bpm) was not different between the 2 groups. HR during CCTA was lower in G1 than G2 ( $50.3 \pm 5.6$  vs.  $53.3 \pm 4.8$ ,  $p = 0.016$ ). HRV was not different between the 2 groups. Per-segment analysis showed better IQ at the mid-segment of the right coronary artery ( $3.0 \pm 0.9$  vs.  $2.5 \pm 1.1$ ,  $p = 0.039$ ) and the first diagonal branch ( $3.4 \pm 0.6$  vs.  $3.1 \pm 0.7$ ,  $p = 0.024$ ), in the G1 than the G2 group, respectively. The IQ was negatively correlated with HR, but no correlation was found between HRV and IQ. The IQs in the per-vessel analysis were not different between the 2 groups.

**Conclusion:** Low-dose  $\beta$ -blocker reduced HR and improved the IQ of CCTA in a few segments, even at a HR of less than 65 bpm. However the effect was limited.

### Index terms

Coronary CT Angiography  
Heart Rate  
Heart Rate Variability  
 $\beta$ -Blocker  
Image Quality

Received August 18, 2014; Accepted November 24, 2014

**Corresponding author:** Tae Hoon Kim, MD  
Department of Radiology and Research Institute of Radiological Science, Gangnam Severance Hospital, Yonsei University College of Medicine, 211 Eonju-ro, Gangnam-gu, Seoul 135-720, Korea.  
Tel. 82-2-2019-3510 Fax. 82-2-3462-5472  
E-mail: thkim1@yuhs.ac

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## INTRODUCTION

Coronary CT angiography (CCTA) with a multidetector row computed tomography (MDCT) scanner holds diagnostic value in coronary artery disease (1-3). Even though many factors can affect the image quality of CCTA, including heart rate (HR), heart rate variability (HRV), breathing, obesity, and coronary calcification (4-7), a relatively low and regular HR is essential to

optimal image quality (4). With advances in CT technology, 64-slice MDCT was designed to accommodate a wider range of HRs with improved temporal resolution; however, studies have shown that HRV during CT scanning has a strong influence on the image quality and can decrease the diagnostic accuracy of CCTA (4, 6).

$\beta$ -blockers reduce sympathetic influences and have been widely used to optimize HR in patients with tachycardia (5).

$\beta$ -blockers can amend HRV by stabilizing HR (6). Therefore, we assumed that administering a low dose  $\beta$ -blocker to patients with low HR could improve the image quality and diagnostic value of a prospective electrocardiography (ECG)-gated CCTA by reducing HR or HRV. We prospectively assessed the effectiveness of a low-dose  $\beta$ -blocker on HR, HRV and image quality of CCTA in subjects with a HR under 65 beats per minute (bpm), using a prospective ECG-triggered protocol (step-and-shoot technique) with a 64-slice MDCT scanner.

## MATERIALS AND METHODS

### Patient Preparation

The study was a prospective study that was approved by our Institutional Review Board. Written informed consent was obtained from all subjects. Healthy subjects with a HR of less than 65 bpm who underwent a prospective ECG-triggered CCTA using a 64-slice CT scanner to screen for coronary artery disease during the study period were enrolled. We excluded those who had a HR of greater than 65 bpm or less than 50 bpm before CT scan, history of coronary artery disease, known arrhythmia, previous allergic reaction to iodinated contrast media or  $\beta$ -blocker, renal insufficiency (serum creatinine > 150  $\mu$ mol/L), hemodynamic instability, congestive heart failure, asthma, chronic obstructive pulmonary disease, systolic blood pressure of less than 100 mm Hg, and those taking an oral  $\beta$ -blocker at baseline (5). Enrolled subjects were randomly assigned to 2 groups (G1 and G2). Group 1 received a low-dose of  $\beta$ -blocker (Atenolol; 12.5 mg, Hyundai Pharmaceutical Ind. Co., Seoul, Korea) 30 minutes before the CCTA examination. Group 2 underwent CCTA without premedication.

### CT Scan Protocol

All cardiac MDCT was performed with a 64-MDCT scanner (Philips Brilliance 64, Philips Medical System, Best, the Netherlands). Data acquisition was performed in the craniocaudal direction within a single breath-hold at end-inspiratory suspension. The scanning range covered the heart from the level of the carina to the diaphragm. A prospective ECG-triggered protocol was used with a step-and-shoot technique. The scanning parameters were as follows: step-and-shoot axial scanning direction, 420-msec gantry rotation time, 120 kV, 210 mAs, 64  $\times$

0.625-mm slice collimation, 4-cm table feed per rotation, and the center of the imaging window set at 70–80% of the R-R interval. Seventy mL of iodinated contrast agent (Optiray 350; Tyco Healthcare, Kantata, Canada) was administered intravenously into an antecubital vein through an 18 gauge catheter at a rate of 5 mL/sec followed by 50 mL of normal saline at a rate of 5 mL/sec using a power injector (Nemoto; Nemoto Kyorindo, Tokyo, Japan). Imaging was performed using a real-time bolus tracking technique. The scans were started 7 seconds after a trigger threshold of 130 Hounsfield units was reached at the proximal descending aorta. The breath-hold maneuver was successfully performed by all patients. An ECG for each patient was recorded simultaneously.

### CT Image Reconstruction

The image reconstruction was performed on the scanner's workstation using commercially available software (Extended Brilliance Workstation, Philips Medical System, Best, the Netherlands). The reconstruction parameters used were as follows: 0.9 mm slice thickness, 0.45 mm increment, 512  $\times$  512 pixels image matrix, XCC kernel, and 15–23 cm field of view. MDCT data were transferred and post processing was performed on an Aquarius Workstation V3.6 (TeraRecon, San Mateo, CA, USA). Multi-planar reformatted (MPR), curved-planar reformatted (CPR), and medial axis reformatted (MAR) images with orthogonal and perpendicular projections to the vessel courses of each coronary segment were generated for the evaluation.

### CT Image Analysis

Fifteen coronary segments were subdivided according to the American Heart Association guidelines (8); proximal right coronary artery (pRCA), middle RCA (mRCA), distal RCA (dRCA), right posterior descending artery (RPD), left main coronary artery, proximal left anterior descending artery (LAD), middle LAD, distal LAD, 1st diagonal branch (D1), 2nd diagonal branch, proximal left circumflex artery (pLCx), distal LCx (dLCx), obtuse marginal artery (OM), posterolateral artery and left posterior descending artery. The segments with a diameter of at least 1.5 mm at their origin were included.

Two radiologists assessed the image quality of 15 coronary segments in consensus. We evaluated all reconstructed images using axial source images, as well as MPR, CPR, and MAR im-

ages. The image quality was assessed using the following 4-point grading scale by the 2 radiologists who were blinded to the HR and HRV: grade 1 (poor), severe degree of image degradation or discontinuation of vessel contour that prevented vessel lumen evaluation; grade 2 (adequate), moderate degree of image degradation with some obstacle to vessel lumen evaluation; grade 3 (good), minor degree of image degradation without compromising vessel lumen evaluation, and grade 4 (excellent), no image degradation (Fig. 1). Any segment with grade 1 was regarded as non-diagnostic.

### Radiation Dose

The dose-length product was recorded for each cardiac CT examination by the CT scanner. The mean effective radiation dose was calculated by multiplying the dose-length product by the conversion coefficient (0.017) for the chest and expressed as millisieverts (9).

### Heart Rate and Heart Rate Variability Analyses

The initial HR ( $HR_{pre}$ ) prior to  $\beta$ -blocker administration, was measured 30 minutes before the CCTA exam after sufficient rest in all patients. HRs were also checked during data acquisition of CCTA. Mean HR during scanning ( $HR_{mean}$ ) was obtained from the ECG-gated data. HR change ( $HR_{chg}$ ) was defined as difference between  $HR_{pre}$  and  $HR_{mean}$  (i.e.,  $HR_{chg} = HR_{pre} - HR_{mean}$ ). HRV was calculated as 1 standard deviation (SD) of the HR dur-

ing CT scanning (10).


### Statistical Analysis

Continuous variables were expressed as mean  $\pm$  SD and categorical variables as frequencies or percentages. Student's *t*-test or Mann-Whitney U test was used to evaluate the statistical significance of differences for the demographic data, HR, HRV and image quality scores between the 2 groups. Paired *t*-test was used to evaluate the statistical significance of difference between  $HR_{pre}$  and  $HR_{mean}$  in each group. Chi-square test was used to evaluate the statistical significance of differences for the number of segments in each score between the 2 groups. A linear regression and Pearson correlation were used to determine the correlations between the mean image quality score and  $HR_{mean}$  or HRV. A *p* value of less than 0.05 was considered as statistically significant. All statistical analyses were performed with the commercially-available software (SPSS 20; Statistical Package for the Social Science, Chicago, IL, USA).

## RESULTS

### Heart Rate and Heart Rate Variability

CCTA was performed in all subjects without complications. Demographic data including HR and HRV were summarized in Table 1. Mean age and body mass index were not statistically different between the 2 groups ( $p > 0.05$ ).  $HR_{pre}$  was not different in




**Fig. 1.** 4-point grading scale of coronary artery image quality. The scores of grade 2 to 4 were regarded diagnostic and grade 1 was non-diagnostic.

- A.** Grade 1 (poor), severe degree of image degradation or discontinuation of vessel contour that prevented vessel lumen evaluation.
- B.** Grade 2 (adequate), moderate degree of image degradation with some obstacle to vessel lumen evaluation.
- C.** Grade 3 (good), minor degree of image degradation without affecting vessel lumen evaluation grade.
- D.** Grade 4 (excellent), no image degradation.

**Table 1. Demographic Data in 75 Patients with a Prospective ECG-Gated CCTA**

Characteristics	Group 1	Group 2	p Value
No. of patients	35	40	
Male to female ratio	28:7	25:15	0.097
Age (year)	61.3 $\pm$ 11.9	57.8 $\pm$ 11.1	0.849
Body mass index (kg/m <sup>2</sup> )	25.2 $\pm$ 3.3	24.1 $\pm$ 2.8	0.094
HR <sub>pre</sub> (beats/min)	60.4 $\pm$ 4.2	58.8 $\pm$ 3.6	0.08
HR <sub>chg</sub> (beats/min)	10.1 $\pm$ 5.4	5.5 $\pm$ 5.5	< 0.001
HR <sub>mean</sub> (beats/min)	50.3 $\pm$ 5.6	53.3 $\pm$ 4.8	0.016

Note.—CCTA = coronary CT angiography, ECG = electrocardiography, HR = heart rate, HR<sub>chg</sub> = HR difference between initial HR and HR during scanning (i.e., HR<sub>chg</sub> = HR<sub>pre</sub> - HR<sub>mean</sub>), HR<sub>mean</sub> = mean HR during data acquisition of CCTA, HR<sub>pre</sub> = initial heart rate measured 30 minutes before the CCTA



**Fig. 2.** HR<sub>pre</sub> and HR<sub>mean</sub> in group 1 and 2. Boxplots show HR<sub>pre</sub> (**A**) and HR<sub>mean</sub> (**B**) in subjects administered a low dose  $\beta$ -blocker (group 1) and those that were not (group 2). (**A**) Mean heart rates (HR) prior to administration of  $\beta$ -blocker (HR<sub>pre</sub>) are not statistically different between group 1 and group 2 ( $p = 0.08$ ). HR change (HR<sub>chg</sub>) was significantly larger in group 1 than in group 2 ( $p < 0.001$ ) and (**B**) mean HR during coronary CT angiography (HR<sub>mean</sub>) in group 1 was significantly lower than group 2 ( $p = 0.016$ ).

Note.—Box = 1st–3rd quartiles, Bold line = median, Whiskers = minimum and maximum values, o = outlier, HR change (HR<sub>chg</sub>) = difference between HR<sub>pre</sub> and HR<sub>mean</sub> (i.e., HR<sub>chg</sub> = HR<sub>pre</sub> - HR<sub>mean</sub>)

group 1 and group 2 (60.4  $\pm$  4.2 bpm vs. 58.8  $\pm$  3.6 bpm,  $p = 0.08$ ). HR<sub>mean</sub> was significantly lower than HR<sub>pre</sub> in group 1 (50.3  $\pm$  5.6 bpm vs. 60.4  $\pm$  4.2 bpm,  $p = 0.011$ ) but not in group 2 (53.3  $\pm$  4.8 bpm vs. 58.8  $\pm$  3.6 bpm,  $p = 0.193$ ). HR<sub>chg</sub> was significantly higher in group 1 than in group 2 (10.1  $\pm$  5.4 bpm vs. 5.5  $\pm$  5.5 bpm,  $p < 0.001$ ). HR<sub>mean</sub> during CCTA in group 1 was significantly lower than that of group 2 (50.3  $\pm$  5.6 bpm vs. 53.3  $\pm$  4.8 bpm,  $p = 0.016$ ) (Table 1, Fig. 2). During CCTA, HRV was not different in group 1 and group 2 (1.1  $\pm$  0.5 vs. 1.3  $\pm$  0.7,  $p = 0.147$ ) (Table 2, Fig. 3). The mean radiation dose of CCTA was 4.0  $\pm$  0.4 mSv (range, 3.7–4.6 mSv). The radiation dose of group 1 was not different with that of group 2 (3.9  $\pm$  0.5 mSv vs. 4.0  $\pm$  0.4 mSv,  $p = 0.98$ ).

### Image Quality

A total of 921 segments with a diameter greater than 1.5 mm were evaluated (431 segments in group 1 and 490 in group 2).

Group 1 was diagnostic in 429 coronary segments (99.3%); excellent (score 4) in 190 (44.1%), good (score 3) in 184 (42.7%), and adequate (score 2) in 54 (12.5%). Three segments (0.7%) were non-diagnostic (score 1) due to severe degradation of vessel contours. Group 2 was diagnostic in 474 (97.7%); excellent (score 4) in 192 (39.2%), good (score 3) in 216 (44.1%), and adequate (score 2) in 66 (13.5%). Sixteen segments (3.3%) were non-diagnostic (score 1). On the segment-based analysis, the image quality scores for mRCA and D1 segments were higher in group 1 than in group 2 (3.0  $\pm$  0.9 vs. 2.5  $\pm$  1.1,  $p = 0.039$  in mRCA, 3.4  $\pm$  0.6 vs. 3.1  $\pm$  0.7,  $p = 0.024$  in D1). However, the mean scores of image quality were not different between group 1 and group 2 (3.3  $\pm$  0.4 vs. 3.2  $\pm$  0.3,  $p = 0.199$ ), and the image quality scores on the vessel-based analysis were not different between group 1 and group 2 (Table 2). The mean image quality score of the coronary artery segments in both groups was nega-

Table 2. Heart Rate, HRV, and Image Quality Score in 75 Patients

	Group 1 (n = 35)	Group 2 (n = 40)	p Value
HR <sub>mean</sub> (beats/min)	50.3 ± 5.6	53.3 ± 4.8	0.016
HRV	1.1 ± 0.5	1.3 ± 0.7	0.147
Image quality: per segment analysis			
Mean of 15 segments	3.3 ± 0.4	3.2 ± 0.3	0.199
pRCA	3.5 ± 0.6	3.6 ± 0.6	0.400
mRCA	3.0 ± 0.9	2.5 ± 1.1	0.039
dRCA	3.4 ± 0.7	3.3 ± 0.8	0.787
RPD	3.5 ± 0.6	3.5 ± 0.6	0.132
LM	3.9 ± 0.3	3.9 ± 0.3	0.835
pLAD	3.7 ± 0.5	3.8 ± 0.5	0.567
mLAD	3.1 ± 0.7	3.1 ± 0.8	0.949
dLAD	2.8 ± 0.9	2.6 ± 0.8	0.459
D1	3.4 ± 0.6	3.1 ± 0.7	0.024
D2	3.3 ± 0.5	3.0 ± 0.6	0.115
pLCx	3.7 ± 0.5	3.5 ± 0.6	0.079
OM	3.1 ± 0.6	3.0 ± 0.5	0.664
dLCx	2.8 ± 0.7	2.8 ± 0.5	0.739
PL	3.1 ± 0.5	2.9 ± 0.4	0.334
LPD	3.3 ± 0.5	3.0 ± 0.5	0.250
Image quality: per vessel analysis			
RCA	3.3 ± 0.7	3.1 ± 0.9	0.262
LAD	3.2 ± 0.8	3.2 ± 0.8	0.835
LCx	3.3 ± 0.8	3.1 ± 0.7	0.268
No. of segments (%)	431 (100)	490 (100)	
No. of segments in each score (%)			
Grade 4 (excellent)	190 (44.1)	192 (39.2)	< 0.05
Grade 3 (good)	184 (42.7)	216 (44.1)	
Grade 2 (adequate)	54 (12.5)	66 (13.5)	
Grade 1 (poor)	3 (0.7)	16 (3.3)	

Note.—dLAD = distal LAD, dLCx = distal LCx, dRCA = distal RCA, D1 = 1st diagonal branch, D2 = 2nd diagonal branch, HR<sub>mean</sub> = mean HR during data acquisition of coronary CT angiography, HRV = heart rate variability, LAD = left anterior descending artery, LCx = left circumflex artery, LM = left main coronary artery, LPD = left posterior descending artery, mLAD = middle LAD, mRCA = middle RCA, OM = obtuse marginal artery, PL = posterolateral artery, pLAD = proximal LAD, pLCx = proximal LCx, pRCA = proximal RCA, RCA = right coronary artery, RPD = right posterior descending artery

tively correlated with HR<sub>mean</sub> ( $r = -0.449$ ,  $p < 0.001$ ), but no significant correlation was found between HRV and mean image quality ( $r = -0.013$ ,  $p = 0.913$ ) (Fig. 4). On the segment-based analysis, the image quality scores of the pRCA, mRCA, dRCA, RPD, D1, pLCx, OM, and dLCx were negatively correlated to HR<sub>mean</sub> with statistical significance ( $p < 0.05$ ). The correlation coefficient of mRCA was  $-0.575$  ( $p < 0.001$ ) and D1 was  $-0.297$  ( $p = 0.013$ ). None of the segments showed significant correlation between image quality and HRV (Table 3).

## DISCUSSION


The aim of this study was to evaluate the effect of a low-dose

β-blocker on HR and HRV, as well as the image quality of a prospective ECG-gated CCTA in healthy subjects with low HR. Our study demonstrated that the use of low dose β-blocker could improve the image quality of mRCA and D1 segments by reducing HR, even when the HR was less than 65 bpm.

The step-and-shoot technique using a prospective ECG-gated algorithm is one of the most powerful methods for radiation dose reduction. Dosage can be reduced up to 70–90% from that of retrospective ECG-gated CCTA (9, 11–13). However, a prospective ECG-gated CCTA mandates a low and regular HR of less than 60 to 65 bpm for optimal images (14). We postulated that the administration of low dose β-blocker could improve the image quality of a prospective ECG-gated CCTA by reducing



HR or HRV, even in low HR subjects. We administered 12.5 mg oral atenolol that is less than the usual dose of 50–100 mg depending on the patient’s weight for heart rate control (15). In group 1, the HR reduction ( $HR_{chg}$ ) was 11 bpm after low dose  $\beta$ -blocker administration. In group 2, the  $HR_{chg}$  was 5.5 bpm without premedication, probably due to the increased vagal tone by breath-holding (16). Zhang et al. (17) reported that breath-holding reduced the mean HR by about 4 bpm. Low dose

$\beta$ -blocker affected the image quality of CCTA. Non-diagnostic segments were lower in group 1 (0.7%) than in group 2 (3.3%). Image quality on segmental analysis was better in group 1 at the mRCA, which was known to be easily degraded by motion artifact (18), and D1 branch. However, image quality improvement was only observed in mRCA and D1 segments. Also, there was no difference in image quality between the 2 groups on vessel based analysis. Leschka et al. (4, 6) reported that 64-slice MDCT had reasonable image quality over a wide range of HRs, and HRV has more effect than the HR. Earls (19) reported that a small dose of  $\beta$ -blocker might be helpful in minimizing HRV in patients with low HR and high HRV. HRV was not related directly to the image quality in our study. These findings may be the result of low HR and small HRV in the study subjects. Low HR provides sufficient duration of diastole for good quality images, and the HRV may have been too small to significantly affect the image quality in patients with low HR (20). Furthermore, there was no difference of HRV between the 2 groups, hence the effect of low dose  $\beta$ -blocker on HRV was hard to compare. The radiation dose of our study was similar to those of previous studies with a prospective ECG-gating technique (9, 21). Lowering HR could increase the radiation dose of prospective ECG-gated CCTA by lengthening the R-R interval. However, the radiation dose in group 1 and 2 were not significantly different.



**Fig. 3.** Heart rate variability (HRV) in group 1 and 2. Boxplots show HRV in subjects with low dose  $\beta$ -blocker (group 1) and without  $\beta$ -blocker (group 2). During coronary CT angiography scanning, HRV was not different in group 1 and group 2 ( $p = 0.147$ ). Note.—Box = 1st–3rd quartiles, Bold line = median, Whiskers = minimum and maximum values, o = outlier

There were several limitations to the study. First, the number



**Fig. 4.** Linear regression plots of mean image quality scores over the mean heart rate and heart rate variability (HRV) in 75 patients.  
**A.** The mean image quality score of the coronary artery segments in both groups is negatively correlated with mean heart rate ( $r = -0.449, p < 0.001$ ).  
**B.** The mean image quality scores for all coronary artery segments and HRV shows no significant correlation ( $r = -0.013, p = 0.913$ ). Dotted lines represent 95% confidence limits.

Table 3. Correlation between Image Quality of Each Segment and Heart Rate/Heart Rate Variability

	Heart Rate		Heart Rate Variability	
	Correlation Coefficient	p Value	Correlation Coefficient	p Value
pRCA	-0.235	0.043	-0.132	0.260
mRCA	-0.575	< 0.001	-0.058	0.621
dRCA	-0.495	< 0.001	-0.65	0.172
RPD	-0.451	0.001	0.104	0.463
LM	0.012	0.916	0.166	0.154
pLAD	-0.135	0.249	0.041	0.729
mLAD	0.070	0.553	0.076	0.516
dLAD	-0.223	0.055	-0.027	0.817
D1	-0.297	0.013	-0.039	0.750
D2	-0.068	0.696	0.010	0.934
pLCx	-0.258	0.026	0.002	0.987
OM	-0.439	< 0.001	0.002	0.987
dLCx	-0.340	0.004	-0.104	0.392
PL	-0.325	0.188	0.243	0.331
LPD	0.055	0.931	0.655	0.230

Note.—dLAD = distal LAD, dLCx = distal LCx, dRCA = distal RCA, D1 = 1st diagonal branch, D2 = 2nd diagonal branch, LAD = left anterior descending artery, LCx = left circumflex artery, LM = left main coronary artery, LPD = left posterior descending artery, mLAD = middle LAD, mRCA = middle RCA, OM = obtuse marginal artery, PL = posterolateral artery, pLAD = proximal LAD, pLCx = proximal LCx, pRCA = proximal RCA, RCA = right coronary artery, RPD = right posterior descending artery

of enrolled subjects was too small to be generalized to all patients. Second, the comparative study was not performed in the same patient groups due to the concern of radiation exposure. Nevertheless, there was no significant difference in demographic data and HRs prior to the administration of  $\beta$ -blocker between the 2 groups. Third, we evaluated the image quality based on motion artifact but not the diagnostic performance for detecting coronary artery disease. A future study has been planned to analyze the diagnostic accuracy of a prospective ECG-gated CCTA under the effect of a low dose  $\beta$ -blocker.

In conclusion, low-dose  $\beta$ -blocker could reduce the HR, including resting HRs of less than 65 bpm that are not usually indicated for  $\beta$ -blocker. The image quality of CCTA using a prospective gating technique was improved in the mRCA and D1 segments with a low-dose  $\beta$ -blocker by reducing HR. However, on the per-vessel analysis, the overall image quality was not changed. Therefore, usefulness of a low-dose  $\beta$ -blocker appears to be limited in healthy subjects with low HR.

## REFERENCES

1. Raff GL, Gallagher MJ, O'Neill WW, Goldstein JA. Diagnostic accuracy of noninvasive coronary angiography using 64-slice spiral computed tomography. *J Am Coll Cardiol* 2005;46:552-557
2. Mollet NR, Cademartiri F, van Mieghem CA, Runza G, McFadden EP, Baks T, et al. High-resolution spiral computed tomography coronary angiography in patients referred for diagnostic conventional coronary angiography. *Circulation* 2005;112:2318-2323
3. Martuscelli E, Romagnoli A, D'Eliseo A, Razzini C, Tomassini M, Sperandio M, et al. Accuracy of thin-slice computed tomography in the detection of coronary stenoses. *Eur Heart J* 2004;25:1043-1048
4. Leschka S, Scheffel H, Husmann L, Gämperli O, Marincek B, Kaufmann PA, et al. Effect of decrease in heart rate variability on the diagnostic accuracy of 64-MDCT coronary angiography. *AJR Am J Roentgenol* 2008;190:1583-1590
5. Pannu HK, Alvarez W Jr, Fishman EK. Beta-blockers for cardiac CT: a primer for the radiologist. *AJR Am J Roentgenol* 2006;186(6 Suppl 2):S341-S345
6. Leschka S, Wildermuth S, Boehm T, Desbiolles L, Husmann L, Plass A, et al. Noninvasive coronary angiography with 64-section CT: effect of average heart rate and heart rate variability on image quality. *Radiology* 2006;241:378-385
7. Brodoefel H, Reimann A, Burgstahler C, Schumacher F, Herberts T, Tsiflikas I, et al. Noninvasive coronary angiography using 64-slice spiral computed tomography in an

- unselected patient collective: effect of heart rate, heart rate variability and coronary calcifications on image quality and diagnostic accuracy. *Eur J Radiol* 2008;66:134-141
8. Austen WG, Edwards JE, Frye RL, Gensini GG, Gott VL, Griffith LS, et al. A reporting system on patients evaluated for coronary artery disease. Report of the Ad Hoc Committee for Grading of Coronary Artery Disease, Council on Cardiovascular Surgery, American Heart Association. *Circulation* 1975;51(4 Suppl):5-40
  9. Shuman WP, Branch KR, May JM, Mitsumori LM, Lockhart DW, Dubinsky TJ, et al. Prospective versus retrospective ECG gating for 64-detector CT of the coronary arteries: comparison of image quality and patient radiation dose. *Radiology* 2008;248:431-437
  10. Brodoefel H, Burgstahler C, Tsiflikas I, Reimann A, Schroeder S, Claussen CD, et al. Dual-source CT: effect of heart rate, heart rate variability, and calcification on image quality and diagnostic accuracy. *Radiology* 2008;247:346-355
  11. Husmann L, Valenta I, Gaemperli O, Adda O, Treyer V, Wyss CA, et al. Feasibility of low-dose coronary CT angiography: first experience with prospective ECG-gating. *Eur Heart J* 2008;29:191-197
  12. Scheffel H, Alkadhi H, Leschka S, Plass A, Desbiolles L, Guber I, et al. Low-dose CT coronary angiography in the step-and-shoot mode: diagnostic performance. *Heart* 2008;94:1132-1137
  13. Earls JP, Berman EL, Urban BA, Curry CA, Lane JL, Jennings RS, et al. Prospectively gated transverse coronary CT angiography versus retrospectively gated helical technique: improved image quality and reduced radiation dose. *Radiology* 2008;246:742-753
  14. Jeong DW, Choo KS, Baik SK, Kim YW, Jeon UB, Kim JS, et al. Step-and-shoot prospectively ECG-gated versus retrospectively ECG-gated with tube current modulation coronary CT angiography using the 128-slice MDCT: comparison of image quality and radiation dose. *Acta Radiol* 2011;52:155-160
  15. College of Physicians and College of Radiologists Joint Writing Committee, Thye HK, Han KB, Chen K, Chua K, Kwok R, et al. Guidelines on cardiac CT in Singapore (2006). *Ann Acad Med Singapore* 2006;35:287-296
  16. Hanly PJ, George CF, Millar TW, Kryger MH. Heart rate response to breath-hold, valsalva and Mueller maneuvers in obstructive sleep apnea. *Chest* 1989;95:735-739
  17. Zhang SZ, Hu XH, Zhang QW, Huang WX. Evaluation of computed tomography coronary angiography in patients with a high heart rate using 16-slice spiral computed tomography with 0.37-s gantry rotation time. *Eur Radiol* 2005;15:1105-1109
  18. Lu B, Mao SS, Zhuang N, Bakhsheshi H, Yamamoto H, Takasu J, et al. Coronary artery motion during the cardiac cycle and optimal ECG triggering for coronary artery imaging. *Invest Radiol* 2001;36:250-256
  19. Earls JP. How to use a prospective gated technique for cardiac CT. *J Cardiovasc Comput Tomogr* 2009;3:45-51
  20. Herzog BA, Husmann L, Burkhard N, Valenta I, Gaemperli O, Tatsugami F, et al. Low-dose CT coronary angiography using prospective ECG-triggering: impact of mean heart rate and heart rate variability on image quality. *Acad Radiol* 2009;16:15-21
  21. Hong YJ, Kim SJ, Lee SM, Min PK, Yoon YW, Lee BK, et al. Low-dose coronary computed tomography angiography using prospective ECG-triggering compared to invasive coronary angiography. *Int J Cardiovasc Imaging* 2011;27:425-431

## 저용량 베타차단제가 낮은 심박수 무증상 환자의 심박수, 심박변화, 그리고 전향적 동조화 심장 CT에 미치는 영향<sup>1</sup>

박철환<sup>1</sup> · 이상민<sup>2</sup> · 홍유진<sup>3</sup> · 김태훈<sup>1</sup>

**목적:** 저용량 베타차단제가 분당 65회 이하의 느린 심박수 무증상 환자에서, 전향적 동조화 심장 CT의 영상 질을 높일 수 있는지 알아본다.

**대상과 방법:** 심박수가 65회 이하인 무증상 환자 75명을 두 군으로 나누어 전향적 동조화 심장 CT를 시행하였다. 1군 35명은 검사 30분 전, 저용량 베타차단제를 구강투여 후 시행하였으며, 2군 40명은 투약 없이 시행하였다. 각 환자에서 검사 전 심박수, 검사 중 심박수, 검사 중 심박변화를 측정하였다. 심장 CT 관상동맥 각 분절마다 영상 질을 평가하여 1점에서 4점까지 점수를 매겼으며, 4점이 가장 좋은 영상을 의미하였다.

**결과:** 두 군의 검사 전 심박수는 차이가 없었다. 심장 CT 시행 중 평균 심박수(bpm)는 1군이 2군에 비해 유의하게 낮았으며( $50.3 \pm 5.6$  vs.  $53.3 \pm 4.8$ ,  $p = 0.016$ ), 검사 중 심박변화는 두 군 간 차이가 없었다. 분절별 분석에서 1군의 우관상동맥 중분절( $3.0 \pm 0.9$  vs.  $2.5 \pm 1.1$ ,  $p = 0.039$ )과 좌전하행동맥 제1사선분절( $3.4 \pm 0.6$  vs.  $3.1 \pm 0.7$ ,  $p = 0.024$ )의 영상 질이 2군에 비해 유의하게 좋았다. 영상 질은 심박수와 음의 상관관계를 보였으나 심박변화와는 상관관계를 보이지 않았다. 혈관별 분석에서 영상 질은 두 군 간 차이가 없었다.

**결론:** 저용량 베타차단제는 느린 심박수 무증상 환자에서 심박수 감소효과가 있으며, 전향적 동조화 심장 CT 일부 분절의 영상 질을 높일 수 있으나, 그 효과는 제한적이다.

<sup>1</sup>연세대학교 의과대학 강남세브란스병원 영상의학과, <sup>2</sup>울산대학교 의과대학 서울아산병원 영상의학과,

<sup>3</sup>연세대학교 의과대학 세브란스병원 영상의학과