



Growth hormone treatment and risk of malignancy

Hyun-Wook Chae, MD¹, Duk-Hee Kim, MD, PhD², Ho-Seong Kim, MD, PhD¹

¹Department of Pediatrics, Endocrine Research Institute, Yonsei University College of Medicine, Seoul, ²Sowha Children's Hospital, Seoul, Korea

Growth hormone (GH) treatment has been increasingly widely used for children with GH deficiencies as the survival rate of pediatric patients with malignancies has increased. Both GH and insulin-like growth factor-I have mitogenic and antiapoptotic activity, prompting concern that GH treatment may be associated with tumor development. In this review, the authors examined the relationship between GH treatment and cancer risk in terms of *de novo* malignancy, recurrence, and secondary neoplasm. Although the results from numerous studies were not entirely consistent, this review of various clinical and epidemiological studies demonstrated that there is no clear evidence of a causal relationship between GH treatment and tumor development. Nonetheless, a small number of studies reported that childhood cancer survivors who receive GH treatment have a small increased risk of developing *de novo* cancer and secondary malignant neoplasm. Therefore, regular follow-ups and careful examination for development of cancer should be required in children who receive GH treatment. Continued surveillance for an extended period is essential for monitoring long-term safety.

Key words: Growth hormone, Neoplasms, Recurrence, Malignancy insulin-like growth factor-I

Introduction

Growth hormone (GH) has been administered to children with GH deficiency since the late 1950s, initially using extracts from human pituitary glands¹. Since the 1980s, GH has been produced by recombinant DNA technology and prescribed for a wide variety of disorders. As the use of synthetic human recombinant GH has increased over the past several decades, there has been a concurrent increase in awareness of the potential safety issues accompanying the treatment². Recently, Carel et al.³ analyzed a French population-based registry of 6,928 children who started GH treatment between 1985 and 1996. The researchers reported that mortality rates increased in children treated with recombinant GH; however, all types of cancer-related mortality did not increase, and most of the increased mortality was due to bone tumors or cerebral hemorrhage. This exemplifies the controversy associated with GH treatment and risk of malignancy.

Survival rates of pediatric patients with malignancies are increasing due to improved treatment modalities, and interest in improved, long-term quality of life is increasing in tandem, including optimizing the final height of a patient with GH deficiency after cancer treatment⁴. For patients with GH deficiency after completing antitumor therapy, GH is often recommended to increase height velocity and improve quality of life⁵.

On its own, however, GH has mitogenic activity⁶. There have been theoretical and clinical concerns that GH may play a part in tumor recurrence^{7,8}. GH treatment may increase an individual's risk of developing cancer, particularly in the case of cancer survivors, by increasing the risk of recurrence or secondary malignancy⁹. Controversies

Corresponding author: Ho-Seong Kim, MD, PhD
Department of Pediatrics, Severance Hospital,
Yonsei University College of Medicine, 50 Yonsei-ro,
Seodaemun-gu, Seoul 120-752, Korea
Tel: +82-2-2228-2069
Fax: +82-2-393-9118
E-mail: kimho@yuhs.ac

Received: 25 August, 2014

Accepted: 6 November, 2014

Copyright © 2015 by The Korean Pediatric Society

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

and debates are currently ongoing regarding GH treatment's possible causal relationship with tumor recurrence.

It is known from *in vitro* and *in vivo* studies that GH and insulin-like growth factor-I (IGF-I) have mitogenic and anti-apoptotic activity, and it is possible that GH treatment may be associated with tumor development^{7,8}. Higher serum levels of IGF-I may also be associated with increased cancer risk, and concern has been raised regarding its potential role as a carcinogen after GH treatment¹⁰. Some epidemiological studies have shown a correlation between high serum levels of IGF-I and occurrence of certain cancers, including carcinomas of the prostate, lung, breast, and colon^{11,12}. GH receptors have been expressed on normal and transformed human white blood cells, and *in vitro* studies have shown that GH may cause transformation of normal cells and proliferation of leukemic cells^{9,13}. Most of IGF-I's physical effects are mediated through the type 1 IGF receptor, which is overexpressed in many different types of cancers¹⁴. Although the results of all studies are not completely consistent, a high concentration of IGF-I may be linked to an increased risk of cancer development¹⁵.

As with IGF-I, there is a concern that GH treatment may be associated with tumor development. Although GH treatment has been shown to be generally safe, it is important to investigate the relationship between GH treatment and cancer risk. The authors reviewed clinical and epidemiological studies that examined cancer risk in patients treated with GH with a focus on *de novo* malignancy, recurrence, and secondary neoplasm.

GH treatment and *de novo* malignancy

1. Leukemia

A possible correlation between GH therapy and increased risk of leukemia was reported in Japan¹⁶; however, the investigators carried out a follow-up analysis of the cohort and could not find a close association between GH therapy and leukemia when subjects with known risk factors for leukemia were excluded¹⁷. The United States' National Hormone Pituitary Program found 3 cases of leukemia in 59,736 patient-years of GH treatment between 1963 and 1985. The rate was not significantly higher than the 1.6 cases expected for an age, ethnicity, and sex-matched population¹⁸. Genentech's National Cooperative Growth Study (NCGS), which enrolled more than 40,000 GH recipients during 20 years of follow-up, showed that the leukemia risk was comparable to that of the general population when subjects with known risk factors for leukemia were excluded¹⁹. Another study from the NCGS reported 3 cases of *de novo* leukemia compared with the 5.6 cases expected in a theoretical, age-matched general population²⁰.

2. Solid tumors

There has been concern that elevated endogenous levels of GH and IGF-I might be associated with an increased risk of certain solid tumors. Swerdlow et al.²¹ studied cancer incidence and mortality in 1,848 patients in the UK who were treated with human pituitary GH during childhood and early adulthood between 1959 and 1985. The incidence and mortality of colorectal cancer and the mortality of Hodgkin's disease initially increased after patients with an otherwise high risk of cancer were excluded. Since then, however, the validity of the results have been debated, because the current GH therapy regimen differs from what was used at the time⁸. All patients received standard GH doses given 2 or 3 times a week, and serum IGF-I levels were not monitored.

Tyden et al.²² reported 2 cases of *de novo* development of cancer in living kidney transplant recipients, although *de novo* development of cancer in renal transplants might be rare. According to the analysis from the Kabi International Growth Study (KIGS) and NCGS databases, however, only 2 cases of renal cell carcinoma in those who did not have renal disease were found among the 43,000 patients in the NCGS registry and 42,000 in the KIGS registry²³. The number of malignancies seemed disproportionately high for the relatively small number of children who had chronic renal failure and who were receiving GH treatment^{8,23}.

Tuffli et al.²⁴ reported that the number of extracranial, non-leukemic neoplasms from 12,209 individuals who were treated with the recombinant GH was no greater than expected cases. Although 10 new cases of malignancies were noted, it was not a greater value than predicted, indicating that GH was not implicated in the occurrence of solid tumors. Another report from NCGS showed no evidence of an increase in the incidence of *de novo* intracranial tumors in children treated with GH²⁵. Bell et al.²⁰ reported, from a recent analysis of the NCGS registry, that *de novo* intracranial and extracranial malignancies were not significantly increased in patients without risk factors (29 confirmed vs. 26 expected). One of the most recent reports from the KIGS compared the incidence of cancer in the cohort to that in the general population by using the standardized incidence ratio (SIR)²⁶. A total of 32 new malignant neoplasms were reported in 58,603 patients, versus the 25.3 expected (SIR, 1.26; 95% confidence interval [CI], 0.86–1.78). Again, patients treated with GH showed no statistically significant difference in cancer incidence compared with the expected number of cases in this study.

GH treatment and recurrence of malignancy

1. Leukemia

Survivors of childhood leukemia are at risk of developing complications, including growth failure, which may require GH treatment⁸⁾. Patients with GH deficiency are particularly common because total body irradiation increases risk of growth failure, although GH deficiency can also result from solely chemotherapeutic regimens²⁷⁾. There has been concern that children treated with GH after therapy against leukemia may be at a higher risk of leukemia recurrence.

Leung et al.²⁸⁾ studied 47 patients who had received GH replacement therapy among 910 patients treated for acute lymphoblastic leukemia and examined recurrence rates at 7 years and 11 years after continuous hematologic remission. There was no statistical evidence that GH therapy was associated with leukemia relapse or development of a second malignancy.

The NCGS has monitored the safety of recombinant human GH since 1985. There have been 3 new cases of leukemia in children without known risk factors for developing leukemia and 5 cases in children with known risk factors; however, there was no evidence of an increased recurrence of leukemia²⁹⁾. Follow-up reports showed no evidence of increased incidence of leukemia among patients without previous risk factors³⁰⁾. Another investigator analyzed 47,000 patients representing 165,000 patient-years from the NCGS and found reassuring evidence that leukemia (*de novo* or relapse), extracranial nonleukemic neoplasms, and central nervous system (CNS) tumor recurrence were not associated with GH therapy³¹⁾.

Sklar et al.³²⁾ investigated 122 acute leukemia survivors from among 13,539 subjects enrolled in the Childhood Cancer Survivor Study (CCSS), a cohort of 5-year survivors of childhood cancer. They found that the relative risk of recurrence was not increased compared to 4,545 children not treated with GH.

2. Brain tumors

GH deficiency is a common disease in patients with hypothalamic-pituitary tumors. The deficiency is caused either by the mass itself or by surgical or irradiation therapy to hypothalamic-pituitary axis⁹⁾. Deficiency can also develop in those who have tumors far from the hypothalamic-pituitary axis, because cerebrospinal irradiation is often needed to treat these tumors³³⁾.

In 1985, Arslanian et al.³⁴⁾ reported the outcome of GH therapy in 34 children with brain tumors who developed hypopituitarism. Of the 34 patients with brain tumors and hypopituitarism, 24 received GH therapy. Eight of the 24 (33%) experienced tumor recurrence, compared with 3 of the 10 (30%) who did not receive GH. Clayton et al.³⁵⁾ reported similar results, namely that the late relapse rate of medulloblastoma and glioma was not altered by

GH therapy, and the therapy might not increase the relapse rate of brain tumors.

Medulloblastoma is one of the most malignant childhood brain tumors. Survivors of medulloblastoma are, however, increasing as progress has been made in the treatment of the tumor³⁶⁾. From a retrospective analysis of 34 children treated with GH for medulloblastoma over 3 years, Chae et al.³⁷⁾ reported no cases of recurrence. Another retrospective review included 545 children with medulloblastoma, of whom 170 patients were treated with GH³⁸⁾. This review found that while GH treatment was underutilized in survivors of medulloblastoma, it was not associated with disease relapse.

Craniopharyngioma is a relatively common brain tumor derived from pituitary gland embryonic tissue in children³⁹⁾. Due to the morbidities associated with damage to the pituitary and hypothalamus from surgical removal and irradiation against craniopharyngioma, GH treatment is commonly needed after therapy to treat this tumor⁴⁰⁾. The recurrence rate of craniopharyngioma was 0.045/treatment year in 488 patients who were enrolled in the KIGS from 1988 to 1996, and the investigators concluded that GH treatment might be safe and effective in children with craniopharyngioma⁴¹⁾. In the NCGS report, children receiving GH after treatment for craniopharyngioma had a recurrence rate of 6.4%, considerably lower than the estimates of 20%–25% in another report^{25,42)}.

Swerdlow et al.⁴³⁾ investigated 180 children with brain tumors treated with GH, and 891 children not treated with GH during 1965–1996. The relative risk of first recurrence in GH-treated patients decreased after adjusting for potentially confounding prognostic variables (0.6; 95% CI, 0.4–0.9). The relative risk of mortality also decreased (0.5; 95% CI, 0.3–0.8). There was no significant trend in relative risk of recurrence with cumulative time for which GH treatment had been given or with time elapsed since GH treatment started. Another study showed, using surveillance imaging, that hypothalamic-pituitary tumor recurrence in GH-treated patients was as low as in non-GH-treated patients⁴⁴⁾. Only 1 patient among 100 consecutive patients showed evidence of slight intrasellar tissue enlargement using pituitary imaging at 6 months. GH replacement was continued, and there was no further change between 6 and 12 months, though the follow-up duration was short and may not be conclusive evidence against recurrence.

GH treatment and risk of secondary malignancy

Childhood cancer survivors might be at increased risk for secondary malignancies compared with the general population⁴⁵⁾. Assessing their risk of second and subsequent malignancies during long term follow-up is thus very important.

Meadows et al.⁴⁶⁾ investigated 14,358 cohort members in the CCSS and analyzed SIRs for occurrences of a second malignant neoplasm. The 30-year cumulative incidence of second malignant neoplasm was 9.3%, and the risk of subsequent neoplasms remained elevated for more than 20 years of follow-up for all diagnoses of primary childhood cancer. Another report from the CCSS showed that the risk of sarcoma was more than 9 folds greater among childhood cancer survivors than among the general population⁴⁷⁾. Ergun-Longmire et al.⁴⁸⁾ analyzed the cohort and reported that the rate of GH-treated survivors developing secondary neoplasms was 2.1 folds greater compared with non-GH-treated survivors; however, the risk appeared to decrease as follow-up time extended. Using large cohort follow-up studies, Carel and Butler⁴⁹⁾ suggested that children treated with GH following childhood cancer treatment might not have a greater number of relapses, but that there might be a higher incidence of secondary tumors in the early years of GH therapy.

There are, however, some more reassuring data. Neglia et al.⁵⁰⁾ analyzed 13,581 children from a retrospective cohort drawn from US and Canadian institutions who were diagnosed with common cancers before the age of 21 and survived at least 5 years. Twenty years after the childhood cancer diagnosis, the estimated cumulative incidence of second malignancy was 3.2%, and only 1.88 excess malignancies occurred during follow-up. Sklar et al.³²⁾ studied 172 brain tumor survivors from among 13,539 survivors enrolled in the CCSS. The relative risk of disease recurrence was 0.83 (95% CI, 0.37–1.86) for survivors who had been treated with GH, and risk was not increased for any of the major cancer diagnoses.

Most recently, Patterson et al.⁵¹⁾ analyzed 12,098 pediatric cancer survivors from the CCSS and reported the incidence of meningioma, glioma, and other solid CNS tumors. The adjusted rate of CNS tumor development in patients treated with GH compared with untreated survivors was 1.0 (95% CI, 0.6–1.8). Overall, there was no statistically significant increase in risk of the occurrence of solid CNS tumors associated with GH exposure. Major recent studies of growth hormone treatment and

malignancy risk from large data registries were summarized in Table 1.

Levels of circulating IGF-I might be associated with an increased risk of common cancers, although the association remains unclear⁵²⁾, as there have been few large cohort studies studying the relationship between GH dosage, IGF-I levels, and risk of tumor recurrence. A meta-analysis and systematic review showed that high concentrations of IGF-I were associated with an increased risk of prostate cancer and premenopausal breast cancer, but the associations were modest⁵²⁾. Nonetheless, monitoring and maintaining IGF-I levels within a normal range might be important, and titrating GH doses to target IGF-I levels (IGF-I based dosing) could be a reasonable approach in this high risk GH-treated group⁴⁹⁾.

Conclusions

The benefits of GH treatment for growth and metabolism in children with GH deficiency are well defined, and replacement therapy with GH is recommended⁵³⁾. Despite theoretical concerns about the effect of GH on tumor development, this review of various clinical and epidemiological studies demonstrated that there is no clear evidence of a causal relationship between GH treatment in patients with GH deficiency and tumor development. Nonetheless, a small number of studies have reported that childhood cancer survivors who have received GH treatment have a small increased risk of *de novo* cancer and second malignant neoplasm. Therefore, regular follow-ups and careful examination for the development of cancer should be required in those who receive GH treatment. Continued surveillance for an extended period is essential to monitor further potential changes.

Conflict of interest

No potential conflict of interest relevant to this article was

Table 1. Major recent studies of growth hormone treatment and malignancy risk (large data registries)

Source	Year of publication	Data registry	No. of patients	No. of observed cases	No. of expected cases	SIR (95% CI)	Type of observed malignancy
Patterson et al. ⁵¹⁾	2014	CCSS	14,358	16	-	1.0 (0.6–1.8)	Secondary CNS malignancy
Wilton et al. ²⁶⁾	2010	KIGS	58,603	32	25.3	1.26 (0.86–1.78)	<i>De novo</i> malignancy
Bell et al. ²⁰⁾	2010	NCGS	54,996	29	26.0	1.12 (0.75–1.61)	<i>De novo</i> nonleukemic malignancy
				3	5.6	0.54 (0.11–1.58)	<i>De novo</i> leukemia
Wyatt ³¹⁾	2004	NCGS	47,000	16	15.3	-	Recurred nonleukemic malignancy
Maneatis et al. ³⁰⁾	2000	NCGS	33,161	20	-	0.73 (0.20–1.86)	Recurred leukemia
				35	-	0.44 (0.24–0.74)	Recurred nonleukemic malignancy

SIR, standardized incidence ratio; CI, confidence interval; CCSS, the Childhood Cancer Survivor Study; CNS, central nervous system; KIGS, the Kabi International Growth Study; NCGS, the National Cooperative Growth Study.

reported.

References

- Blizzard RM. History of growth hormone therapy. *Indian J Pediatr* 2012;79:87-91.
- Laron Z. Growth hormone therapy: emerging dilemmas. *Pediatr Endocrinol Rev* 2011;8:364-73.
- Carel JC, Ecosse E, Landier F, Meguellati-Hakkas D, Kaguelidou F, Rey G, et al. Long-term mortality after recombinant growth hormone treatment for isolated growth hormone deficiency or childhood short stature: preliminary report of the French SAGhE study. *J Clin Endocrinol Metab* 2012;97:416-25.
- Noorda EM, Somers R, van Leeuwen FE, Vulmsa T, Behrendt H; Dutch Late Effects Study Group. Adult height and age at menarche in childhood cancer survivors. *Eur J Cancer* 2001;37:605-12.
- Adan L, Sainte-Rose C, Souberbielle JC, Zucker JM, Kalifa C, Brauner R. Adult height after growth hormone (GH) treatment for GH deficiency due to cranial irradiation. *Med Pediatr Oncol* 2000;34:14-9.
- Wabitsch M, Braun S, Hauner H, Heinze E, Ilondo MM, Shymko R, et al. Mitogenic and antiadipogenic properties of human growth hormone in differentiating human adipocyte precursor cells in primary culture. *Pediatr Res* 1996;40:450-6.
- Ogilvy-Stuart AL, Gleeson H. Cancer risk following growth hormone use in childhood: implications for current practice. *Drug Saf* 2004;27:369-82.
- Banerjee I, Clayton PE. Growth hormone treatment and cancer risk. *Endocrinol Metab Clin North Am* 2007;36:247-63.
- Sklar CA. Growth hormone treatment: cancer risk. *Horm Res* 2004;62 Suppl 3:30-4.
- Pekic S, Popovic V. GH therapy and cancer risk in hypopituitarism: what we know from human studies. *Eur J Endocrinol* 2013;169:R89-97.
- Khandwala HM, McCutcheon IE, Flyvbjerg A, Friend KE. The effects of insulin-like growth factors on tumorigenesis and neoplastic growth. *Endocr Rev* 2000;21:215-44.
- Furstenberger G, Senn HJ. Insulin-like growth factors and cancer. *Lancet Oncol* 2002;3:298-302.
- Mercola KE, Cline MJ, Golde DW. Growth hormone stimulation of normal and leukemic human T-lymphocyte proliferation in vitro. *Blood* 1981;58:337-40.
- Baserga R, Peruzzi F, Reiss K. The IGF-1 receptor in cancer biology. *Int J Cancer* 2003;107:873-7.
- Cohen P, Clemmons DR, Rosenfeld RG. Does the GH-IGF axis play a role in cancer pathogenesis? *Growth Horm IGF Res* 2000;10:297-305.
- Watanabe S, Tsunematsu Y, Fujimoto J, Komiyama A. Leukemia in patients treated with growth-hormone. *Lancet* 1988;1:1159-60.
- Nishi Y, Tanaka T, Takano K, Fujieda K, Igarashi Y, Hanew K, et al. Recent status in the occurrence of leukemia in growth hormone-treated patients in Japan. GH Treatment Study Committee of the Foundation for Growth Science, Japan. *J Clin Endocrinol Metab* 1999;84:1961-5.
- Fradkin JE, Mills JL, Schonberger LB, Wysowski DK, Thomson R, Durako SJ, et al. Risk of leukemia after treatment with pituitary growth hormone. *JAMA* 1993;270:2829-32.
- Allen DB, Rundle AC, Graves DA, Blethen SL. Risk of leukemia in children treated with human growth hormone: review and reanalysis. *J Pediatr* 1997;131(1 Pt 2):S32-6.
- Bell J, Parker KL, Swinford RD, Hoffman AR, Maneatis T, Lippe B. Long-term safety of recombinant human growth hormone in children. *J Clin Endocrinol Metab* 2010;95:167-77.
- Swerdlow AJ, Higgins CD, Adlard P, Preece MA. Risk of cancer in patients treated with human pituitary growth hormone in the UK, 1959-85: a cohort study. *Lancet* 2002;360:273-7.
- Tyden G, Wernersson A, Sandberg J, Berg U. Development of renal cell carcinoma in living donor kidney grafts. *Transplantation* 2000;70:1650-6.
- Mehls O, Wilton P, Lilien M, Berg U, Broyer M, Rizzoni G, et al. Does growth hormone treatment affect the risk of post-transplant renal cancer? *Pediatr Nephrol* 2002;17:984-9.
- Tuffli GA, Johanson A, Rundle AC, Allen DB. Lack of increased risk for extracranial, nonleukemic neoplasms in recipients of recombinant deoxyribonucleic acid growth hormone. *J Clin Endocrinol Metab* 1995;80:1416-22.
- Moshang T Jr, Rundle AC, Graves DA, Nickas J, Johanson A, Meadows A. Brain tumor recurrence in children treated with growth hormone: the National Cooperative Growth Study experience. *J Pediatr* 1996;128(5 Pt 2):S4-7.
- Wilton P, Mattsson AF, Darendeliler F. Growth hormone treatment in children is not associated with an increase in the incidence of cancer: experience from KIGS (Pfizer International Growth Database). *J Pediatr* 2010;157:265-70.
- Haddy TB, Mosher RB, Nunez SB, Reaman GH. Growth hormone deficiency after chemotherapy for acute lymphoblastic leukemia in children who have not received cranial radiation. *Pediatr Blood Cancer* 2006;46:258-61.
- Leung W, Rose SR, Zhou Y, Hancock ML, Burstein S, Schriock EA, et al. Outcomes of growth hormone replacement therapy in survivors of childhood acute lymphoblastic leukemia. *J Clin Oncol* 2002;20:2959-64.
- Blethen SL, Allen DB, Graves D, August G, Moshang T, Rosenfeld R. Safety of recombinant deoxyribonucleic acid-derived growth hormone: The National Cooperative Growth Study experience. *J Clin Endocrinol Metab* 1996;81:1704-10.
- Maneatis T, Baptista J, Connelly K, Blethen S. Growth hormone safety update from the National Cooperative Growth Study. *J Pediatr Endocrinol Metab* 2000;13 Suppl 2:1035-44.
- Wyatt D. Lessons from the national cooperative growth study. *Eur J Endocrinol* 2004;151 Suppl 1:S55-9.
- Sklar CA, Mertens AC, Mitby P, Occhiogrosso G, Qin J, Heller G, et al. Risk of disease recurrence and second neoplasms in survivors of childhood cancer treated with growth hormone: a report from the Childhood Cancer Survivor Study. *J Clin Endocrinol Metab* 2002;87:3136-41.
- Bernier V. Technical aspects in cerebrospinal irradiation. *Pediatr Blood Cancer* 2004;42:447-51.
- Arslanian SA, Becker DJ, Lee PA, Drash AL, Foley TP Jr. Growth hormone therapy and tumor recurrence. Findings in children with brain neoplasms and hypopituitarism. *Am J Dis Child* 1985;139:347-50.
- Clayton PE, Shalet SM, Gattamaneni HR, Price DA. Does growth hormone cause relapse of brain tumours? *Lancet* 1987;1:711-3.
- Ranke MB, Price DA, Lindberg A, Wilton P, Darendeliler F, Reiter EO. Final height in children with medulloblastoma treated with growth hormone. *Horm Res* 2005;64:28-34.
- Chae HW, Park YS, Kim DS, Kwon AR, Kim HS, Kim DH. Final height and insulin-like growth factor-1 in children with medulloblastoma treated with growth hormone. *Childs Nerv Syst* 2013;29:1859-63.
- Packer RJ, Boyett JM, Janss AJ, Stavrou T, Kun L, Wisoff J, et

- al. Growth hormone replacement therapy in children with medulloblastoma: use and effect on tumor control. *J Clin Oncol* 2001; 19:480-7.
39. Price DA, Jonsson P. Effect of growth hormone treatment in children with craniopharyngioma with reference to the KIGS (Kabi International Growth Study) database. *Acta Paediatr Suppl* 1996;417:83-5.
 40. Hogeveen M, Noordam C, Otten B, Wit JM, Massa G. Growth before and during growth hormone treatment in children operated for craniopharyngioma. *Horm Res* 1997;48:258-62.
 41. Price DA, Wilton P, Jonsson P, Albertsson-Wikland K, Chatelain P, Cutfield W, et al. Efficacy and safety of growth hormone treatment in children with prior craniopharyngioma: an analysis of the Pharmacia and Upjohn International Growth Database (KIGS) from 1988 to 1996. *Horm Res* 1998;49:91-7.
 42. Weiss M, Sutton L, Marcial V, Fowble B, Packer R, Zimmerman R, et al. The role of radiation therapy in the management of childhood craniopharyngioma. *Int J Radiat Oncol Biol Phys* 1989;17: 1313-21.
 43. Swerdlow AJ, Reddingius RE, Higgins CD, Spoudeas HA, Phipps K, Qiao Z, et al. Growth hormone treatment of children with brain tumors and risk of tumor recurrence. *J Clin Endocrinol Metab* 2000;85:4444-9.
 44. Frajese G, Drake WM, Loureiro RA, Evanson J, Coyte D, Wood DF, et al. Hypothalamo-pituitary surveillance imaging in hypopituitary patients receiving long-term GH replacement therapy. *J Clin Endocrinol Metab* 2001;86:5172-5.
 45. Henderson TO, Rajaraman P, Stovall M, Constine LS, Olive A, Smith SA, et al. Risk factors associated with secondary sarcomas in childhood cancer survivors: a report from the childhood cancer survivor study. *Int J Radiat Oncol Biol Phys* 2012;84:224-30.
 46. Meadows AT, Friedman DL, Neglia JP, Mertens AC, Donaldson SS, Stovall M, et al. Second neoplasms in survivors of childhood cancer: findings from the Childhood Cancer Survivor Study cohort. *J Clin Oncol* 2009;27:2356-62.
 47. Henderson TO, Whitton J, Stovall M, Mertens AC, Mitby P, Friedman D, et al. Secondary sarcomas in childhood cancer survivors: a report from the Childhood Cancer Survivor Study. *J Natl Cancer Inst* 2007;99:300-8.
 48. Ergun-Longmire B, Mertens AC, Mitby P, Qin J, Heller G, Shi W, et al. Growth hormone treatment and risk of second neoplasms in the childhood cancer survivor. *J Clin Endocrinol Metab* 2006;91:3494-8.
 49. Carel JC, Butler G. Safety of recombinant human growth hormone. *Endocr Dev* 2010;18:40-54.
 50. Neglia JP, Friedman DL, Yasui Y, Mertens AC, Hammond S, Stovall M, et al. Second malignant neoplasms in five-year survivors of childhood cancer: childhood cancer survivor study. *J Natl Cancer Inst* 2001;93:618-29.
 51. Patterson BC, Chen Y, Sklar CA, Neglia J, Yasui Y, Mertens A, et al. Growth hormone exposure as a risk factor for the development of subsequent neoplasms of the central nervous system: a report from the childhood cancer survivor study. *J Clin Endocrinol Metab* 2014;99:2030-7.
 52. Renehan AG, Zwahlen M, Minder C, O'Dwyer ST, Shalet SM, Egger M. Insulin-like growth factor (IGF)-I, IGF binding protein-3, and cancer risk: systematic review and meta-regression analysis. *Lancet* 2004;363:1346-53.
 53. Kelnar CJ. Which children should receive growth hormone treatment. Cost-benefit analysis is the key. *Arch Dis Child* 2000;83:176-8.