

Uterine Arterial Embolization for the Treatment of Leiomyomas Accompanying with Adenomyosis

Jin Beum Jang, Sang Wook Bai, Jae Hak Lim, Do Yeon Lee^{*}, Jung Yeon Kim,
Kyung Ah Jeong, Sei Kwang Kim, Ki Hyun Park

Department of Obstetrics and Gynecology, Department of Radiology^{},
Yonsei university, College of Medicine, Seoul, Korea*

Objective: The purpose of this study was to make a guideline of uterine artery embolization for the treatment of uterine leiomyomas accompanying with adenomyosis in Korea.

Materials and Methods: We performed the retrospective study for 37 women who had uterine leiomyomas accompanying with adenomyosis. Bilateral uterine artery embolization was performed in 37 patients (age range 25-65) during 17 months with pain, hypermenorrhea, urinary frequency etc due to leiomyomas. Ultrasound imaging was performed before the procedure and at mean 6.9 months after the procedure.

Results: All procedures were technically successful. Mean clinical follow-up was 12.8 months. Minor complication occurred in 82% patients after the procedure. After imaging follow-up (mean, 6.9 months postprocedure), median uterine volume decreased 34.4%, and dominant myoma volume decreased 86%. There was no statistical difference in uterine volume reduction and dominant myoma size reduction whether occluding agents was polyvinyl alcohol, polyvinyl alcohol plus gelfoam, and gelfoam, and whether ultrasound measured Resistance Index value before the procedure was low or high.

Conclusion: Primary candidates for uterine artery embolization include those with symptomatic uterine leiomyomas who no longer desire fertility but wish to avoid surgery or are poor surgical risks. To our study, uterine volume reduction and dominant myoma size reduction in patients who had adenomyosis were similar to previous other studies in patients who had not adenomyosis. Therefore adenomyosis should not be considered as a contraindication for uterine artery embolization. Because there is little data about subsequent reproductive potential after this procedure, it should not be routinely advocated for infertile women. Further investigation is warranted for occluding agents and Resistance Index.

Key Words: Uterine artery embolization, Leiomyoma, Adenomyosis, Occluding agent, Resistance index

가
 , .40 가 가
 40% 가 가 , 2000 2,000 6,000
 , 39 . ,
 , 가가 , ,
 . , .1 .5 ,
 , 600,000 ,
 , 73,000 가 가 ,
 , 25 30% 가 가 , 1998 1
 . 7 10 2000 10
 , 2 3 37
 , 가 가
 , ! ,
 , Resistance index (RI) 가

(Gonadotropin regulating hormone agonist)

1.
 2000 WM Liu가 1998 5 2000 10
 ,² Nd-YAG
 , Pirfenidone 37
 .³
 7 Lariboisiere JH Ravina가
 1988 가 40 , Doppler
 Resistance index
 Resistance index (RI)
 1995 Lancet 4 , ,
 , , alcohol , polyvinyl alcohol gelfoam

Table 1. Characteristics of Patients Undergoing Uterine Artery Embolization

Total Number	37
Age	40.0 years (25~65)
Initial Hemoglobin	9.9 mg/dL (4.8~14)
Initial Hematocrit	30.9% (15.7~41.7)
Uterine Volume Reduction	34.4%
Dorminant Myoma Size Reduction	86%
Duration of Follow-up Sonogram	6.9 months (1~28)
Duration of Total Follow-up	12.8 months (1~31)

Table 2. Current Symptoms of Patients

	Number	Percent
Bleeding	21	57%
Dysmenorrhea	9	24%
Abdominal Pain	8	22%
Back Pain	3	8%
Urinary Symptoms	4	11%
No Symptoms	4	11%

Table 3. Indications of Uterine Artery Embolization

	Number	Percent
High Operation Risks	10	27%
Adjuvant to Myomectomy	5	14%
Previous Myomectomy	3	8%
Previous GnRH analogue treatment	3	8%

gelfoam

2.

29 , 3

5 .4

French 5 French Glide cobra catheter (Terumo, Tokyo, Japan)

(Tracker 325, Boston Scientific, Watertown, Miami, USA)

6 polyvinyl alcohol (contour, Boston Scientific, Watertown, Miami, USA) 350~500 ì m 500~700 ì m 19 , gelfoam (spongostan, Johnson & Johnson, Gargrave, UK)

6 , 7 가

가

1

30 .

3.

one-way

ANOVA , p value <0.05

40 (25~65)

6

1

(Table 1).

57% (n=21) ,

54% (n=20),

11% (n=4) ,

가 11% (n=4) (Table 2).

(Gonadotropin regulating hormone agonist)

8% (n=3)

8%

가 27%

(n=10) ,

14% (n=5) (Table 3).

11 , 73% (n=29)

1.59 ± 0.8

59% (n=22), 3%

(n=1), 24% (n=9),

가 11% (n=4) . 10% (n=3)

가 가 , 90% (n=26)

. 14% (n=3)

가 가 , 86% (n=19) 가

Table 4. Correlations between Uterine Volume Reduction (UVR), Dominant Myoma Size Reduction (MSR) and Resistance index (RI) changes of Myoma Feeding Vessels (MFV), Right Uterine Artery (RUI), Left Uterine Artery (LUI)

	RI changes of MFV	RI changes of RUI	RI changes of LUI
UVR	0.389 (p=0.300)	0.344 (p=0.210)	-0.469 (p=0.078)
MSR	-0.253 (p=0.681)	0.362 (p=0.304)	-0.066 (p=0.857)

Table 5. Uterine Volume Reduction (UVR) and Dominant Myoma Size Reduction (MSR) After Using Different Embolizing Material (Polyvinyl alcohol, Polyvinyl alcohol and gelfoam, gelfoam)

	Number	Mean (%)	p value
UVR			
PVA	13	32.2	0.856
PVA+gelfoam	7	27.0	
Gelfoam	5	34.0	
MSR			
PVA	9	58.9	0.296
PVA+gelfoam	6	33.5	
Gelfoam	4	40.0	

polyvinyl alcohol, polyvinyl alcohol gelfoam, gelfoam

polyvinyl alcohol gelfoam, 가 (Table 5), 79%, 가 25%, 14%

Heaston DK 1979, gelatin sponge, 34.4%, 86%, (1~28), 12.8 (1~31), Resistance index (RI) (Pearson correlation coefficient = 0.389) (p value = 0.300), 18, 가, Yamashita Y 1994, 17, 7, 10, McIvor J 1996, 7, 10, Volgelzang RL 1991, 4, 11, 8, 가, (Pearson correlation coefficient = -0.469, -0.066, Table 4).

JH Ravina 1995 34 49 16 20 5
(11 48) 75% 가 ,
11
, 10 , 2000 1,730
2 32 7
, ⁴ Goodwin SC 1997 11
5.8 (2 가 ,
9) 가 40%
, 가 가 60%
65% 가 .¹² JH Ravina 가 2000 JH
1997 88 Ravina 184
6 60 9 12 7
9 , 6 , 4
69% ,¹³ Bradley 3 ,
EA 1998 31 48 8 20 2 37 41
가 3 1179.0 cm³ 583.3 가 .¹⁶ ,
cm³ .¹⁴ 1% 2%
6.9 14% (n=3) , 2000 Spies
가 가 66 4
, 34% , (Follicle stimulating hormone) 3.5 mIU/ml
86% 가 .¹⁷
1998 Worthington-Kirsch RL 1997 Goo-
dwin SC 3 gelfoam 1997
가 85% 90% 1 6
,¹⁵ 1998 Bradley EA 7 , McLvor J
6 .¹⁴ 1996 7 polyvinyl alcohol,
coil, gelfoam 3
가 .¹⁰ ,
. 7 cm , ,
, 가 가
가
, 8 cm , Bradley EA,
가 , JH Ravina, Goodwin SC

가 13
 1 1
 가 , 가 .²² , Goodwin Sc 1999
 가 . 가 3 ,¹⁸ 2000 Siskin GP
 Bradley JH . 1998 가 14 , 11 10
 25 , 6 47%
 ,¹⁴ 1999 Goodwin Sc 59 .²³
 가 ,
 1390 cm³ (370 cm³~2880 cm³)
 510 cm³ .¹⁸
 , 1998 Goodwin SC
 Djabbari M 1999
 ,¹⁹ 1999 Mclucas 300
 15% ,²⁴
 8.7 cm 가 ,
²⁰ 가 가 가
 가 ,
 가 가
 Resistance
 90% index (RI) 가 ,
 86% 가
 가
 Resistance index (RI)
 34.4% 가
 40%~69% .
 20% 가
 가
 polyvinyl alcohol gel-
 foam 가 polyvinyl alcohol
 gelfoam
 1999 Smith SJ
 ,²¹ 1999 Goodwin
 SC 5 , polyvinyl alcohol gelfoam
 3 가
 ,¹⁸ 1999 Stancato-Pasik 가
 가

가 가 .5
, 2000
Sidney Royal North Shore RW Dover
가 가
, 3 1
가 ,
, ,
Mallory-Weiss .0
Abulapia 1999 가 , 가
, guide wire , ,
, ,
.25 가
, 1999
Chelsea & Westminster Vashisht ,
가 51 20 ,
7 ,
21 ,
.26 ,
, , , , , ,
1999 Georgetown JB Spies ,
, , ,
가 ,
, , ,
.27 Roxborough 가 가
FL Hutchins Jr. 2000 가
, , , ,
, , ,
.28 North Carolina BS Hurst ,
2000 , ,
(hydrosalpinx), 가 ,
, , , , ,

1. Reidy JF, Bradley EA. Uterine artery embolization for fibroid disease. Cardiovasc Intervent Radiol 1998; 21: 357-60.

2. Liu WM. Laparoscopic bipolar coagulation of uterine vessels to treat symptomatic leiomyomas. J Am Assoc Gynecol Laparosc 2000; 7(1): 125-9.

3. Lee BS, Margolin SB, Nowak RA. A novel pharmacological agent that inhibits leiomyoma cell proliferation. American society of reproductive medicine; 1996.

4. Ravina JH, Herbreteau D, Ciraru-Vigneron N, Bouret JM, Haudart E, Aymard A, et al. Arterial embolisation to treat uterine myomata. *Lancet* 1995; 345: 671-2.
5. Hurst BS, Stackhouse DJ, Matthews ML, Marshburn PB. Uterine artery embolization for symptomatic uterine myomas. *Fertil Steril* 2000; 74(5): 855-69.
6. , , , , , , , . 1999; 41: 481-5.
7. Heaston DK, Mineau DE, Brown BJ, Miller FJ. Transcatheter arterial embolization for control of persistent massive puerperal hemorrhage after bilateral surgical hypogastric artery ligation. *Am J Roentgenol* 1979; 133: 152-4.
8. Oliver JA, Lance JS. Selective embolization to control massive hemorrhage following pelvic surgery. *Am J Obstet Gynecol* 1979; 135: 431-2.
9. Yamashita Y, Harada M, Yamamoto H, Miyazaki T, Takahashi M, Miyazaki K, et al. Transcatheter arterial embolization of obstetric and gynaecological bleeding: Efficacy and clinical outcome. *Br J Radiol* 1994; 67: 530-4.
10. McIvor J, Cameron EW. Pregnancy after uterine artery embolisation to control haemorrhage from gestational trophoblastic tumor. *Br J Radiol* 1996; 69: 624-9.
11. Volgezang RL, Nemcek AA Jr., Skrtic Z, Gorrell J, Lurain JR. Uterine arteriovenous malformations: Primary treatment with therapeutic embolization. *J Vasc Interv Radiol* 1991; 2: 517-22.
12. Goodwin SC, Vedantham S, McLucas B, Forno AE, Perrella R. Preliminary experience with uterine artery embolization for uterine fibroids. *J Vasc Interv Radiol* 1997; 8: 517-26.
13. Ravina JH, Bouret JM, Ciraru-Vigneron N, Repiquet D, Herbreteau D, Aymard A, et al. Application of particulate arterial embolization in the treatment of uterine fibromyomata. *Bull Acad National Med* 1997; 181: 233-43.
14. Bradley EA, Reidy JF, Forman RG, Jarosz J, Braude PR. Transcatheter uterine artery embolisation to treat large uterine fibroids. *Br J Obstet Gynaecol* 1998; 105: 235-40.
15. Worthington-Kirsch RL, Popky GL, Hutchins FL Jr. Uterine arterial embolization for the management of leiomyoma: Quality-of-life assessment and clinical response. *Radiology* 1998; 208: 625-9.
16. Ravina JH, Vigneron NC, Aymard A, LeDref O, Merland JJ. Pregnancy after embolization of uterine myoma. *Fertil Steril* 2000; 73: 1241-3.
17. Spies JB, Walsh SM, Roth AM. Assessment of ovarian function via serum follicle stimulating hormone (FSH) assay before and after uterine fibroid embolization (UFE): Preliminary results [abstract]. Society of Cardiovascular and Interventional Radiology 25th Annual Scientific Meeting. San Diego, March 2000.
18. Goodwin SC, McLucas B, Lee M, Chen G, Perrella R, Vedantham S, et al. Uterine artery embolization for the treatment of uterine leiomyomata: midterm results. *J Vasc Interv Radiol* 1999; 10: 1159-65.
19. Goodwin SC, Walker WJ. Uterine artery embolization for the treatment of uterine fibroids. *Current Opinion Obstet Gynecol* 1998; 10: 315-20.
20. McLucas B, Adler L, Perrella R. Predictive factors for success in uterine fibroid embolization [abstract]. SMIT/CIMIT 11th Annual Scientific Meeting. Boston, Sept. 1999.
21. Smith ST, Sewell LE, Handelsman A. A clinical failure of uterine fibroid embolization due to adenomyosis. *J Vasc Interv Radiol* 1999; 10: 1171-4.
22. Stancato-Pasik A, Katz R, Mitty HA. Uterine artery embolization of myomas: Preliminary results of gelatin sponge pledgets as the embolitic agent [abstract]. SMIT/CIMIT 11th Annual Scientific Meeting. Boston, Sept. 1999.
23. Siskin GP, Tublin ME, Stainken BF, Dowling K, Ahn J, Dolen EG. Bilateral uterine artery embolization for the treatment of menorrhagia due to adenomyosis. Society of Cardiovascular and Interventional Radiology 25th Annual Scientific Meeting. San

- Diego, March 2000.
24. Djabbari M, Denys AL, Anquetil C, Levardon M, Menu YM. Preoperative bilateral uterine arterial embolization before multiple myomectomies: Is it unusual to reduce preoperative bleeding [abstract]. Radiological Society of North American Scientific Assembly and Annual meeting. Chicago, Nov. 1999.
 25. Abulafia O, Sherer DM. Transcatheter uterine artery embolization for the management of symptomatic uterine leiomyomas. *Obstet Gynecol Surv* 1999; 54 (12): 745-53.
 26. Vashisht A, Studd J, Carey A, Burn P. Fetal septicaemia after fibroid embolisation. *Lancet* 1999; 354: 307-8.
 27. Spies JB, Scialli AR, Jha RC, Imaoka I, Ascher SM, Fraga VM, et al. Initial results from uterine fibroid embolization for symptomatic leiomyomata. *J Vasc Interv Radiol* 1999; 10: 1149-57.
 28. Hutchins FL, Worthington-Kirsch R. Embolotherapy for myoma-induced menorrhagia. *Obstet Gynecol Clin North Am* 2000; 27(2): 397-405.
 29. Dover RW, Torode HW, Briggs GM. Uterine artery embolisation for symptomatic fibroid. *Med J Australia* 2000; 172: 233-6.
-