

Original Article
Psychiatry & Psychology



Association between Parents' Country of Birth and Adolescent Depressive Symptoms: the Early Stages of Multicultural Society

Jieun Jang ^{1,2} Eun-Cheol Park ^{2,3} Sang Ah Lee ^{1,2} Young Choi ^{1,2}
Yoon Soo Choy ^{1,2} Woorim Kim ^{1,2} and Sung-In Jang ^{2,3}

¹Department of Public Health, Graduate School, Yonsei University, Seoul, Korea

²Institute of Health Services Research, Yonsei University, Seoul, Korea

³Department of Preventive Medicine, Yonsei University College of Medicine, Seoul, Korea



Received: Jul 24, 2017

Accepted: Feb 26, 2018

Address for Correspondence:

Sung-In Jang, MD, PhD

Department of Preventive Medicine &
Institute of Health Services Research, Yonsei
University College of Medicine, 50 Yonsei-ro,
Seodaemun-gu, Seoul 03722, Korea.
E-mail: JANGSI@yuhs.ac

© 2018 The Korean Academy of Medical
Sciences.

This is an Open Access article distributed
under the terms of the Creative Commons
Attribution Non-Commercial License ([https://
creativecommons.org/licenses/by-nc/4.0/](https://creativecommons.org/licenses/by-nc/4.0/))
which permits unrestricted non-commercial
use, distribution, and reproduction in any
medium, provided the original work is properly
cited.


ORCID iDs

Jieun Jang 


<https://orcid.org/0000-0003-1797-8649>

Eun-Cheol Park 


<https://orcid.org/0000-0002-2306-5398>

Sang Ah Lee 


<https://orcid.org/0000-0001-5415-0141>

Young Choi 


<https://orcid.org/0000-0002-8314-6130>

Yoon Soo Choy 

<https://orcid.org/0000-0002-0301-8703>

Woorim Kim 

<https://orcid.org/0000-0002-1199-6822>

Sung-In Jang 

<https://orcid.org/0000-0002-0760-2878>

ABSTRACT

Background: This study aims to determine whether significant associations exist between the parents' country of birth and adolescent depressive symptoms in the early stages of a multicultural society.

Methods: We used data from the 2012–2016 Korea Youth Risk Behavior Web-based Survey, which included responses from 327,357 individuals. Participants were classified into groups according to their parent's country of birth. Logistic regression analysis was used to examine the significance of the associations.

Results: Adolescents whose parents were born abroad are more likely to have depressive symptoms (odds ratio [OR] = 1.68; 95% confidence interval [CI], 1.33–2.12) than adolescents whose parents were native Koreans. Respondents whose father was born in North Korea or Japan or Taiwan show greater odds of depressive symptoms than respondents whose parents were native Korean.

Conclusion: Adolescents whose parents were born abroad are more likely to have depressive symptoms. Multicultural family support policies should be implemented in consideration of the characteristics of the parents' country of birth.

Keywords: Multicultural; Depression; Adolescent; Parents' Birth of Country; Migration

INTRODUCTION

The pace of globalization has rapidly increased with the progress in information and communication technologies.¹ With globalization, people from different countries have become multilingual and multicultural,² resulting in the increased movement of populations internationally.³ According to the United Nations Population Division statistics, the number of immigrants in Korea has grown rapidly from 43,107 in 1990 to 1,327,324 in 2015.⁴ Considering that the current population of Korea is about 51 million, migrants account for a large proportion.⁵ Studies claim that globalization may have also led to an increase in the number of international marriages.⁶ The rate of marriage between Koreans and foreigners has nearly doubled, from 3.5% in 2000 to 7.3% in 2016.⁷

Disclosure

The authors have no potential conflicts of interest to disclose.

Author Contributions

Conceptualization: Jang J, Jang SI.
Investigation: Jang J, Park EC, Lee SA, Choi Y, Choy YS, Kim W. Supervision: Jang SI.
Validation: Jang J, Park EC, Lee SA, Choi Y, Choy YS, Kim W, Jang SI. Writing - original draft: Jang J. Writing - review & editing: Jang J, Park EC, Lee SA.

In 2015, the birth rate of multicultural families accounted for 4.5% of all births in Korea.⁸ As such, adolescents in multicultural families are likely to be at greater risk for social dysfunction than adolescents in monocultural families.⁹ For instance, peer relationships could be a particularly common problem for multicultural adolescents because of the uncertainty surrounding racial status, which may lead to behavioral and psychosocial problems such as social isolation and delinquent behavior.⁹⁻¹¹ Recent studies have consistently reported that multicultural adolescents tend to have more psychological problems than monoracial adolescents.^{12,13} Because adolescence is the period of ego development and self-esteem formation, these psychological issues warrant more attention.¹⁴

While industrialization and economic development have brought about significant advances in convenience, deteriorating mental health is becoming an increasingly relevant issue in our competitive modern society.¹⁵ The World Health Organization defines health as a state of complete physical, mental, and social well-being and not merely the absence of disease or infirmity.¹⁶ Thus, mental health is considered a notable health issue; depression is a particularly important mental health condition.¹⁶ In fact, depression is a disease of great burden. A study showed that depression is likely to have the highest disease burden worldwide by 2030.¹⁷ In addition, more than 60% of suicides can be attributed to mental health status including depressive disorder and other mood disorders.¹⁸ In particular, Korea has the highest suicide rate among Organisation for Economic Co-operation and Development (OECD) countries (29 suicides per 100,000 persons, in 2014).¹⁹ Thus, depression and its treatment requires attention.²⁰

Although several studies have investigated depression in adolescents from multicultural families, most of them involved Caucasians, and rarely Asians.²¹ Thus, the purpose of this study was to investigate whether a significant association exists between the parents' country of birth and adolescent depressive symptoms in a nationally representative sample of South Korean adolescents.

METHODS

Study population

Data from the Korea Youth Risk Behavior Web-based Survey (KYRBWS) 2012, 2013, 2014, 2015, and 2016 were used for this study. The KYRBWS used a cross-sectional study design and comprised a nationwide web-based survey of Korean adolescent health behaviors. The KYRBWS was the result of collaboration between the Korea Centers for Disease Control and Prevention, Ministry of Health and Welfare, and Ministry of Education. The study design included multistage sampling, stratification, and clustering. The samples were stratified by 44 regional and school type variables and the participants were selected from two sources: school (first sampling unit) and class (second sampling unit). The validity and reliability of the data collected using the questionnaire survey analysis has been approved by several studies.^{22,23} The study population consisted of 74,186 students in 2012, 72,435 students in 2013, 72,060 students in 2014, 68,043 students in 2015 and 65,528 students in 2016 from 400 middle schools and 400 high schools in Korea. Students who did not have a mother or father or both mother and father ($n = 24,894$) were excluded from our study population. After excluding missing data ($n = 1$), our total sample size comprised 327,357 individuals.

Variables

The dependent variable was adolescent depressive symptoms. All participants were asked the question, “Within the last year, did you experience any emotions such as sadness or despair continuously for 2 whole weeks, which was enough to hinder your daily life?” Respondents who answered “Yes” were classified as having depressive symptoms, and those who answered “No” were classified as having no depressive symptoms.

The main independent variable of interest in this study was the parents' country of birth. We examined the parents' country of birth using a questionnaire; the data were subjected to two stages of analyses. At the first stage, the students were asked to state whether the father and/or mother was born in Korea or abroad using a “Yes” or “No” response. Subsequently, the participants were divided into three groups: i.e., both parents were native Korean, only one parent was born abroad, both parents were born abroad. In the second stage of the analysis, the participants were asked, “In which country was your father/mother born?” Accordingly, the participants were classified into six groups on the basis of their fathers' or mothers' country of birth and considering the possibility that both parents were native Korean: Korea, China, North Korea, developing countries, Japan, Taiwan, and Others. Developing countries included Vietnam, Philippines, Mongolia, Thailand, Cambodia, and Malaysia, which ranked below 25 in GDP globally after excluding other country groups.

We controlled for sociodemographic, socioeconomic, health-behavior, and health-condition covariates in the analysis. The sociodemographic factors were grade (middle school student, high school) and gender (male, female). The socioeconomic factors were school achievement (low, middle, high), region (metropolitan, urban, rural area), father's/mother's educational level (high school or less, college or over, unknown), household income level (low, middle, high). Metropolitan areas included capital and metropolitan cities, urban areas included cities except metropolitan areas, and rural areas included the remaining geographic area, except for the metropolitan and urban regions. Health-behavior factors included physical activity (inactive, low, middle, high), smoking status (ever or never), and alcohol consumption (ever or no), Health-condition factors included suicidal thoughts (yes or no), sleep duration for overcoming fatigue (enough, normal, not enough), perceived health status (good, normal, bad), perceived happiness status (good, normal, bad), stress level (high or low) and year (2012, 2013, 2014, 2015, and 2016).

Statistical analysis

First of all, χ^2 tests were used to confirm significant differences in the presence of adolescents' depressive symptoms according to the parents' migration background. We also conducted logistic regression analysis to determine adjusted odds ratios (ORs) and 95% confidence intervals (CIs). Subgroup analysis was performed according to grade, gender, region, household income level, father's educational level, and mother's educational level. Model fitting was conducted using the PROC SURVEYLOGISTIC procedure using SAS software, version 9.4 (SAS Institute, Cary, NC, USA). All analyses were performed by applying strata, cluster, and weight procedures. A P value < 0.05 was considered to indicate a statistically significant result.

RESULTS

Table 1 demonstrates the general characteristics of the study population. Of the 327,357 participants, 88,341 (27.0%) have depressive symptoms and 239,016 (73.0%) do not. The rate of depressive symptoms is higher among adolescents born in families wherein only

Parent Country of Birth and Offspring Depression

Table 1. General characteristics of the study population

Variables	Total	Depressive symptoms		P value
		Yes	No	
Total	327,357 (100.0)	88,341 (27.0)	239,016 (73.0)	
Parents' country of birth				< 0.001
Parents were native Korean	323,934 (99.0)	87,330 (27.0)	236,604 (73.0)	
Only one was born abroad	2,971 (0.9)	813 (27.4)	2,158 (72.6)	
Parents were born abroad	452 (0.1)	198 (43.8)	254 (56.2)	
Grade				< 0.001
Middle school	165,124 (50.4)	40,705 (24.7)	124,419 (75.3)	
High school	162,233 (49.6)	47,636 (29.4)	114,597 (70.6)	
Gender				< 0.001
Male	167,092 (51.0)	36,894 (22.1)	130,198 (77.9)	
Female	160,265 (49.0)	51,447 (32.1)	108,818 (67.9)	
Region				0.047
Metropolitan area	170,132 (52.0)	46,335 (27.2)	123,797 (72.8)	
Urban area	137,551 (42.0)	36,730 (26.7)	100,821 (73.3)	
Rural area	19,674 (6.0)	5,276 (26.8)	14,398 (73.2)	
Household income level				< 0.001
Low	56,386 (17.2)	20,261 (35.9)	36,125 (64.1)	
Middle	157,654 (48.2)	40,195 (25.5)	117,459 (74.5)	
High	113,317 (34.6)	27,885 (24.6)	85,432 (75.4)	
School achievement				< 0.001
Low	35,712 (10.9)	13,081 (36.6)	22,631 (63.4)	
Middle	252,014 (77.0)	66,801 (26.5)	185,213 (73.5)	
High	39,631 (12.1)	8,459 (21.3)	31,172 (78.7)	
Father's educational level				< 0.001
High school or less	114,451 (35.0)	32,373 (28.3)	82,078 (71.7)	
College or over	159,585 (48.7)	43,341 (27.2)	116,244 (72.8)	
Unknown	53,321 (16.3)	12,627 (23.7)	40,694 (76.3)	
Mother's educational level				< 0.001
High school or less	140,787 (43.0)	39,765 (28.2)	101,022 (71.8)	
College or over	134,792 (41.2)	36,550 (27.1)	98,242 (72.9)	
Unknown	51,778 (15.8)	12,026 (23.2)	39,752 (76.8)	
Physical activity				< 0.001
Inactive	78,521 (24.0)	21,854 (27.8)	56,667 (72.2)	
Low	128,422 (39.2)	34,882 (27.2)	93,540 (72.8)	
Middle	72,448 (22.1)	19,075 (26.3)	53,373 (73.7)	
High	47,966 (14.7)	12,530 (26.1)	35,436 (73.9)	
Smoking status				< 0.001
Ever	61,298 (18.7)	22,497 (36.7)	38,801 (63.3)	
Never	266,059 (81.3)	65,844 (24.7)	200,215 (75.3)	
Alcohol consumption				< 0.001
Ever	136,328 (41.6)	45,937 (33.7)	90,391 (66.3)	
Never	191,029 (58.4)	42,404 (22.2)	148,625 (77.8)	
Suicidal thoughts				< 0.001
Yes	45,940 (14.0)	33,484 (72.9)	12,456 (27.1)	
No	281,417 (86.0)	54,857 (19.5)	226,560 (80.5)	
Sleeping duration for overcoming fatigue				< 0.001
Enough	89,334 (27.3)	15,185 (17.0)	74,149 (83.0)	
Normal	106,652 (32.6)	26,063 (24.4)	80,589 (75.6)	
Not enough	131,371 (40.1)	47,093 (35.8)	84,278 (64.2)	
Perceived health status				< 0.001
Good	232,074 (70.9)	52,914 (22.8)	179,160 (77.2)	
Normal	75,082 (22.9)	25,589 (34.1)	49,493 (65.9)	
Bad	20,201 (6.2)	9,838 (48.7)	10,363 (51.3)	
Perceived happiness status				< 0.001
Good	205,088 (62.6)	37,326 (18.2)	167,762 (81.8)	
Normal	91,241 (27.9)	31,563 (34.6)	59,678 (65.4)	
Bad	31,028 (9.5)	19,452 (62.7)	11,576 (37.3)	

(continued to the next page)

Table 1. (Continued) General characteristics of the study population

Variables	Total	Depressive symptoms		P value
		Yes	No	
Stress level				< 0.001
High	125,598 (38.4)	57,822 (46.0)	67,776 (54.0)	
Low	201,759 (61.6)	30,519 (15.1)	171,240 (84.9)	
Year				< 0.001
2012	68,989 (21.1)	20,810 (30.2)	48,179 (69.8)	
2013	66,951 (20.5)	20,338 (30.4)	46,613 (69.6)	
2014	66,955 (20.5)	17,482 (26.1)	49,473 (73.9)	
2015	63,376 (19.4)	14,557 (23.0)	48,819 (77.0)	
2016	61,086 (18.7)	15,154 (24.8)	45,932 (75.2)	

Data shown are number (percentage).

one was born abroad (27.4%) than among adolescents with native Korean parents (27.0%). Furthermore, the rate of depressive symptoms is the highest among adolescents from families wherein both parents were born abroad (43.8%).

Table 2 shows the results for the factors associated with adolescent depressive symptoms. After controlling for covariates, the parents' country of birth is significantly associated with adolescent depressive symptoms. Respondents whose parents were born abroad are more likely to have depressive symptoms than respondents whose parents were native Korean and

Table 2. Factors associated with adolescents' depressive symptoms

Variables	Depressive symptoms	
	Adjusted OR	95% CI
Parents' country of birth		
Parents were native Korean	1.00	-
Only one was born abroad	1.08	0.97-1.20
Parents were born abroad	1.68	1.33-2.12
Grade		
Middle school	1.00	-
High school	1.03	1.01-1.06
Gender		
Male	1.00	-
Female	1.56	1.52-1.59
Region		
Metropolitan area	1.00	0.96-1.05
Urban area	0.98	0.93-1.02
Rural area	1.00	-
Household income level		
Low	1.00	-
Middle	0.90	0.87-0.92
High	1.00	0.97-1.03
School achievement		
Low	1.39	1.33-1.45
Middle	1.19	1.16-1.23
High	1.00	-
Father's educational level		
High school or less	1.00	-
College or over	1.05	1.02-1.07
Unknown	0.93	0.89-0.98
Mother's educational level		
High school or less	1.00	-
College or over	1.08	1.05-1.11
Unknown	0.92	0.88-0.96

(continued to the next page)

Table 2. (Continued) Factors associated with adolescents' depressive symptoms

Variables	Depressive symptoms	
	Adjusted OR	95% CI
Physical activity		
Inactive	0.70	0.67–0.72
Low	0.84	0.81–0.86
Middle	0.94	0.91–0.98
High	1.00	-
Smoking status		
Ever	1.00	-
Never	0.73	0.71–0.75
Alcohol consumption		
Ever	1.00	-
Never	0.74	0.72–0.76
Suicidal thoughts		
Yes	6.15	5.98–6.33
No	1.00	-
Sleeping duration for overcoming fatigue		
Enough	1.00	-
Normal	1.20	1.16–1.23
Not enough	1.37	1.33–1.41
Perceived health status		
Good	1.00	-
Normal	1.13	1.10–1.16
Bad	1.31	1.26–1.36
Perceived happiness status		
Good	1.00	-
Normal	1.42	1.39–1.45
Bad	2.10	2.03–2.18
Stress level		
High	1.00	-
Low	0.41	0.40–0.42
Year		
2012	1.00	-
2013	1.09	1.06–1.13
2014	0.99	0.96–1.03
2015	0.87	0.84–0.90
2016	0.98	0.94–1.01

OR = odds ratio, CI = confidence interval.

respondents with only one was born abroad (only one was born abroad: OR = 1.08; 95% CI, 0.97–1.20; parents were born abroad: OR = 1.68; 95% CI, 1.33–2.12).

Table 3 demonstrates the results of the subgroup analysis of the effect of parents' country of birth on the incidence of adolescent depressive symptoms, stratified by grade, gender, region, household income level, father's educational level and mother's educational level. The analysis stratified by grade revealed that, compared with middle school students, high school students with one or both non-native Korean parents have a greater magnitude of depressive symptoms (only one was born abroad: OR = 1.15; 95% CI, 0.97–1.36; parents were born abroad: OR = 2.04; 95% CI, 1.45–2.87). Overall, the multiculturalism of the family is found to be directly proportional to the magnitude of depressive symptoms in the adolescents.

Table 4 shows a more detailed analysis of the correlation between the parents' country of birth and the adolescents' depressive symptoms. Respondents whose father was North Korean are more likely to have depressive symptoms (father born in North Korea and mother was native Korean: OR = 4.70; 95% CI, 1.27–17.44) compared to respondents from other groups. Also, adolescents whose father was from Japan or Taiwan are more likely to be have depressive

Table 3. Subgroup analysis of depressive symptoms with parents' country of birth by demographic factors

Variables	Parents' country of birth					
	Parents were native Korean		Only one was born abroad		Parents were born abroad	
	OR		OR	95% CI	OR	95% CI
Grade						
Middle school	1.00		1.02	0.88–1.17	1.40	1.01–1.94
High school	1.00		1.15	0.97–1.36	2.04	1.45–2.87
Gender						
Male	1.00		1.09	0.93–1.28	1.65	1.22–2.24
Female	1.00		1.06	0.92–1.23	1.72	1.19–2.48
Region						
Metropolitan area	1.00		1.06	0.89–1.25	1.63	1.19–2.23
Urban area	1.00		1.08	0.92–1.27	1.67	1.15–2.42
Rural area	1.00		1.12	0.81–1.56	2.38	0.77–7.29
Household income level						
Low	1.00		1.05	0.87–1.26	1.50	0.98–2.30
Middle	1.00		1.00	0.84–1.18	2.00	1.33–3.00
High	1.00		1.26	1.00–1.58	1.61	1.11–2.35
Father's educational level						
High school or less	1.00		1.06	0.91–1.24	1.69	1.18–2.43
College or over	1.00		1.14	0.92–1.41	1.55	1.01–2.36
Unknown	1.00		1.00	0.81–1.24	1.75	1.08–2.83
Mother's educational level						
High school or less	1.00		1.08	0.90–1.28	1.88	1.33–2.66
College or over	1.00		1.10	0.92–1.32	1.44	0.96–2.16
Unknown	1.00		1.00	0.81–1.24	1.49	0.90–2.49

OR = odds ratio, CI = confidence interval.

Table 4. Adolescents' depressive symptoms associated with parents' country of birth^a

Variables	Total, No.	Depressive symptoms	
		Adjusted OR ^a	95% CI
Father's country of birth (under mother was native Korean)			
Korea	323,934	1.00	-
China	54	0.93	0.45–1.93
North Korea	15	4.70	1.27–17.44
Developing countries	23	1.85	0.44–7.73
Japan, Taiwan	119	1.79	1.15–2.77
Others	95	1.94	1.20–3.15
Mother's country of birth (under father was native Korean)			
Korea	323,934	1.00	-
China	1,084	0.90	0.75–1.08
North Korea	18	2.79	0.92–8.48
Developing countries	605	1.31	1.02–1.70
Japan, Taiwan	866	0.97	0.78–1.20
Others	92	0.79	0.47–1.34

OR = odds ratio, CI = confidence interval.

^aAdjusted by grade, gender, region, household income level, school achievement, father's educational level, mother's educational level, physical activity, smoking status, alcohol consumption, suicidal thoughts, sleeping duration for overcoming fatigue, perceived health status, perceived happiness status, stress level, year.

symptoms than other groups (OR = 1.49; 95% CI, 1.15–2.77). Additionally, depressive symptoms are observed more frequently in adolescents with mother was from a developing country and father was native Korean (OR = 1.31; 95% CI, 1.02–1.70) than other adolescents.

Finally, in **Supplementary Table 1**, the general characteristics of the study population are demonstrated according to parental country of birth. We found that father's educational level and mother's educational level are higher among native Korean parents than among parents born abroad. Furthermore, the frequency of low household income level is 17.2% in the families with native Korean parents and 34.6% in families wherein the parents were born abroad group.

DISCUSSION

The aim of this study was to identify significant correlations between the parents' country of birth and adolescent depressive symptoms in Korea. The findings revealed that, compared with adolescents whose parents are both native Korean, adolescents whose parents were born abroad are more likely to have depressive symptoms. In high school students, the parents' country of birth has a greater influence on the existence of depressive symptoms than that in middle school students. By subdividing the parents' country of birth, we found that if father was born in North Korea or Japan or Taiwan and mother was native Korean, the adolescents are more likely to have depressive symptoms. When the mother's country of birth was developing countries, adolescents are more likely to have depressive symptoms.

In this study, adolescents whose parents were born abroad are more likely to have depressive symptoms than adolescents whose parents were both native Korean. This could be explained by several reasons. Adolescents of immigrants experience many conflicts and unfamiliar situations as they grow in a dual culture where the values and attitudes of their fathers and mothers are different.²⁴⁻²⁷ Moreover, because the adolescents experience different lifestyles at home and in society, they often experience identity crises and value conflicts. Bilingual usage within the family could lead to difficulty with language competence. Additionally, adolescents of immigrants were more likely to experience poor parenting and family functioning and discrimination such as school violence. These factors can affect the mental health of multicultural adolescents.²⁴⁻²⁷ In addition, we found that low parental education levels and low household income were more commonly observed among multicultural families. Several previous studies have reported that lower parental education levels are associated with a higher probability of impaired mental health in the child. Moreover, children belonging to families with low household income levels are more likely to be depressed.^{28,29}

The present study shows that high school students demonstrated a greater magnitude of depressive symptoms than did middle school students, when analyzed according to the parents' birth country. In Korea, where educational success is very important, in order to go to university, high school students have to take an entrance examination.³⁰ High school students therefore experience enormous amounts of educational success; several reports of high school students committing suicide owing to poor entrance examination performance have been reported.³⁰ Because high school students have vulnerable mental health owing to this educational success, they may be more influenced by their parents' birth of country.

In multicultural families, adolescents with father was from North Korea are more likely to have depressive symptoms. This may be associated with the traumatic events experienced while escaping from North Korea to South Korea. People who escaped from North Korea to South Korea were often crossing through China or other Asian countries.³¹ If arrested during this process, the people were deported to North Korea again and would face severe punishment or a death penalty. These traumatic events may cause serious mental health problems for North Korean defectors. Moreover, North Korean defectors commonly experience discrimination and political problems in South Korea.^{31,32} Therefore, adolescents with one North Korean father may experience mental health problems owing to the unstable parenting and home environment.

Adolescents with their mother from a developing country tend to have depressive symptoms more frequently than other adolescents. It is necessary to consider that many immigrants from developing countries are migration workers.^{32,33} Most migration workers are given dirty, dangerous, difficult, and low-paying jobs that South Koreans refuse. Therefore, migration workers are likely to maintain a low economic level. In addition, there were many international marriages between women from developing countries and men who were not married to native Korean women. These families tended to maintain a low economic level; in particular, the women who migrated had several difficulties adapting to the South Korean culture.^{32,33} Furthermore, people from low-income countries experienced more severe discriminations than people from high-income countries.³⁴ Thus, adolescents with their mother from developing countries were more likely to experience lower incomes and several discriminations. These factors may have contributed to the adolescents' depressive symptoms.

This study had some limitations. First, the data were self-reported by the participants. It is possible that the responses did not match the actual depressive symptoms rate. Second, we could not examine several parental factors such as their characters and domestic violence exposure. Parental characteristics are known as important factors that determine the adolescent's mental health.^{35,36} Third, the study was based on a cross-sectional survey. Causality could not be confirmed clearly and the association could be confirmed. These limitations will be considered in a future study. Despite the above limitations, this study also had strengths. First, this study used the most recent nationally, multistage, stratified collected data. Therefore, the results obtained are representative of Korean adolescents. Second, although most previous studies were conducted in Caucasians who already have formed a multicultural society, we conducted research in Asian society, which is experiencing the early stages of a multicultural society. Third, to the best of our knowledge, this study offers new insights into the association between the parents' country of birth and adolescent depressive symptoms rates.

In conclusion, in our investigation of the correlation between the parents' country of birth and adolescent depressive symptoms, we found that, compared with adolescents whose parents were both native Korean, adolescents whose parents were born abroad are more likely to have depressive symptoms. Korea is in the early stages of forming a multicultural society; therefore, the number of studies on multicultural adolescents and mental health in Korea is limited. From a long-term point of view, mental health examinations for adolescents from multicultural families should be conducted, and adolescent mental health should be given attention. In addition, the existing multicultural policy does not include a classification according to the parent's country of origin; the immigrant populations with various nationalities are included in a single policy. The findings of our study confirm that the rates of depressive symptoms differ according to the parent's country of birth. Further studies should investigate the relationship between detailed characteristics of the parent's country of birth and their offspring's mental health. In addition, multicultural family support policies should be implemented, considering the characteristics of the parent's country of birth. Also, schools, government, and communities should propose and implement the multicultural education policies so that the growing youth could understand and accept the multicultural society. Overall, the findings also demonstrate what could happen in an early multicultural era. The results could be regarded as a prequel to the current situation in advanced multicultural societies such as the United States and Canada,³⁷⁻³⁹ and could have implications in many countries experiencing the early stages of a multicultural society.⁴⁰

ACKNOWLEDGMENTS

We thank the Korea Centers for Disease Control and Prevention, Ministry of Health and Welfare, and Ministry of Education that provided data.

SUPPLEMENTARY MATERIAL

Supplementary Table 1

General characteristic of study population by parental country of birth

[Click here to view](#)

REFERENCES

- Arnett JJ. The psychology of globalization. *Am Psychol* 2002;57(10):774-83.
[PUBMED](#) | [CROSSREF](#)
- Chen SX, Benet-Martinez V, Harris Bond M. Bicultural identity, bilingualism, and psychological adjustment in multicultural societies: immigration-based and globalization-based acculturation. *J Pers* 2008;76(4):803-38.
[PUBMED](#) | [CROSSREF](#)
- Castles S, de Haas HG, Miller MJ. *The Age of Migration: International Population Movements in the Modern World*. 5th ed. New York, NY: Palgrave Macmillan; 2013.
- International migrant stock 2015. <http://www.un.org/en/development/desa/population/migration/data/estimates2/estimates15.shtml>. Updated 2015. Accessed March 15, 2017.
- Population, households and housing units. http://kosis.kr/statHtml/statHtml.do?orgId=101&tblId=DT_1IN1502&conn_path=I2&language=en. Updated 2017. Accessed March 16, 2017.
- Lee HK. International marriage and the state in South Korea: focusing on governmental policy. *Citizensh Stud* 2008;12(1):107-23.
[CROSSREF](#)
- Marriages to foreigners for city, county, and district. http://kosis.kr/statHtml/statHtml.do?orgId=101&tblId=DT_1B83A24&conn_path=I2&language=en. Updated 2017. Accessed April 19, 2017.
- Multicultural Live live births by province, si(city), gun(county) and gu(borough). http://kosis.kr/statHtml/statHtml.do?orgId=101&tblId=DT_1BB0006&conn_path=I2&language=en. Updated 2017. Accessed April 20, 2016.
- Gibbs JT, Moskowitz-Sweet G. Clinical and cultural issues in the treatment of biracial and bicultural adolescents. *Fam Soc* 1991;72(10):579-92.
- Brown PM. Biracial identity and social marginality. *Child Adolesc Social Work J* 1990;7(4):319-37.
[CROSSREF](#)
- Gibbs JT, Huang LN. *Children of Color: Psychological Interventions with Minority Youth*. San Francisco, CA: Jossey-Bass; 1991.
- Cheng S, Lively KJ. Multiracial self-identification and adolescent outcomes: a social psychological approach to the marginal man theory. *Soc Forces* 2009;88(1):61-98.
[CROSSREF](#)
- Udry JR, Li RM, Hendrickson-Smith J. Health and behavior risks of adolescents with mixed-race identity. *Am J Public Health* 2003;93(11):1865-70.
[PUBMED](#) | [CROSSREF](#)
- Allen JP, Hauser ST, Bell KL, O'Connor TG. Longitudinal assessment of autonomy and relatedness in adolescent-family interactions as predictors of adolescent ego development and self-esteem. *Child Dev* 1994;65(1):179-94.
[PUBMED](#) | [CROSSREF](#)
- Desjarlais R. *World Mental Health: Problems and Priorities in Low-income Countries*. New York, NY: Oxford University Press; 1995.

16. World Health Organization. *Preamble to the Constitution of the World Health Organization as Adopted by the International Health Conference, New York, 19–22 June, 1946*. Geneva, Switzerland: World Health Organization; 1948.
17. Mathers C, Fat DM, Boerma JT. *The Global Burden of Disease: 2004 Update*. Geneva, Switzerland: World Health Organization; 2008.
18. Cavanagh JT, Carson AJ, Sharpe M, Lawrie SM. Psychological autopsy studies of suicide: a systematic review. *Psychol Med* 2003;33(3):395-405.
[PUBMED](#) | [CROSSREF](#)
19. Organisation for Economic Co-operation and Development. *Society at a Glance 2016: OECD Social Indicators*. Paris, France: OECD Publishing; 2016.
20. Kim JL, Cho J, Park S, Park EC. Depression symptom and professional mental health service use. *BMC Psychiatry* 2015;15:261.
[PUBMED](#) | [CROSSREF](#)
21. Campbell ME, Eggerling-Boeck J. "What about the children?" The psychological and social well-being of multiracial adolescents. *Sociol Q* 2006;47(1):147-73.
[CROSSREF](#)
22. Bae J, Joung H, Kim JY, Kwon KN, Kim Y, Park SW. Validity of self-reported height, weight, and body mass index of the Korea Youth Risk Behavior Web-based Survey questionnaire. *J Prev Med Public Health* 2010;43(5):396-402.
[PUBMED](#) | [CROSSREF](#)
23. Bae J, Joung H, Kim JY, Kwon KN, Kim YT, Park SW. Test-retest reliability of a questionnaire for the Korea Youth Risk Behavior Web-based Survey. *J Prev Med Public Health* 2010;43(5):403-10.
[PUBMED](#) | [CROSSREF](#)
24. Stevens GW, Vollebergh WA. Mental health in migrant children. *J Child Psychol Psychiatry* 2008;49(3):276-94.
[PUBMED](#) | [CROSSREF](#)
25. Bursztein Lipsicas C, Henrik Mäkinen I. Immigration and suicidality in the young. *Can J Psychiatry* 2010;55(5):274-81.
[PUBMED](#) | [CROSSREF](#)
26. Kim JH, Kim JY, Kim SS. School violence, depressive symptoms, and help-seeking behavior: a gender-stratified analysis of biethnic adolescents in South Korea. *J Prev Med Public Health* 2016;49(1):61-8.
[PUBMED](#) | [CROSSREF](#)
27. Bourque F, van der Ven E, Malla A. A meta-analysis of the risk for psychotic disorders among first- and second-generation immigrants. *Psychol Med* 2011;41(5):897-910.
[PUBMED](#) | [CROSSREF](#)
28. Angold A. Childhood and adolescent depression. I. Epidemiological and aetiological aspects. *Br J Psychiatry* 1988;152(5):601-17.
[PUBMED](#) | [CROSSREF](#)
29. Goodman E, Slap GB, Huang B. The public health impact of socioeconomic status on adolescent depression and obesity. *Am J Public Health* 2003;93(11):1844-50.
[PUBMED](#) | [CROSSREF](#)
30. Lee S, Hong JS, Espelage DL. An ecological understanding of youth suicide in South Korea. *Sch Psychol Int* 2010;31(5):531-46.
[CROSSREF](#)
31. Jeon W, Hong C, Lee C, Kim DK, Han M, Min S. Correlation between traumatic events and posttraumatic stress disorder among North Korean defectors in South Korea. *J Trauma Stress* 2005;18(2):147-54.
[PUBMED](#) | [CROSSREF](#)
32. Moon S. Multicultural and global citizenship in a transnational age: the case of South Korea. *Int J Multicult Educ* 2010;12(1):1-15.
[CROSSREF](#)
33. Kim AE. Global migration and South Korea: foreign workers, foreign brides and the making of a multicultural society. *Ethn Racial Stud* 2009;32(1):70-92.
[CROSSREF](#)
34. Borrell C, Muntaner C, Gil-González D, Artazcoz L, Rodríguez-Sanz M, Rohlfs I, et al. Perceived discrimination and health by gender, social class, and country of birth in a Southern European country. *Prev Med* 2010;50(1-2):86-92.
[PUBMED](#) | [CROSSREF](#)
35. Hagan MJ, Roubinov DS, Adler NE, Boyce WT, Bush NR. Socioeconomic adversity, negativity in the parent child-relationship, and physiological reactivity: an examination of pathways and interactive processes affecting young children's physical health. *Psychosom Med* 2016;78(9):998-1007.
[PUBMED](#) | [CROSSREF](#)

36. Smith A, Lalonde RN, Johnson S. Serial migration and its implications for the parent-child relationship: a retrospective analysis of the experiences of the children of Caribbean immigrants. *Cultur Divers Ethnic Minor Psychol* 2004;10(2):107-22.
[PUBMED](#) | [CROSSREF](#)
37. Berry JW. Multicultural policy in Canada: a social psychological analysis. *Can J Behav Sci* 1984;16(4):353-70.
[CROSSREF](#)
38. Alba R. Immigration and the American realities of assimilation and multiculturalism. *Sociol Forum (Randolph N J)* 1999;14(1):3-25.
[CROSSREF](#)
39. Joppke C. Multiculturalism and immigration: a comparison of the United States, Germany, and Great Britain. *Theory Soc* 1996;25(4):449-500.
[CROSSREF](#)
40. Vertovec S. Introduction: new directions in the anthropology of migration and multiculturalism. *Ethn Racial Stud* 2007;30(6):961-78.
[CROSSREF](#)