



Klebsiella pneumoniae Carbapenemase Producers in South Korea between 2013 and 2015

Eun-Jeong Yoon¹, Jung Ok Kim¹, Dokyun Kim¹, Hyukmin Lee¹, Ji Woo Yang²,
Kwang Jun Lee^{2*} and Seok Hoon Jeong^{1*}

¹ Department of Laboratory Medicine, Yonsei University College of Medicine, Seoul, South Korea, ² Centers for Disease Control and Prevention, National Institute of Health, Cheongju, South Korea

OPEN ACCESS

Edited by:

Dongsheng Zhou,
Beijing Institute of Microbiology and
Epidemiology, China

Reviewed by:

Mariagrazia Perilli,
University of L'Aquila, Italy
Amy J. Mathers,
University of Virginia, United States

*Correspondence:

Kwang Jun Lee
kwangjun@korea.kr
Seok Hoon Jeong
kscpjsh@yuhs.ac

Specialty section:

This article was submitted to
Infectious Diseases,
a section of the journal
Frontiers in Microbiology

Received: 16 October 2017

Accepted: 10 January 2018

Published: 25 January 2018

Citation:

Yoon E-J, Kim JO, Kim D, Lee H,
Yang JW, Lee KJ and Jeong SH
(2018) *Klebsiella pneumoniae*
Carbapenemase Producers in South
Korea between 2013 and 2015.
Front. Microbiol. 9:56.
doi: 10.3389/fmicb.2018.00056

Between 2014 and 2015, the *Klebsiella pneumoniae* carbapenemase (KPC) was becoming endemic in South Korea. To assess this period of transition, we analyzed KPC producers in terms of molecular epidemiology. A total of 362 KPC-producing *Enterobacteriaceae* strains, including one from 2013, 13 from 2014, and 348 from 2015, were actively collected from 60 hospitals throughout the peninsula. Subtypes of KPC were determined by PCR and direct sequencing, and isotypes of Tn4401 (the transposon flanking the *bla*_{KPC} gene) were specified by PCR using isotype-specific primers and direct sequencing. Sporadic occurrence of KPC-producing *Enterobacteriaceae* was initially observed around Seoul, which is the most crowded district of the country, and these strains rapidly disseminated in 2014, to the other parts of the country in 2015. The bacterial clones responsible for the extreme epidemiological transition were *K. pneumoniae* ST307 (46.2%) and ST11 (21.3%). Less frequently, *E. coli* (4.7%), *Enterobacter* spp. (1.4%), and other *Enterobacteriaceae* members (1.7%) producing the enzyme were identified. The *bla*_{KPC-2} gene bracketed by Tn4401a (72.1%) was the most prevalent mobile genetic element responsible for the dissemination, and the same gene carried either by Tn4401b (2.2%) or Tn4401c (6.6%) was identified at a lesser frequency. The genes *bla*_{KPC-3} (1.6%) and *bla*_{KPC-4} (6.4%), both flanked by Tn4401b, were occasionally identified. This study showed endemic dissemination of KPC producers in 2015 due to a clonal spread of two *K. pneumoniae* strains. Further systemic surveillance is needed to monitor dissemination of KPC-producers.

Keywords: KPC, Tn4401, carbapenem resistance, *Klebsiella pneumoniae* ST307, carbapenemase producing *Enterobacteriaceae*

INTRODUCTION

Carbapenemases often jeopardize chemotherapy for infectious diseases caused by common nosocomial pathogens. Currently, *Klebsiella pneumoniae* carbapenemases (KPC) belonging to the Ambler class A carbapenemases are the most clinically worrisome enzymes because of their global spreads and a broad-spectrum of substrates including most of the β -lactam drugs, except cephamycins (Munoz-Price et al., 2013; Pitout et al., 2015). KPC producers were first identified for *K. pneumoniae* from the United States in 1996 (Yigit et al., 2001) and, so far, a total of 23 KPC variants have been identified (<http://www.lahey.org/Studies>). KPC producers, particularly among *Enterobacteriaceae* isolates, are now being extensively reported worldwide. In some regions of the

world, such as North America (some regions of the USA), Latin America (Colombia, Brazil, and Argentina), Europe (Greece and Italy), the Middle East (Israel), and Asia (some regions of China), KPC dissemination has reached endemic levels (Albiger et al., 2015; Mehta et al., 2015).

The rapid dissemination of the *bla*_{KPC} gene is known to be derived from the clonal spread of *K. pneumoniae* sequence type 258 (ST258) (Munoz-Price et al., 2013; Pitout et al., 2015). Besides, horizontal gene transfer drives KPC dissemination via a Tn3-type active transposon Tn4401 enclosing the *bla*_{KPC} gene. The transposon is bracketed by 39-bp imperfect inverted repeat sequences capable of mobilizing the *bla*_{KPC} gene at high frequency and it generates a 5-bp target duplication by insertion without target site specificity (Cuzon et al., 2011). At present, nine isotypes of Tn4401 (Tn4401a to Tn4401i) differing by the upstream sequences of the *bla*_{KPC} gene were identified. The *bla*_{KPC} gene promoters varied by the Tn4401 isotypes control the level of gene expression resulting in varied levels of carbapenem resistance (Naas et al., 2012; Cheruvanky et al., 2017).

The National Laboratory Surveillance System (NLSS) for CPEs started in November 2010 by the Korea Center for Disease Prevention and Control (KCDC) with 100 sentinel general hospitals (unpublished data). Through the surveillance study, an alarming transition of KPC epidemiological stages to inter-regional spread was observed between 2014 and 2015. The population of KPC producers and the inter-regional movement of core clones needed to be further investigated and, in this study, the molecular epidemiology was assessed with actively collected KPC producers located throughout the peninsula.

MATERIALS AND METHODS

Strains

A total of 362 non-duplicate KPC-producing *Enterobacteriaceae* strains (one from 2013, 13 from 2014, and 348 from 2015) were collected voluntarily from 60 hospitals: 16 NLSS-non-participating hospitals including 12 general hospitals, two convalescent hospitals, and two clinics, and 44 NLSS-participating general hospitals. Further testing including the species identification by 16S rDNA sequencing was performed in a reference laboratory.

Antimicrobial Susceptibility and Resistance Analyses

Antimicrobial susceptibility to eight drugs (aztreonam, cefotaxime, ceftazidime, cefoxitin, gentamicin, amikacin, ciprofloxacin, and tigecycline) was evaluated by the disk diffusion method on Mueller-Hinton (MH) agar (Difco Laboratories, Detroit, MI), following CLSI guidelines (Clinical and Laboratory Standards Institute, 2015b). The MICs for colistin were assessed by the broth microdilution method using MH broth (Difco Laboratories), following the recommendations of the Joint CLSI-EUCAST Polymyxin Breakpoints Working Group (The joint CLSI-EUCAST and Polymyxin Breakpoints Working Group, 2016). The MICs for imipenem and meropenem were determined by the agar dilution method, following CLSI guidelines (Clinical and Laboratory Standards Institute, 2015a).

Both *Escherichia coli* ATCC 25922 and *Pseudomonas aeruginosa* ATCC 27853 were used for quality control. The *bla*_{KPC} gene was detected by PCR using gene-specific primers (Supplementary Table), and direct sequencing was carried out for subtyping KPC. Additional resistance determinants were identified by PCR and direct sequencing: *bla*_{CTX-M-1-like}, *bla*_{CTX-M-9-like}, and *bla*_{CTX-M-25-like} for CTX-M-type extended-spectrum β -lactamases (ESBLs); *bla*_{ACT}, *bla*_{ACC}, *bla*_{CMY}, and *bla*_{DHA} for plasmid-mediated AmpCs; and *armA*, *rmtA*, and *rmtB* for 16S ribosomal methyltransferases (Perez-Perez and Hanson, 2002; Ryoo et al., 2005). Multiple-carbapenemase producers of positive amplification either of *bla*_{OXA-48-like}, *bla*_{IMP}, *bla*_{VIM}, *bla*_{GES}, and *bla*_{NDM} genes were eliminated from the study.

Isotyping the Tn4401 Transposon

To determine the isotypes of Tn4401, isotype-specific forward primers for the five most common isotypes (*a* to *e*) (Naas et al., 2012) were newly designed (Supplementary Table) and PCR was carried out with a universal reverse primer targeting the *bla*_{KPC} gene. The amplicons were sequenced for verification.

Multilocus Sequence Typing (MLST)

PCR and sequencing were carried out for seven housekeeping genes: *gapA*, *infB*, *mdh*, *pgi*, *phoE*, *rpoB*, and *tonB* for *K. pneumoniae* (Diancourt et al., 2005), and *adk*, *fumC*, *gyrB*, *icd*, *mdh*, *purA*, and *recA* for *E. coli* (Wirth et al., 2006). The nucleotide sequences obtained for both DNA strands were compared with sequences in the MLST database for each species (<http://bigsd.b.pasteur.fr/klebsiella> for *K. pneumoniae* and <http://mlst.warwick.ac.uk/mlst/dbs/Ecoli> for *E. coli*), to determine allelic numbers and STs.

RESULTS

The Epidemiology of KPC-Producing *Enterobacteriaceae*

Among the 362 KPC-producing *Enterobacteriaceae* strains, *K. pneumoniae* (334/362, 92.2%) was the predominant bacterial species and *E. coli* (17/362, 4.7%) was followed by *Enterobacter* spp. (5/362, 1.4%), *Citrobacter* spp. (3/362, 0.8%), *Klebsiella oxytoca* (1/362, 0.3%), *Serratia marcescens* (1/362, 0.3%), and *Hafnia alvei* (1/362, 0.3%) were identified less frequently.

Three subtypes KPC-2 (334/362, 92.3%), KPC-3 (6/362, 1.7%), and KPC-4 (22/362, 6.1%) were identified. Major isotype of the Tn4401 bracketing the *bla*_{KPC} gene was Tn4401a (261/362, 72.0%) and, less frequently, Tn4401b (36/362, 9.9%) and Tn4401c (24/362, 6.6%) were identified. Tn4401b was linked with all three subtypes of the *bla*_{KPC} gene, while Tn4401a and Tn4401c carried only the *bla*_{KPC-2} gene. For the non-isotypable 41 transposons (11.3%) by the Tn4401-isotyping PCR, amplification between the 3'-end of the upstream ISKpn7 and the *bla*_{KPC} gene was tried and only one of the 41 was identified having a truncated Tn3 at the isotype-determining region, while none produced any amplicons. The non-isotype-determined (ND) Tn4401s were coupled only with *bla*_{KPC-2} gene, whether in *K. pneumoniae* (*n* = 38) or in *E. coli* (*n* = 3).

The *bla*_{KPC-2} gene bracketed by Tn4401a (Tn4401a[*bla*_{KPC-2}]) was harbored by *K. pneumoniae* (*n* = 243) of varied STs [mainly ST307 (*n* = 155), ST11 (*n* = 37), and ST789 (*n* = 21)], *E. coli* (*n* = 11) [mostly of ST1624 (*n* = 7)], and other *Enterobacteriaceae* [*Enterobacter* spp. (*n* = 2), *C. freundii* (*n* = 2), *C. koseri* (*n* = 1), *K. oxytoca* (*n* = 1), and *S. marcescens* (*n* = 1)] (Table 1). Tn4401b[*bla*_{KPC-2}] was carried either by *K. pneumoniae* (*n* = 7), five ST11 and two ST307, or by *Enterobacter* spp. (*n* = 1). Tn4401c[*bla*_{KPC-2}] was carried only by *K. pneumoniae* (*n* = 24) belonging to ST307 (*n* = 8), ST789 (*n* = 6), ST11 (*n* = 5), ST48 (*n* = 3), ST392 (*n* = 1), and ST2230 (*n* = 1). The Tn4401b[*bla*_{KPC-3}] was identified in *E. coli* ST393 (*n* = 2) and in *K. pneumoniae* (*n* = 4, specifically two ST314 and one each of ST15 and ST697) and Tn4401b[*bla*_{KPC-4}] was identified in *K. pneumoniae* (specifically five ST392, four ST101, three ST11, two ST307, and one of each ST4, ST48, ST54, and ST627), *E. coli* ST2847 (*n* = 1), *Enterobacter* spp. (*n* = 2), and *Hafnia alvei* (*n* = 1).

Dissemination of KPC Producers throughout the South Korean Peninsula from 2013 to 2015

One *K. pneumoniae* isolate ST2230/Tn4401c[*bla*_{KPC-2}] was collected in Seoul in 2013. In 2014, 13 KPC producers were identified including six *K. pneumoniae* strains, i.e., two ST307/Tn4401a[*bla*_{KPC-2}], one ST11/ND[*bla*_{KPC-2}], one ST392/Tn4401c[*bla*_{KPC-2}], one ST392/Tn4401b[*bla*_{KPC-4}], and one ST15/Tn4401b[*bla*_{KPC-3}], along with seven *E. coli* ST1642/Tn4401a[*bla*_{KPC-2}] strains from a single-hospital outbreak in a suburb of Seoul (Incheon). In 2015, a total of 348 KPC producers were identified including 327 *K. pneumoniae* [mainly ST307/Tn4401a[*bla*_{KPC-2}] (153/327, 46.8%), ST11/Tn4401a[*bla*_{KPC-2}] (37/327, 11.3%), ST25/ND[*bla*_{KPC-2}] (25/327, 7.6%), and ST789/Tn4401a[*bla*_{KPC-2}] (21/327, 6.4%)] and 10 *E. coli* strains of nine subclones. Notably, five KPC-producing *K. pneumoniae* isolates were collected in 2015 from other healthcare facilities than general hospitals: ST307/Tn4401a[*bla*_{KPC-2}], ST11/Tn4401a[*bla*_{KPC-2}], and ST290/Tn4401a[*bla*_{KPC-2}] from two convalescent hospitals and ST48/Tn4401c[*bla*_{KPC-2}] and ST307/Tn4401a[*bla*_{KPC-2}] from two clinics.

The *K. pneumoniae* ST307/Tn4401a[*bla*_{KPC-2}], ST11/ND[*bla*_{KPC-2}], and ST392/Tn4401b[*bla*_{KPC-4}] clones were identified over two consecutive years. In 2014, two *K. pneumoniae* ST307/Tn4401a[*bla*_{KPC-2}] were first identified in Seoul. In 2015, the same clone was identified with 153 strains in Seoul (*n* = 49) and three more districts including a suburb of Seoul (Gyeonggi, *n* = 12), a southeastern peninsula (Busan, *n* = 91), and a southwestern peninsula (Jeonbuk, *n* = 1; Figure 1). Besides, the *K. pneumoniae* ST11/ND[*bla*_{KPC-2}] clone was first identified in Seoul in 2014 and, in 2015, the same clone was identified in Seoul (*n* = 12), two suburbs of Seoul (Gyeonggi, *n* = 1; Incheon, *n* = 1), and in southeastern peninsula (Gyeongbuk, *n* = 2; Busan, *n* = 1). The *K. pneumoniae* ST392/Tn4401b[*bla*_{KPC-4}] clone from Incheon in 2014 was

subsequently identified with four isolates in 2015 from the same district.

Carbapenem Resistance Conferred by the *bla*_{KPC} Gene

Overall MIC₅₀s of imipenem and meropenem in KPC producers were 16 mg/L and 8 mg/L, respectively (Figure 2), and the values met the three predominant factors in the KPC populations: the host bacterial species (*K. pneumoniae*), the resistance gene (*bla*_{KPC-2}), and the transposon encoding the gene (Tn4401a). The KPC-producing *K. pneumoniae* had three-dilution higher MIC₅₀s of the two carbapenems than *E. coli* and KPC-2 producers had three- and four-dilution higher MIC₅₀ values than to the KPC-3 and KPC-4 producers, respectively. The KPC producers carrying Tn4401a had one- or two-dilution higher MIC₅₀ values of the carbapenemase than KPC producers carrying Tn4401b, whereas those carrying Tn4401c had the same MIC₅₀ values as KPC producers carrying Tn4401a. The ND transposon encoding *bla*_{KPC-2} (*n* = 40) had the highest MIC₅₀ of meropenem 32 mg/L that was two- to three-dilution higher than the three isotypes of Tn4401, whereas that of imipenem looked alike.

Cross- and Co-resistance to Other Drugs

As a result of cross resistance from producing KPC, all or most of the 362 KPC producers were non-susceptible to aztreonam (100%), cefotaxime (100%), ceftazidime (*n* = 359, 99.2%), and cefoxitin (*n* = 325, 89.8%) (Table 2).

A total of 192 KPC producers (53.0%) produced at least one CTX-M-type ESBL (Table 3): 46.7% (169/362) CTX-M type 1 and 8.0% (29/362) CTX-M type 9. Of note, the *bla*_{CTX-M} gene was often possessed by KPC-2 (183/334, 54.8%) and KPC-4 (9/22, 40.9%) producers, whereas none were identified in KPC-3 producers. Plasmid-mediated AmpC was rarely co-produced (13/362, 3.4%) by the KPC producers: DHA (*n* = 9), ACT (*n* = 1), or CMY (*n* = 1) production in KPC-2 producers and ACC (*n* = 2) production in KPC-4 producers.

Among the 362 KPC producers, 220 (60.8%) and 46 (12.7%) were resistant to gentamicin and amikacin, respectively, but only few KPC-2 producers possessed the 16S ribosomal methyltransferase *armA* (1/362, 0.3%) and *rmtB* (22/362, 6.1%) genes.

A total of 348 isolates (96.7%) were non-susceptible to ciprofloxacin and 80 (22.1%) and eight (2.2%) isolates were non-susceptible to tigecycline and colistin, respectively. None of the eight colistin-resistant strains possessed the *mcr* gene.



DISCUSSION

KPC enzymes are able to hydrolyze penicillins, cephalosporins, monobactams, carbapenems, and even β -lactamase inhibitors. Therefore, the wide spread of the producers is one of the worrisome global issues encountering antimicrobial resistance. In accordance of occasional reports (Rhee et al., 2010; Hong et al., 2013; Yoo et al., 2013) and data from NLSS (unpublished data), only infrequent occurrence of the KPC producers were identified in South Korea until 2013. However, after a sudden expansion of KPC producers in 2014, rapid nation-wide spread supported

TABLE 1 | The *bla*_{KPC} mobile elements carried by KPC producers in South Korea.

	<i>E. coli</i> belonging to ST										<i>K. pneumoniae</i> belonging to ST										<i>Enterobacter</i> spp.		<i>C. freundii</i>	<i>C. koseri</i>	<i>K. oxytoca</i>	<i>S. marcescens</i>	<i>H. alvei</i>	Total						
	1642	131	2448	3306	4683	1193	410	393	2847	307	11	789	48	392	290	101	2538	29	314	524	628	15							1035	1393	1413	147	17	2230
2013																																		
KPC-2																																		
Tn4401c																																		
2014																																		
KPC-2																																		
Tn4401a																																		
Tn4401c																																		
ND																																		
KPC-3																																		
Tn4401b																																		
KPC-4																																		
Tn4401b																																		
2015																																		
KPC-2																																		
Tn4401a																																		
Tn4401b																																		
Tn4401c																																		
ND																																		
KPC-3																																		
Tn4401b																																		
KPC-4																																		
Tn4401b																																		
Total	7	2	1	1	1	1	1	1	2	1	167	77	34	14	7	6	6	3	2	2	2	1	1	1	1	1	1	1	1	1	1	1	362	

Numbers in *italics*, sum of the sub-titled values.



by single-hospital outbreaks (Kim et al., 2017), frequently associated with inter-hospital or inter-regional spread (Jeong et al., 2016) led the endemic stage of KPC in 2015. Actually,

the KPC producer in 2013 has never been identified again and the four *K. pneumoniae* subclones, ST307/Tn4401a[*bla*_{KPC-2}], ST307/Tn4401b[*bla*_{KPC-4}], ST392/Tn4401b[*bla*_{KPC-4}], and

TABLE 2 | Resistance phenotypes of KPC producers.

Total		Number (%) of non-susceptible strains to								
		Aztreonam	Cefotaxime	Ceftazidime	Cefoxitin	Gentamicin	Amikacin	Ciprofloxacin	Tigecycline	Colistin
KPC-2	334	334 (100)	334 (100)	332 (99.4)	305 (91.3)	203 (60.8)	44 (13.2)	323 (96.7)	77 (23.1)	8 (2.4)
Tn4401a	261	261 (100)	261 (100)	259 (99.2)	235 (90.0)	160 (61.3)	17 (6.5)	250 (95.8)	72 (27.6)	4 (1.5)
Tn4401b	8	8 (100)	8 (100)	8 (100)	8 (100)	8 (100)	5 (62.5)	8 (100)	0 (0.0)	1 (12.5)
Tn4401c	24	24 (100)	24 (100)	24 (100)	22 (91.7)	12 (50.0)	2 (8.3)	24 (100)	4 (16.7)	1 (4.2)
ND	41	41 (100)	41 (100)	41 (100)	40 (97.6)	23 (56.1)	20 (48.8)	41 (100)	1 (2, 0.4)	2 (4.9)
KPC-3	6	6 (100)	6 (100)	5 (83.3)	6 (100)	2 (33.3)	0 (0.0)	6 (100)	0 (0.0)	0 (0.0)
Tn4401b	6	6 (100)	6 (100)	5 (83.3)	6 (100)	2 (33.3)	0 (0.0)	6 (100)	0 (0.0)	0 (0.0)
KPC-4	22	22 (100)	22 (100)	22 (100)	14 (63.6)	15 (68.2)	2 (9.1)	19 (86.4)	3 (13.6)	0 (0.0)
Tn4401b	22	22 (100)	22 (100)	22 (100)	14 (63.6)	15 (68.2)	2 (9.1)	19 (86.4)	3 (13.6)	0 (0.0)
Total	362	362 (100)	362 (100)	359 (99.2)	325 (89.8)	220 (60.8)	46 (12.7)	348 (96.1)	80 (22.1)	8 (2.2)

TABLE 3 | Co-producing drug CTX-M-type extended-spectrum beta-lactamase.

Total		CTX-M-type extended-spectrum beta-lactamase		
		Type 1	Type 9	Both types 1 and 9
KPC-2	334	156 (46.7)	21 (6.3)	6 (1.8)
Tn4401a	261	136 (52.1)	9 (3.4)	6 (1.8)
Tn4401b	8	2 (25.0)	1 (12.5)	0 (0.0)
Tn4401c	24	5 (20.8)	0 (0.0)	0 (0.0)
ND	41	13 (31.7)	11 (26.8)	0 (0.0)
KPC-3	6	0 (0.0)	0 (0.0)	0 (0.0)
Tn4401b	6	0 (0.0)	0 (0.0)	0 (0.0)
KPC-4	22	7 (31.8)	2 (9.1)	0 (0.0)
Tn4401b	22	7 (31.8)	2 (9.1)	0 (0.0)
Total	362	163 (45.0)	23 (6.4)	6 (1.7)

ST11/ND[*bla*_{KPC-2}] in 2014 triggered the inter-regional dissemination of KPC producers.

In agreement with the Healthcare Delivery System in South Korea, convalescent hospitals are in charge post-acute care and general hospitals are responsible for acute care, leading a habitual inter-hospital patient transfer. In general, the infection control system in convalescent hospitals is poorly operated compared to that in general hospitals and their role as a reservoir of CPE has been highly suspected. Indeed, the core KPC-producing subclone *K. pneumoniae* ST307/Tn4401a[*bla*_{KPC-2}] was identified in all three types of healthcare facilities: 153 of the 362 KPC producers from general hospitals, one of the three isolates from the two convalescent hospitals and one of the two isolates from clinics. It suggested that the KPC has disseminated during the inter-hospital patient transfer and expanded up to the community level. Further study with more isolates from the two types of healthcare facility is needed.

The well-known KPC-producing clone in a global level is *K. pneumoniae* ST258 (Pitout et al., 2015; Sheppard et al., 2016), however, it was not the case in South Korea. The foremost

South Korean KPC producer was *K. pneumoniae* ST307 clone and the ST11 was followed by presenting half the prevalence of ST307. Indeed, the ST258 (*gapA-infB-mdh-pgi-phoE-rpoB-tonB*, 3-3-1-1-1-1-79) is a deduced variant of ST11 (3-3-1-1-1-1-43) manufactured by multiple events of a genetic rearrangement (Breurec et al., 2013). *K. pneumoniae* ST307, which has gained attention as a potentially high-risk KPC producer, has been a notorious clone producing CTX-M-15 (Villa et al., 2017) and the clone co-producing KPC and CTX-M-15 was recently identified in Italy (Geraci et al., 2015). The South Korean ST307 clone are assumed to be close to the Italian clone in an evidence of 71.9% KPC-producing *K. pneumoniae* ST307 co-produced CTX-M enzyme. The second-most disseminated *K. pneumoniae* ST11 is one of the main KPC-producing clones in China, and it frequently co-harbors the *rmtB* genes (Cheng et al., 2016). The 22 RmtB-co-producers were all *K. pneumoniae* ST11 representing 28.7% (22/77) the clone.

KPC-2 and KPC-3 are the prevailing global subtypes of KPC, with regional variation (Munoz-Price et al., 2013). Among the South Korean KPC producers, KPC-4 was the second most prevalent subtype, although the number of KPC-4 producers was only one fifteenth that of the KPC-2 producers. Since the bacterial host species varied, the Tn4401b-associated plasmid (along with sporadic occurrence) was likely responsible for its observed dissemination. KPC-3, which was produced by fewer bacterial hosts, was limited to one district. KPC-2, KPC-3, and KPC-4 each confer a similar level of carbapenem resistance (Mehta et al., 2015). However, among the KPC producers studied, KPC-2 had the highest MIC₅₀s for carbapenems, while KPC-3 and KPC-4 had 3- and 4-dilution-reduced MIC₅₀s, respectively (Figure 2). It is likely that multiple factors (such as small relative group size, or low gene expression in the Tn4401b isotype) accounted for this unexpected result.

Although group size varied (334 vs. 17), the KPC-producing *K. pneumoniae* isolates had 3-dilution higher MIC₅₀s for carbapenems compared to the KPC-producing *E. coli* isolates (Figure 2), pronouncing the importance of bacterial host background for revealing resistance. As well, the Tn4401 isotypes, differing the promoter of the *bla*_{KPC} gene, is closely associated

with the level of gene expression resulting in varied carbapenem susceptibility (Naas et al., 2012; Cheruvanky et al., 2017). For example, Tn4401a and Tn4401d possessing two promoters upstream from the gene confer high carbapenem resistance whereas Tn4401c and Tn4401e, with one promoter, confer low levels of resistance and the Tn4401b, with three promoters, confers moderate resistance (Naas et al., 2012; Cheruvanky et al., 2017). Three isotypes of the transposon that are Tn4401a, Tn4401b, and Tn4401c were identified in our study. Tn4401a had the highest MIC₅₀s for carbapenems and the Tn4401b exhibited the lowest values. Tn4401c of similar MIC₅₀ values to those of Tn4401a could be explained by the bacterial host, as *K. pneumoniae* likely, somehow, has the most efficient KPC production. The relatively high meropenem MIC₅₀ by ND transposon encoding the *bla*_{KPC-2} compared to the other isotypes of Tn4401 needed to be further studied.

Though the global spread of KPC-producing *Enterobacteriaceae* is becoming more common, local epidemiology patterns need to be actively assessed. An awareness of the details of regional epidemiology is fundamental for understanding specific situations. Most endemic cases have occurred via the clonal dissemination of notorious KPC-producing clones, and the South Korean situation was no exception. The disseminated *K. pneumoniae* clones in South Korea have been linked to the Italian ST307 and Chinese ST11 strains, confirming the importance of international and intercontinental travel for clonal spread. There remains a paucity of antimicrobial options to treat pathogens capable of KPC production, which raises mortality rates. Only a few new drug candidates are in development at present. Combination therapy using both β -lactams and β -lactamase inhibitors that are stable in the presence of KPC enzymes (such as avibactam) is available

in other parts of the world, but is not yet accessible in South Korea. Because South Korea is facing endemic dissemination, the situation is critical. Only systematic surveillance will empower researchers to control and prevent the spread of KPC producers.

ETHICS STATEMENT

The research, which has no involvement of human subject but the clinical isolates, does meet the exempt category without approval from Ethics Committee on Human Research of the Health Ministry in South Korea and the study design has not been reviewed by the committee.

AUTHOR CONTRIBUTIONS

SJ and E-JY: contributed to idea conception, designed the study, conducted the experiments, interpreted the data, and wrote the manuscript; JK, DK, HL, JY, and KL: conducted the experiments and interpreted the data.

ACKNOWLEDGMENTS

This research was supported by a fund (KCDC2017E4400100#) from the Research of Korea Centers for Disease Control and Prevention.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2018.00056/full#supplementary-material>

REFERENCES

- Albiger, B., Glasner, C., Struelens, M. J., Grundmann, H., Monnet, D. L., and European Survey of Carbapenemase-Producing Enterobacteriaceae Working Group (2015). Carbapenemase-producing Enterobacteriaceae in Europe: assessment by national experts from 38 countries, May 2015. *Euro Surveill* 20. doi: 10.2807/1560-7917.ES.2015.20.45.30062
- Breurec, S., Guessennd, N., Timinouni, M., Le, T. A., Cao, V., Ngandjio, A., et al. (2013). *Klebsiella pneumoniae* resistant to third-generation cephalosporins in five African and two Vietnamese major towns: multiclonal population structure with two major international clonal groups, CG15 and CG258. *Clin. Microbiol. Infect.* 19, 349–355. doi: 10.1111/j.1469-0691.2012.03805.x
- Cheng, L., Cao, X. L., Zhang, Z. F., Ning, M. Z., Xu, X. J., Zhou, W., et al. (2016). Clonal dissemination of KPC-2 producing *Klebsiella pneumoniae* ST11 clone with high prevalence of *oqxAB* and *rmtB* in a tertiary hospital in China: results from a 3-year period. *Ann. Clin. Microbiol. Antimicrob* 15:1. doi: 10.1186/s12941-015-0109-x
- Cheruvanky, A., Stoesser, N., Sheppard, A. E., Crook, D. W., Hoffman, P. S., Weddle, E., et al. (2017). Enhanced *Klebsiella pneumoniae* carbapenemase expression from a novel Tn4401 deletion. *Antimicrob. Agents Chemother.* 61:e00025-17. doi: 10.1128/AAC.00025-17
- Clinical and Laboratory Standards Institute (2015a). *Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically*. (Wayne, PA: Clinical and Laboratory Standards Institute).
- Clinical and Laboratory Standards Institute (2015b). *Performance Standards for Antimicrobial Susceptibility Testing: Twenty-Fifth Informational Supplement*. (Wayne, PA: Clinical and Laboratory Standards Institute).
- Cuzon, G., Naas, T., and Nordmann, P. (2011). Functional characterization of Tn4401, a Tn3-based transposon involved in *bla*_{KPC} gene mobilization. *Antimicrob. Agents Chemother.* 55, 5370–5373. doi: 10.1128/AAC.05202-11
- Diancourt, L., Passet, V., Verhoef, J., Grimont, P. A., and Brisse, S. (2005). Multilocus sequence typing of *Klebsiella pneumoniae* nosocomial isolates. *J. Clin. Microbiol.* 43, 4178–4182. doi: 10.1128/JCM.43.8.4178-4182.2005
- Geraci, D. M., Bonura, C., Giuffrè, M., Saporito, L., Graziano, G., Aleo, A., et al. (2015). Is the monoclonal spread of the ST258, KPC-3-producing clone being replaced in southern Italy by the dissemination of multiple clones of carbapenem-nonsusceptible, KPC-3-producing *Klebsiella pneumoniae*? *Clin. Microbiol. Infect.* 21, e15–e17. doi: 10.1016/j.cmi.2014.08.022
- Hong, S. K., Yong, D., Kim, K., Hong, S. S., Hong, S. G., Khosbayan, T., et al. (2013). First outbreak of KPC-2-producing *Klebsiella pneumoniae* sequence type 258 in a hospital in South Korea. *J. Clin. Microbiol.* 51, 3877–3879. doi: 10.1128/JCM.01730-13
- Jeong, S. H., Kim, H. S., Kim, J. S., Shin, D. H., Kim, H. S., Park, M. J., et al. (2016). Prevalence and molecular characteristics of carbapenemase-producing Enterobacteriaceae from five hospitals in Korea. *Ann. Lab. Med.* 36, 529–535. doi: 10.3343/alm.2016.36.6.529
- Kim, J. O., Song, S. A., Yoon, E. J., Shin, J. H., Lee, H., Jeong, S. H., et al. (2017). Outbreak of KPC-2-producing Enterobacteriaceae caused by clonal dissemination of *Klebsiella pneumoniae* ST307 carrying an IncX3-type plasmid harboring a truncated Tn4401a. *Diagn. Microbiol. Infect. Dis.* 87, 343–348. doi: 10.1016/j.diagmicrobio.2016.12.012
- Mehta, S. C., Rice, K., and Palzkill, T. (2015). Natural variants of the KPC-2 Carbapenemase have evolved increased catalytic efficiency for ceftazidime

- hydrolysis at the cost of enzyme stability. *PLoS Pathog.* 11:e1004949. doi: 10.1371/journal.ppat.1004949
- Munoz-Price, L. S., Poirel, L., Bonomo, R. A., Schwaber, M. J., Daikos, G. L., Cormican, M., et al. (2013). Clinical epidemiology of the global expansion of *Klebsiella pneumoniae* carbapenemases. *Lancet Infect. Dis.* 13, 785–796. doi: 10.1016/S1473-3099(13)70190-7
- Naas, T., Cuzon, G., Truong, H. V., and Nordmann, P. (2012). Role of ISKpn7 and deletions in *bla*_{KPC} gene expression. *Antimicrob. Agents Chemother.* 56, 4753–4759. doi: 10.1128/AAC.00334-12
- Pérez-Pérez, F. J., and Hanson, N. D. (2002). Detection of plasmid-mediated AmpC β -lactamase genes in clinical isolates by using multiplex PCR. *J. Clin. Microbiol.* 40, 2153–2162. doi: 10.1128/JCM.40.6.2153-2162.2002
- Pitout, J. D., Nordmann, P., and Poirel, L. (2015). Carbapenemase-producing *Klebsiella pneumoniae*, a key pathogen set for global nosocomial dominance. *Antimicrob. Agents Chemother.* 59, 5873–5884. doi: 10.1128/AAC.01019-15
- Rhee, J. Y., Park, Y. K., Shin, J. Y., Choi, J. Y., Lee, M. Y., Peck, K. R., et al. (2010). KPC-producing extreme drug-resistant *Klebsiella pneumoniae* isolate from a patient with diabetes mellitus and chronic renal failure on hemodialysis in South Korea. *Antimicrob. Agents Chemother.* 54, 2278–2279. doi: 10.1128/AAC.00011-10
- Ryoo, N. H., Kim, E. C., Hong, S. G., Park, Y. J., Lee, K., Bae, I. K., et al. (2005). Dissemination of SHV-12 and CTX-M-type extended-spectrum beta-lactamases among clinical isolates of *Escherichia coli* and *Klebsiella pneumoniae* and emergence of GES-3 in Korea. *J. Antimicrob. Chemother.* 56, 698–702. doi: 10.1093/jac/dki324
- Sheppard, A. E., Stoesser, N., Wilson, D. J., Sebra, R., Kasarskis, A., Anson, L. W., et al. (2016). Nested Russian doll-like genetic mobility drives rapid dissemination of the carbapenem resistance gene *bla*_{KPC}. *Antimicrob. Agents Chemother.* 60, 3767–3778. doi: 10.1128/AAC.00464-16
- The joint CLSI–EUCAST and Polymyxin Breakpoints Working Group (2016). *Recommendations for MIC Determination of Colistin (polymyxin E) as Recommended by the Joint CLSI–EUCAST Polymyxin Breakpoints Working Group*. Available online at: http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/General_documents/Recommendations_for_MIC_determination_of_colistin_March_2016.pdf (Accessed March 22, 2016).
- Villa, L., Feudi, C., Fortini, D., Brisse, S., Passet, V., Bonura, C., et al. (2017). Diversity, virulence, and antimicrobial resistance of the KPC-producing *Klebsiella pneumoniae* ST307 clone. *Microb. Genom.* 3:e000110. doi: 10.1099/mgen.0.000110
- Wirth, T., Falush, D., Lan, R., Colles, F., Mensa, P., Wieler, L. H., et al. (2006). Sex and virulence in *Escherichia coli*: an evolutionary perspective. *Mol. Microbiol.* 60, 1136–1151. doi: 10.1111/j.1365-2958.2006.05172.x
- Yigit, H., Queenan, A. M., Anderson, G. J., Domenech-Sanchez, A., Biddle, J. W., Steward, C. D., et al. (2001). Novel carbapenem-hydrolyzing beta-lactamase, KPC-1, from a carbapenem-resistant strain of *Klebsiella pneumoniae*. *Antimicrob. Agents Chemother.* 45, 1151–1161. doi: 10.1128/AAC.45.4.1151-1161.2001
- Yoo, J. S., Kim, H. M., Yoo, J. I., Yang, J. W., Kim, H. S., Chung, G. T., et al. (2013). Detection of clonal KPC-2-producing *Klebsiella pneumoniae* ST258 in Korea during nationwide surveillance in 2011. *J. Med. Microbiol.* 62, 1338–1342. doi: 10.1099/jmm.0.059428-0

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Yoon, Kim, Kim, Lee, Yang, Lee and Jeong. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.