

# Management of postoperative ileus after robot-assisted laparoscopic prostatectomy

Jee Soo Park, MD<sup>a</sup>, Jongchan Kim, MD<sup>a</sup>, Won Sik Jang, MD<sup>a</sup>, Ji Eun Heo, MD<sup>a</sup>, Ahmed Elghiaty, MD<sup>a,b</sup>, Koon Ho Rha, MD, PhD<sup>a</sup>, Young Deuk Choi, MD, PhD<sup>a</sup>, Won Sik Ham, MD, PhD<sup>a,\*</sup>

## Abstract

To compare different postoperative management methods on the recovery of bowel function after robot-assisted laparoscopic prostatectomy (RALP).

This is a prospective study of 716 patients who underwent RALP at Severance Hospital in Seoul, South Korea, between March 2017 and February 2018. Instructions for the different postoperative management methods (mobilization, abdominal massage, hot pack therapy, and gum chewing) were presented to patients, who subsequently reported when these activities were performed as well as the time to first flatus on a designated form.

There were no significant differences in age, height, weight, body mass index, body surface area, prevalence of hypertension and diabetes mellitus, and in American Society of Anesthesiologists (ASA) scores with respect to early bowel recovery. Prolonged times of surgery and anesthesia significantly caused delays in bowel recovery. The total number and time of mobilization, total time of hot pack therapy, and number of gum chewing were significantly and positively associated with bowel recovery. A Kaplan–Meier analysis showed that all of the postoperative management methods were positively associated with the mean time to first flatus.

Methods of postoperative management (mobilization, abdominal massage, hot pack therapy, and gum chewing) have positive effect on bowel motility after RALP. Furthermore, reductions in the times of surgery and anesthesia could significantly decrease prolonged delays in bowel recovery.

**Abbreviations:** ASA = American Society of Anesthesiologists, BMI = body mass index, BSA = body surface area, EBL = estimated blood loss, RALP = robot-assisted laparoscopic prostatectomy.

**Keywords:** management, postoperative ileus, robot-assisted laparoscopic prostatectomy

## 1. Introduction

Livingston and Passaro<sup>[1]</sup> define postoperative ileus as an “uncomplicated ileus occurring following surgery, resolving spontaneously within 2 to 3 days.” In postoperative ileus, the stomach recovers within 24 to 48 hours, whereas the motility of the colon returns after 48 to 72 hours.<sup>[1]</sup> Postoperative ileus remains the most common minor postoperative complication to

cause morbidity and delays in patient discharge from the hospital, leading to an increased economic burden on the healthcare system.<sup>[2]</sup> Therefore, many researchers have focused on the management of postoperative ileus: Many studies have investigated treatment approaches such as motility agents, early feeding, gum chewing, nasogastric intubation, fluid restriction, epidural anesthesia and analgesia, mobilization, and physical therapy.<sup>[2–7]</sup> However, these therapies have not been routinely used in the clinic because of their limited clinical efficacy.<sup>[2]</sup>

Most studies have focused on colorectal surgery; however, only a few studies have examined the postoperative management of a major urological surgery.<sup>[3,8,9]</sup> Robot-assisted surgery can better minimize blood loss and morbidity than traditional open surgical techniques.<sup>[10]</sup> No studies have determined the efficacy of postoperative ileus management after robot-assisted laparoscopic prostatectomy (RALP). The purpose of this study is to compare the effects of different postoperative management methods (mobilization, abdominal massage, hot pack therapy, and gum chewing) on the reduction of postoperative ileus and improved bowel recovery after RALP.

## 2. Materials and methods

### 2.1. Study population

This study is a prospective study of patients who underwent RALP at Severance Hospital in Seoul, South Korea, between March 2017 and February 2018. This study was approved by the Institutional Review Board of the Yonsei University Health System (project no: 4-2017-0225). We used the technique described by Menon et al.<sup>[11,12]</sup> The routine postoperative

Editor: Johannes Mayr.

Ethical approval: All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee, and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent was obtained after a full explanation of the study. This study did not involve patient intervention or the use of human tissue samples.

The authors have no conflicts of interest to disclose.

<sup>a</sup> Department of Urology and Urological Science Institute, Yonsei University College of Medicine, Seoul, Republic of Korea, <sup>b</sup> Department of Urology, Tanta University College of Medicine, Tanta, Egypt.


\* Correspondence: Won Sik Ham, Department of Urology and Urological Science Institute, Yonsei University College of Medicine, Seoul 03722, Republic of Korea (e-mail: uroham@yuhs.ac).

Copyright © 2018 the Author(s). Published by Wolters Kluwer Health, Inc. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CC BY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Medicine (2018) 97:44(e13036)

Received: 2 August 2018 / Accepted: 8 October 2018

<http://dx.doi.org/10.1097/MD.00000000000013036>



**Figure 1.** Flow chart of excluding patients for the patients who underwent robot-assisted laparoscopic prostatectomy (RALP).

analgesia was commenced with intravenous pethidine (meperidine) usually by patient-controlled methods (PCA) and oral acetaminophen were used for additional analgesia as required. We removed the transurethral catheter after 2 weeks after the surgery. Patients were excluded for the following reasons: history of inflammatory bowel disease, prior abdominal radiation, or surgery, and concurrent surgery during RALP (Fig. 1). Seven hundred sixteen patients were enrolled in the study. The details of the study were explained to the subjects, and written informed consent was obtained.

## 2.2. Measurement and classification of variables

The time at the end of surgery was defined as the zero hour. To monitor the recovery of bowel function, all the patients were instructed to write down when a bowel-related event occurred. Instructions for mobilization, abdominal massage, hot pack therapy, and gum chewing were presented to all patients, who were instructed to document the exact time they performed those activities. These management activities were discontinued when the passage of flatus first occurred. The end of postoperative ileus was defined as the passage of flatus in this study, and the study endpoint was the time to the first postoperative passage of flatus.

The time from the first incision to the final closure of the wound was defined as the time of surgery. Demographic data, including patient age, height, weight, body mass index (BMI), body surface area (BSA), American Society of Anesthesiologists (ASA) score, and history of hypertension and diabetes mellitus, were recorded. Intraoperative parameters including time of surgery, time of anesthesia, and estimated blood loss (EBL) were analyzed.

The group with the first postoperative passage of flatus within 24 hours, which is the mean time for the time to first flatus, was classified as the early bowel recovery group, whereas those who had the first postoperative passage of flatus after 24 hours were considered the late bowel recovery group. Patients who performed mobilization, abdominal massage, hot pack therapy, and gum chewing more than once were considered the group who

performed that management method and were compared with the control group, which was defined as those who did not perform this management method. Mobilization was further divided into groups who walked more than 30 and 60 minutes per day, respectively.

## 2.3. Statistical analysis

The results are reported as the mean (standard deviation) for continuous variables and as a percentage for categorical variables. For the univariate analysis, the *t* test was used to compare continuous variables. The multivariate analysis used multivariate models of logistic regression, including all risk factors that were significantly associated in the univariate analysis. The times to the different events were compared among groups using Kaplan–Meier analyses and the log-rank test. SPSS software, version 23.0 (IBM Corp, Armonk, NY), was used for statistical analyses. All statistical tests were 2-tailed, and a *P* value less than .05 was considered statistically significant.

## 3. Results

The baseline characteristics of the study population are shown in Table 1. The mean age of the patients who participated in this study was  $65.9 \pm 4.5$  years. The prevalence of hypertension and diabetes mellitus were 39.5% and 14.4%, respectively. Most patients were in the ASA score 2 (46.4%) and score 3 (37.6%) groups. The mean time of surgery was  $142.6 \pm 70.3$  minutes. The mean time to the first postoperative passage of flatus was  $29.1 \pm 16.2$  hours.

Five hundred fifty-six (77.7%) patients have undergone lymphadenectomy and the mean number of lymph nodes removed were  $11.9 \pm 8.3$ . Most of the patients (70.7%) had nerve sparing surgery on both the sides. One hundred fifty (20.9%) patients had nerve sparing surgery on 1 side and 60 (8.4%) patients did not have nerve sparing surgery. Performance of lymphadenectomy and nerve sparing surgery did not differ

**Table 1**  
Characteristics of the study population.

	Study population (n=716)
Patient characteristics	
Age, y	65.9 (4.5)
Height, cm	168.3 (5.6)
Weight, kg	69.3 (8.1)
BMI, kg/m <sup>2</sup>	24.4 (2.3)
BSA, m <sup>2</sup>	1.8 (0.1)
ASA score	
1	115 (16.1%)
2	332 (46.4%)
3	269 (37.6%)
HTN, n, %	283 (39.5%)
DM, n, %	103 (14.4%)
Time to first flatus (hours)	29.1 (16.2)
Intraoperative parameters	
Time of surgery, min	142.6 (70.3)
Time of anesthesia, min	182.8 (66.4)
EBL, cc	421.4 (347.2)
Postoperative management	
Mobilization	
Number	2.6 (2.2)
Total time, min	47.3 (43.8)
Abdominal massage	
Number	0.2 (0.5)
Total time, min	3.0 (9.1)
Hot pack therapy	
Number	0.8 (1.2)
Total time, min	14.7 (25.1)
Gum chewing	
Number	0.8 (1.2)
Total time, min	25.4 (45.8)

Data are shown as the mean (SD) or number of subjects (%).

ASA = American Society of Anesthesiologists, BMI = body mass index, BSA = body surface area, DM = diabetes mellitus, EBL = estimated blood loss, HTN = hypertension.

between early and late bowel recovery group ( $P=.120$  and  $P=.473$ , respectively).

There were no significant differences in age, height, weight, BMI, and BSA between early and late bowel recovery groups, although, as expected, those in the late bowel recovery group were older (Table 2). There were no differences in the prevalence of hypertension and diabetes mellitus between groups. ASA scores were not significantly different between the 2 groups (Table 2). Times of surgery and anesthesia were significantly longer in the late bowel recovery group in the multivariate analysis. EBL was greater in the late bowel recovery group but was not statistically significant in multivariate analysis. The total time of mobilization, abdominal massage, hot pack therapy were significantly associated with bowel recovery in the univariate analysis, and mobilization and hot pack therapy showed significant association in multivariate analyses. The number of gum chewing was significantly associated with improved bowel recovery. The Gleason score and pathological stage of the study population are shown in Table 3. Only the pathological T stage was significantly associated with bowel recovery both in the univariate and multivariate analysis ( $P=.019$ ).

The mean time to first flatus was significantly decreased ( $P<.001$ ) in patients in the mobilization group [38.2 (0.6) hours,  $n=668$ ] compared with the control group [56.3 (1.4) hours,  $n=48$ ], as shown by the Kaplan-Meier curves in Figure 2A. For the group who walked less than 30 minutes, the mean time to first

flatus was significantly prolonged compared with the control group [36.6 (0.7),  $n=514$  vs. 46.8 (0.9) hours,  $n=202$ ,  $P<.001$ ] (Fig. 2B). Similarly, there was significant difference between the group who walked more than 60 minutes and the control group [27.1 (0.2),  $n=228$  vs. 42.4 (0.6) hours,  $n=488$ ,  $P<.001$ ] (Fig. 2C). There were significant differences between the group who performed hot pack therapy or abdominal massage and the control group [37.0 (0.9),  $n=320$  vs. 42.0 (0.8) hours,  $n=396$ ,  $P<.001$ ; 34.3 (1.3),  $n=108$  vs. 41.3 (0.7) hours,  $n=608$ ,  $P<.001$ ] (Fig. 2D and E). Furthermore, the mean time to first flatus was significantly prolonged ( $P<.001$ ) in patients who performed gum chewing [36.2 (1.0) hours,  $n=294$ ] compared with the control group [42.8 (0.8) hours,  $n=422$ ] (Fig. 2F).

#### 4. Discussion

This is the first study to assess the effects of the different management methods (mobilization, abdominal massage, hot pack therapy, and gum chewing) on reducing postoperative ileus after RALP. Many studies have proven treatment efficacy in the management of postoperative ileus,<sup>[2–7]</sup> but surgeons are unlikely to suggest such treatments to patients. Postoperative ileus causes patient discomfort and pain. Moreover, it can prolong hospitalization, which leads to increases in hospital costs.<sup>[13,14]</sup> No studies have evaluated the effects of mobilization, abdominal massage, hot pack therapy, and gum chewing on reductions in postoperative ileus after RALP.

Similar to the results of most studies, mobilization was positively associated with bowel recovery. Kaplan-Meier analyses also showed that the mobilization group had a significantly decreased time to first flatus compared with the control group. The group with more time of mobilization had the faster time for bowel recovery, as shown by the Kaplan-Meier analysis (Fig. 2A–C). Some studies reported that physical exercise does not improve colonic mobility.<sup>[15,16]</sup> However, this study showed significantly positive relationship between mobilization and bowel recovery time both in multivariate and Kaplan-Meier analysis. There are studies that reported the prolonged immobilization after surgery causing postoperative complications and delayed recovery.<sup>[17–19]</sup>

No studies have demonstrated the effects of hot pack therapy on the management of postoperative ileus. However, some studies reported that hot pack therapy increases regional blood flow.<sup>[20,21]</sup> Increases in blood flow to the gastrointestinal tract would result in early recovery of bowel function. Our results showed that the mean time to first flatus in patients who utilized hot pack therapy was significantly decreased. Takayama et al<sup>[21]</sup> showed that thermal stimulation of the para-umbilical region increased blood flow to the superior mesenteric artery within 20 minutes of thermal stimulation. This finding explains why patients utilizing hot pack therapy passed gas early, as shown by the Kaplan-Meier curve in Figure 2D.

Gum chewing is considered a quick and safe method to enhance the recovery of bowel function after surgery with medium-level evidence.<sup>[22,23]</sup> Chewing increases the levels of several hormones, such as gastrin, neuropeptides, pancreatic polypeptide, and cholecystokinin, which are all associated with indirect vagal afferent stimulation.<sup>[24]</sup> Moreover, chewing increases salivary and gastric secretions.<sup>[3]</sup> Asao et al<sup>[24]</sup> first reported the use of chewing gum after surgery in 2002. These authors described gum chewing as a form of sham feeding, which stimulates the motility of the gastrointestinal tract and results in reductions in the times to first flatus and first bowel

**Table 2**

Characteristics of the study population according to the first postoperative passage of flatus in 24 h.

	First postoperative passage of flatus		<i>P</i> value*	<i>P</i> value†
	Before 24 hours (n = 315)	After 24 hours (n = 401)		
Patient characteristics				
Age, y	65.6 (3.9)	66.0 (5.0)	.202	
Height, cm	168.0 (5.6)	168.6 (5.6)	.174	
Weight, kg	69.2 (8.1)	69.3 (8.1)	.907	
BMI, kg/m <sup>2</sup>	24.5 (2.3)	24.4 (2.3)	.407	
BSA, m <sup>2</sup>	1.8 (0.1)	1.8 (0.1)	.705	
ASA score				
1	50 (15.9%)	65 (16.2%)		
2	151 (47.9%)	181 (45.1%)	.743	
3	114 (36.2%)	155 (38.7%)		
HTN, n (%)	121 (38.4%)	162 (40.4%)	.589	
DM, n (%)	38 (12.1%)	65 (16.2%)	.117	
Intraoperative parameters				
Time of surgery, min	103.1 (23.3)	173.6 (78.9)	<.001	<.001
Time of anesthesia, min	147.8 (24.6)	210.3 (75.4)	<.001	.009
EBL, cc	322.5 (211.9)	499.1 (408.0)	<.001	.605
Postoperative management				
Mobilization				
Number	2.8 (2.7)	2.4 (1.8)	.005	.002
Total time, min	65.1 (56.1)	33.3 (22.7)	<.001	<.001
Abdominal massage				
Number	0.3 (0.6)	0.2 (0.5)	.104	.382
Total time, min	4.9 (12.4)	1.5 (4.9)	<.001	.627
Hot pack therapy				
Number	1.0 (1.2)	0.6 (1.2)	<.001	.014
Total time, min	20.4 (27.0)	10.2 (22.4)	<.001	
Gum chewing				
Number	0.9 (1.3)	0.7 (1.1)	.041	<.001
Total time, min	29.1 (50.1)	22.6 (42.0)	.065	

Data are shown as the mean (SD) or number of subjects (%).

ASA = American Society of Anesthesiologists, BMI = body mass index, BSA = body surface area, DM = diabetes mellitus, EBL = estimated blood loss, HTN = hypertension.


\* *P* value calculated using the *t* test (continuous data) and  $\chi^2$  test (categorical data).† *P* value calculated using logistic regression for multivariate analysis.**Table 3**

Gleason score and pathological stage of the study population according to the first postoperative passage of flatus in 24 h.

Study population (n = 716)	First postoperative passage of flatus		<i>P</i> value*	<i>P</i> value†
	Before 24 h (n = 315)	After 24 h (n = 401)		
Biopsy Gleason score				
Mean (SD)	7.2 (1.0)	7.3 (1.0)	.270	
≤6	189 (26.4%)	72 (22.9%)	.386	
3+4	164 (22.9%)	79 (25.1%)		
4+3	120 (16.8%)	55 (17.5%)		
8	154 (21.5%)	68 (21.6%)		
9-10	89 (12.4%)	41 (13.0%)		
pT stage				
T2	365 (51.0%)	143 (45.4%)		.019
T3a	223 (31.1%)	117 (37.1%)		.019
T3b	110 (15.4%)	47 (14.9%)		
T4	18 (2.5%)	8 (2.5%)		
pN stage				
N0	672 (93.9%)	300 (95.2%)		.172
N1	44 (6.1%)	15 (4.8%)		
pM stage				
M0	696 (97.2%)	304 (96.5%)		.315
M1	20 (2.8%)	11 (3.5%)		

Data are shown as the mean (SD) or number of subjects (%).

\* *P* value calculated using the *t* test (continuous data) and  $\chi^2$  test (categorical data).† *P* value calculated using logistic regression for multivariate analysis.



**Figure 2.** Kaplan–Meier curves of the first passage of flatus postoperatively between the control group and each group performing a type of postoperative management. A, Mobilization. B, Walking more than 30min. C, Walking more than 60min. D, Hot pack therapy. E, Abdominal massage. F, Gum chewing.

movement.<sup>[24]</sup> Our study also showed that gum chewing enhanced the bowel function recovery.

The mean time of abdominal massage was less than 3 minutes, which may be related to the reluctance of patients to perform abdominal massage due to surgical pain. However, our results showed that abdominal massage also supports bowel recovery (Fig. 2E). Prolonged times of surgery and anesthesia resulted in delayed bowel recovery. It is well known that all drugs used for the induction and maintenance of general anesthesia depress bowel motility.<sup>[25]</sup>

There are several limitations of this study. First, flatus may not be an ideal end point, since it requires a conscious patient who must report its occurrence to investigators. We could not measure times to first bowel movement, the most reliable end point. Many patients have bowel movements almost immediately after the passage of flatus. Moreover, most patients were discharged before having a bowel movement. Second, this study is based on patients' self-reported questionnaires. As patients report the number and duration of mobilization, abdominal massage, hot pack therapy, and gum chewing after surgery, there could be some misreported and/or missing information.

This study showed that the different postoperative management methods (mobilization, abdominal massage, hot pack

therapy, and gum chewing) that were recommended to patients to help bowel recovery have positive effects on the recovery of bowel motility after RALP. Prolonged times of surgery and anesthesia resulted in delayed bowel recovery. Surgeons should thus focus on reducing the time of surgery to help patients recover bowel function. Moreover, all postoperative management methods have been found to be effective in reducing bowel recovery time, especially, mobilization has been significantly found to be helpful.

## Author contributions

- Conceptualization: Jee Soo Park, Won Sik Ham.
- Data curation: Jee Soo Park, Jongchan Kim, Ji Eun Heo, Ahmed Elghiaty.
- Formal analysis: Jee Soo Park, Won Sik Jang.
- Investigation: Jee Soo Park.
- Methodology: Jee Soo Park.
- Project administration: Koon Ho Rha, Young Deuk Choi, Won Sik Ham.
- Supervision: Koon Ho Rha, Young Deuk Choi, Won Sik Ham.
- Writing – original draft: Jee Soo Park.
- Writing – review & editing: Jee Soo Park, Koon Ho Rha, Young Deuk Choi, Won Sik Ham.

## References

- [1] Livingston EH, Passaro EP. Postoperative ileus. *Dig Dis Sci* 1990;35:121–32.
- [2] Matros E, Rocha F, Zinner M, et al. Does gum chewing ameliorate postoperative ileus? Results of a prospective, randomized, placebo-controlled trial. *J Am Coll Surg* 2006;202:773–8.
- [3] Choi H, Kang SH, Yoon DK, et al. Chewing gum has a stimulatory effect on bowel motility in patients after open or robotic radical cystectomy for bladder cancer: a prospective randomized comparative study. *Urology* 2011;77:884–90.
- [4] Asgeirsson T, El-Badawi KI, Mahmood A, et al. Postoperative ileus: it costs more than you expect. *J Am Coll Surg* 2010;210:228–31.
- [5] Noble EJ, Harris R, Hosie KB, et al. Gum chewing reduces postoperative ileus? A systematic review and meta-analysis. *Int J Surg* 2009;7:100–5.
- [6] Bisanz A, Palmer JL, Reddy S, et al. Characterizing postoperative paralytic ileus as evidence for future research and clinical practice. *Gastroenterol Nurs* 2007;31:336–44.
- [7] Holte K, Kehlet H. Postoperative ileus: a preventable event. *Br J Surg* 2000;87:1480–93.
- [8] Kouba EJ, Wallen EM, Pruthi RS, et al. Gum chewing stimulates bowel motility in patients undergoing radical cystectomy with urinary diversion. *Urology* 2007;70:1053–6.
- [9] Maffezzini M, Campodonico F, Parodi D, et al. Current perioperative management of radical cystectomy with intestinal urinary reconstruction for muscle-invasive bladder cancer and reduction of the incidence of postoperative ileus. *Surg Oncol* 2008;17:41–8.
- [10] Guru KA, Wilding GE, Piacente P, et al. Robot-assisted radical cystectomy vs. open radical cystectomy: assessment of postoperative pain. *Can J Urol* 2007;14:3753–6.
- [11] Menon M, Hemal AK. Vattikuti Institute prostatectomy: a technique of robotic radical prostatectomy: experience in more than 1000 cases. *J Endourol* 2004;18:611–9.
- [12] Menon M, Tewari A, Peabody J. Vattikuti Institute prostatectomy: technique. *J Urol* 2003;169:2289–92.
- [13] Senagore A. Pathogenesis and clinical and economic consequences of postoperative ileus. *Clin Exp Gastroenterol* 2010;3:87–9.
- [14] Saclarides TJ. Current choices good or bad for the proactive management of postoperative ileus: a surgeon's view. *J Perianesthesia Nurs* 2006;21:7–15.
- [15] Rao SS, Beaty J, Chamberlain M. Effects of acute graded exercise on human colonic motility. *Am J Physiol* 1999;276:1221–6.
- [16] Waldhausen JH, Schirmer BD. The effect of ambulation on recovery from postoperative ileus. *Ann Surg* 1990;212:671–7.
- [17] Allen C, Glasziou P, Del Mar C. Bed rest: a potentially harmful treatment needing more careful evaluation. *Lancet* 1999;354:1229–33.
- [18] Harper CM, Lyles YM. Physiology and complications of bed rest. *J Am Geriatr Soc* 1988;36:1047–54.
- [19] Kehlet H. Multimodal approach to control postoperative pathophysiology and rehabilitation. *Br J Anaesth* 1997;78:606–17.
- [20] Okada K, Yamaguchi T, Minowa K, et al. The influence of hot pack therapy on the blood flow in masseter muscles. *J Oral Rehabil* 2005;32:480–6.
- [21] Takayama S, Seki T, Watanabe M, et al. Changes of blood flow volume in the superior mesenteric artery and brachial artery with abdominal thermal stimulation. *Evid Based Complement Alternat Med* 2011;2011:214089.
- [22] Gustafsson UO, Scott MJ, Schwenk W, et al. Guidelines for perioperative care in elective colonic surgery: enhanced recovery after surgery (ERAS<sup>®</sup>) society recommendations. *World J Surg* 2013;37:259–84.
- [23] Nygren J, Thacker J, Carli F, et al. Guidelines for perioperative care in elective rectal/pelvic surgery: enhanced recovery after surgery (ERAS<sup>®</sup>) society recommendations. *World J Surg* 2013;37:285–305.
- [24] Asao T, Kuwano H, Nakamura J, et al. Gum chewing enhances early recovery from postoperative ileus after laparoscopic colectomy. *J Am Coll Surg* 2002;195:30–2.
- [25] Ogilvy AJ, Smith G. The gastrointestinal tract after anaesthesia. *Eur J Anaesthesiol Suppl* 1995;10:35–42.