

Predictive factors for the development of renal insufficiency following partial nephrectomy and subsequent renal function recovery

A multicenter retrospective study

Kwang Suk Lee, MD, MMS^a, Dae Keun Kim, MD, MMS^b, Kwang Hyun Kim, MD, PhD^c, Woo Jin Bang, MD, PhD^d, Hyung Joon Kim, MD, PhD^e, Sung Yul Park, MD, PhD^f, Koon Ho Rha, MD, PhD^g, Byung Ha Chung, MD, PhD^a, Jin Seon Cho, MD, PhD^d, Kyo Chul Koo, MD, PhD^{a,*}

Abstract

Patients who undergo partial nephrectomy (PN) may exhibit renal function insufficiency, and a subset of these patients achieves renal function recovery. We evaluated the predictors of renal insufficiency and subsequent renal function recovery following PN. Data on 393 patients who underwent PN for solid renal tumors between March 2001 and November 2013, obtained from 6 institutions, were retrospectively reviewed. Renal insufficiency was defined as new onset of chronic kidney disease stage ≥ 3 postoperatively on the second of 2 consecutive tests. Renal function recovery was defined as an estimated glomerular filtration rate ≥ 60 ml/minute/1.73 m² following renal insufficiency. Tumor complexity was stratified according to the RENAL classification system. The median (interquartile range) age, tumor size, and follow-up period were 53 (45–63) years, 2.6 (1.9–3.8) cm, and 36 (12–48) months, respectively. Tumors were of low complexity in 258/393 (65.6%) of cases. Renal insufficiency developed in 54/393 (13.5%) patients, in which age ≥ 60 years and preoperative creatinine ≥ 1.1 mg/ml were independent predictors. Tumor complexity, clamp type, and operative method were not significant prognostic factors. Among patients with newly developed renal insufficiency, 18/54 (33.3%) patients exhibited renal function recovery within a median period of 18 months, of which preoperative creatinine < 1.1 mg/ml was an independent predictor. Age ≥ 60 years and preoperative creatinine ≥ 1.1 mg/ml were risk factors for renal insufficiency following PN. Patients with renal insufficiency whose preoperative creatinine was < 1.1 mg/ml were likely to have renal function recovery.

Abbreviations: CT = computed tomography, EBL = estimated blood loss, eGFR = estimated glomerular filtration rate, HR = hazard ratio, MRI = magnetic resonance imaging, PN = partial nephrectomy, PSM = positive surgical margin, RCC = renal cell carcinoma, WIT = warm ischemia time.

Keywords: carcinoma, complications, partial nephrectomy, renal cell, warm ischemia time

Editor: Giuseppe Lucarelli.

All procedures involving human participants were performed in accordance with the ethical standards of the institutional research committee and with the 1964 Declaration of Helsinki and subsequent amendments or comparable ethical standards.

Informed consent was obtained from all individual participants included in the study.

All the authors declare that there is no conflict of interest.

^a Department of Urology, Yonsei University College of Medicine, Gangnam Severance Hospital, ^b Department of Urology, CHA Seoul Station Medical Center, CHA University, ^c Department of Urology, Ewha Women's University College of Medicine, ^d Department of Urology, Hallym University College of Medicine, Chuncheon, ^e Department of Urology, Konyang University College of Medicine, Daejeon, ^f Department of Urology, Hanyang University College of Medicine, ^g Department of Urology, Yonsei University College of Medicine, Shinchon Severance Hospital, Seoul, Korea.

* Correspondence: Kyo Chul Koo, Department of Urology, Yonsei University College of Medicine, 211 Eonjuro, Gangnam-gu, Seoul 135-720, Korea (e-mail: gckoo@yuhs.ac).

Copyright © 2019 the Author(s). Published by Wolters Kluwer Health, Inc. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial License 4.0 (CCBY-NC), where it is permissible to download, share, remix, transform, and buildup the work provided it is properly cited. The work cannot be used commercially without permission from the journal.

Medicine (2019) 98:18(e15516)

Received: 23 October 2018 / Received in final form: 10 April 2019 / Accepted: 11 April 2019

<http://dx.doi.org/10.1097/MD.00000000000015516>

1. Introduction

The National Comprehensive Cancer Network guidelines recommend partial nephrectomy (PN) as the treatment of choice for cT1 tumors.^[1] PN was originally indicated if radical nephrectomy would render the patient functionally anephric, necessitating dialysis, such as for renal cell carcinoma (RCC) in a solitary kidney, RCC in 1 kidney with inadequate contralateral renal function, or bilateral synchronous RCC. There is accumulating evidence that the oncologic outcomes of PN are comparable to those of radical nephrectomy.^[2,3] Moreover, PN allows preservation of renal function, decrease in overall mortality, and reduced risk of cardiovascular events.^[4,5] Hence, over the past decade, PN has become the standard of care for most technically resectable renal tumors.^[6] In Korea, the proportion of PN performed has rapidly increased from 2008 to 2014.^[7]

As a surrogate for the success of the PN, the trifecta outcome was investigated, which represents no complications, negative surgical margins, and warm ischemia time (WIT) < 25 minutes.^[8] In addition to the trifecta outcome, long-term renal function was another parameter used to prove the success of PN as one of the pentafecta outcomes.^[9] Several factors, including WIT, loss of parenchymal volume, and ischemic damage in preserved tissue during the operation, are known renal function risk factors after PN. These factors are closely related to various

surgical factors such as clamping methods, surgical methods, and renorrhaphy. However, published data from a multicenter database of patients with an initial diagnosis of RCC after PN is scarce. Moreover, to the best of our knowledge, no study has reported predictors of renal function recovery following renal insufficiency in patients who underwent PN. The aims of this multicenter analysis were to predict renal insufficiency and to identify the prognostic factors of subsequent renal function recovery following PN.

2. Materials and methods

2.1. Study population and data collection

A total of 393 consecutive Korean patients with RCC treated with PN were selected from the multi-center, Severance Urological Oncology Group PN registry. Data on patients who underwent PN for RCC between March 2001 and November 2013, obtained from 6 institutions, were retrospectively reviewed. The present study's retrospective protocol was reviewed and approved by the Yonsei University Health System Ethics Committee, which waived the requirement for informed consent (2019-005-001). The patient records were anonymized and de-identified prior to analysis. Patient demographics, including age, sex, body mass index, and a history of hypertension or diabetes mellitus, were collected. Perioperative and postoperative outcomes were assessed, including American Society of Anesthesiologists score, tumor size, RENAL nephrometry score, intraoperative surgical complications, operative time, clamp type, ischemic time, estimated blood loss (EBL), and serum chemistry.

Preoperative radiological examination data (computed tomography [CT] or magnetic resonance imaging [MRI]) of the enrolled patients were interpreted by radiologists in the urology department at each participating hospital. The final pathology was determined using surgical specimens and reported by the uropathologists at each institution. Pathological data included pathological tumor size, TNM stage, Fuhrman grade, positive surgical margin (PSM), and histological subtype.

Tumor complexity was stratified according to the RENAL classification system^[10]: low (RENAL nephrometry score ≤ 7) or high (RENAL nephrometry score ≥ 8). Postoperative complications (≤ 30 days after surgery) were graded according to the modified Clavien-Dindo classification.^[11] Trifecta achievement was defined as a WIT of < 25 minutes, negative surgical margins, and no complications intraoperatively or postoperatively (Clavien-Dindo complication grade ≥ 3) as a surrogate of surgical quality. The estimated glomerular filtration rate (eGFR) was determined using the Modification of Diet in Renal Disease formula.^[12] Renal insufficiency was defined as a new onset eGFR < 60 ml/minute/1.73 m² (chronic kidney disease stage ≥ 3) postoperatively on the second of 2 consecutive tests at least 3 months apart. Renal function recovery was defined as eGFR ≥ 60 ml/minute/1.73 m² following renal insufficiency.

Patients without notable complications were followed up with serum chemistries every 3 months for 2 years. Thereafter, the decision of follow up protocols after surgical treatment was based on the surgeons' discretion. All patients received standard care according to contemporary guidelines for the duration of follow-up.

2.2. Study endpoints

The endpoint was the predictors associated with renal insufficiency and subsequent renal function recovery.

2.3. Statistical analysis

Categorical variables were evaluated by the Fisher exact test. Differences in variables with a continuous distribution across categories were assessed using the Mann-Whitney *U* test. Multivariate Cox regression analyses were performed on predictors of renal insufficiency and subsequent renal function recovery that had a *P* value $< .05$ in the univariate analyses. The signed-rank test for each group was used to compare median serum creatinine and eGFR at different time points. All reported *P* values are two-sided, and statistical significance was set at *P* $< .05$. Statistical analyses were performed using the Statistical Package for Social Sciences, version 23.0, for Windows (IBM Corp., Armonk, NY, USA).

3. Results

The baseline clinical and demographic characteristics of the patients are shown in Table 1. The median (interquartile range) age, tumor size, and follow-up period were 53 (45–63) years, 2.6 (1.9–3.8) cm, and 36 (12–48) months, respectively. Tumors of low complexity were found in 258/393 (65.6%) cases. Clavien-Dindo complication grade ≥ 3 occurred in 5.1% (20/393) of patients. The main reasons for Clavien-Dindo complication grade ≥ 3 were hemorrhage (8/20, 40%) followed by pseudoaneurysm (5/20, 25%). Radical nephrectomy conversion occurred in 2/20 (0.5%) cases. The median ischemic time was 25.0 (18.0–31.0) minutes, and there were no differences according to tumor complexity. The achievement rates of trifecta and pentafecta were 43.8% and 37.9%, respectively.

Patients with high complexity had a larger tumor size (*P* = .002) and higher Fuhrman grade (≥ 3) (*P* = .011) than those with low complexity. Tumor complexity was significantly associated with complications (*P* = .005), ischemic time (*P* = .005), and PSM (*P* = .014), but not with postoperative renal insufficiency (*P* = .943). The achievement rate of trifecta for high and low complexities was 31.1% and 50.4%, respectively (*P* $< .001$), and the achievement rate of pentafecta was 28.1%, 43.0%, respectively (*P* = .004). There were no differences in operative time, type of pedicle clamp, EBL, proportion of renal insufficiency, or follow-up period between the groups according to renal complexity.

Table 2 shows the change in median eGFR over time according to clamping type, RENAL nephrometry score, and WIT. Decreased eGFR at 3 months postoperatively finally recovered to near preoperative levels at 2 years (*P* = .865). Although the eGFR level at 2 years recovered to the preoperative level regardless of clamping method, the patterns of the change in eGFR varied. The patients with WIT ≥ 25 minutes had no recovery of renal function at 2 years compared to preoperative levels.

Renal insufficiency developed in 54/393 (13.7%) patients, in which age ≥ 60 years (hazard ratio [HR], 3.04; confidence interval [CI], 1.681–5.512; *P* $< .001$) and preoperative creatinine ≥ 1.1 mg/ml (HR, 3.57; 95% CI, 2.050–6.202; *P* $< .001$) were independent predictors (Table 3).

Table 1**Baseline clinical and demographic characteristics.**

| Variables | Total | RENAL score ≤ 7 | RENAL score ≥ 8 | P value |
|--|-------------------|-------------------|-------------------|---------|
| | 393 | 258 (65.6) | 135 (34.4) | |
| Age (y) | 53.0 (45.0–63.0) | 53.5 (45.0–64.0) | 53.0 (45.0–61.0) | .346 |
| Sex | | | | |
| Male | 261 (66.4) | 91 (35.3) | 41 (30.4) | .330 |
| Female | 132 (33.6) | 167 (64.7) | 94 (69.6) | |
| BMI (kg/m ²) | 24.3 (22.2–26.5) | 24.5 (22.1–26.6) | 24.0 (22.4–26.1) | .865 |
| Comorbidity | | | | |
| ASA ≥ 2 | 150 (38.2) | 99 (38.4) | 51 (37.8) | .916 |
| Method | | | | |
| Open | 118 (30.0) | 100 (38.8) | 18 (13.3) | <.001 |
| Laparoscopy | 53 (13.5) | 44 (17.1) | 9 (6.7) | |
| Robot | 222 (56.5) | 114 (44.2) | 108 (80.0) | |
| Tumor size (cm) | 2.6 (1.9–3.8) | 2.3 (1.6–3.0) | 3.8 (2.7–5.0) | .002 |
| Fuhrman grade | | | | |
| ≥ 3 | 95 (24.2) | 52 (20.2) | 95 (70.4) | .011 |
| Preoperative Cr (mg/ml) | 0.9 (0.73–1.02) | 0.88 (0.70–1.00) | 0.91 (0.74–1.04) | .239 |
| Preoperative eGFR (ml/minute/1.73 m ²) | 82.2 (68.0–104.7) | 82.8 (67.7–104.1) | 81.0 (68.1–105.7) | .596 |
| Operative time (minute) | 172 (130–219) | 176 (135–220) | 169 (126–212) | .398 |
| Pedicle clamp | | | | |
| Total | 294 (74.8) | 193 (74.8) | 101 (74.8) | .373 |
| Selective | 63 (16.0) | 36 (14.0) | 27 (20.0) | |
| No | 36 (9.2) | 29 (11.2) | 7 (5.2) | |
| Estimated blood loss (ml) | 400 (200–600) | 400 (200–600) | 400 (200–625) | .226 |
| Complications (Clavien-Dindo classification) | | | | |
| G1 | 25 (6.4) | 12 (4.7) | 13 (9.6) | .005 |
| G2 | 14 (3.6) | 5 (1.9) | 9 (6.7) | |
| G3 | 20 (5.1) | 10 (3.9) | 10 (7.4) | |
| Ischemic time (minute) | 25.0 (18.0–31.0) | 23.0 (17.0–30.0) | 27.5 (22.8–32.3) | .005 |
| Positive surgical margin | 27 (6.9) | 11 (4.3) | 16 (11.9) | .014 |
| Renal insufficiency | 54 (13.7) | 35 (13.6) | 19 (14.1) | .943 |
| Renal function recovery | 18 (4.6) | 11 (4.3) | 7 (5.2) | .694 |
| Trifecta achievement | 172 (43.8) | 130 (50.4) | 42 (31.1) | <.001 |
| Pentafecta achievement | 149 (37.9) | 111 (43.0) | 38 (28.1) | .004 |
| Follow-up period (months) | 24.0 (6.0–36.0) | 24.0 (6.0–36.0) | 24.0 (6.0–48.0) | .916 |

Data are n (%) or median (interquartile range).

ASA=American Society of Anesthesiologists, BMI=body mass index, Cr=creatinine, eGFR=estimated glomerular filtration rate.

Among these patients, 18/54 (33.3%) patients exhibited renal function recovery within a median (interquartile range) period of 18 (12–36) months, of which preoperative creatinine <1.1 mg/ml was an independent predictor (HR, 4.38; 95% CI, 1.203–15.911; $P=.025$) (Table 4).

4. Discussion

The increasing use of abdominal imaging including ultrasonography, CT, and MRI is the main cause for the increasing detection rate of renal masses.^[13] With the dramatically changing epidemiology of renal masses, PN has been considered the

Table 2**Change in median eGFR over time according to clamping type, RENAL nephrometry score, and WIT.**

| | Median (IQR) eGFR, ml/minute/1.73 m ² | | | P value for Wilcoxon signed-rank test | | |
|-------------------------|--|-------------------|------------------|---------------------------------------|-------------------|---------------------|
| | Preoperative | 3 months | 2 years | 2 years vs preop | 3 months vs preop | 2 years vs 3 months |
| Overall | 82.2 (68.0–104.7) | 75.9 (64.4–91.6) | 79.4 (66.3–94.4) | .865 | .005 | .026 |
| Clamping type | | | | | | |
| Total | 81.6 (67.6–104.5) | 75.2 (63.9–89.3) | 78.8 (65.8–93.8) | .652 | .003 | .026 |
| Selective | 83.7 (68.0–100.1) | 85.4 (70.2–109.4) | 77.3 (61.6–85.9) | .394 | .480 | .754 |
| No | 87.5 (74.6–121.4) | 77.3 (61.6–85.9) | 76.8 (63.3–87.4) | .695 | .073 | .110 |
| RENAL nephrometry score | | | | | | |
| <7 | 82.8 (67.7–104.1) | 77.9 (61.5–95.0) | 80.8 (66.9–95.7) | .606 | .046 | .104 |
| ≥ 8 | 81.0 (68.1–105.7) | 72.7 (64.1–86.8) | 78.2 (64.6–89.0) | .390 | .046 | .122 |
| WIT | | | | | | |
| < 25 minutes | 83.2 (68.9–105.4) | 77.9 (66.2–93.2) | 80.9 (70.0–97.1) | .050 | .324 | .020 |
| ≥ 25 minutes | 81.6 (67.6–104.5) | 74.4 (64.0–89.2) | 76.5 (63.8–90.6) | .043 | .005 | .366 |

eGFR=estimated glomerular filtration rate, IQR=interquartile range, WIT=warm ischemic time.

Table 3
Multivariate analysis for predicting renal insufficiency after partial nephrectomy.

| | Univariate | | | Multivariate | | |
|-------------------------------------|------------|-------------|---------|--------------|-------------|---------|
| | HR | 95% CI | P value | HR | 95% CI | P value |
| Age ≥ 60 y | 3.90 | 2.214–6.881 | <.001 | 3.04 | 1.681–5.512 | <.001 |
| Male | 1.60 | 0.859–2.990 | .139 | | | |
| BMI | 1.02 | 0.952–1.095 | .564 | | | |
| Diabetes mellitus | 1.54 | 0.479–4.934 | .470 | | | |
| ASA ≥ 2 | 1.89 | 1.093–3.253 | .023 | 1.49 | 0.851–2.621 | .162 |
| Operative methods | | | .189 | | | |
| Open | Reference | | | | | |
| Laparoscopy | 0.56 | 0.058–5.341 | .611 | | | |
| Robot | 2.33 | 0.671–8.119 | .183 | | | |
| Tumor size | 0.93 | 0.776–1.111 | .417 | | | |
| Pedicle clamp | 1.62 | 0.875–2.995 | .125 | | | |
| WIT > 25 minutes | 1.25 | 0.731–2.137 | .416 | | | |
| Preoperative creatinine ≥ 1.1 mg/ml | 3.88 | 2.264–6.635 | <.001 | 3.57 | 2.050–6.202 | <.001 |
| RENAL nephrometry score (≥8) | 1.08 | 0.618–1.889 | .786 | | | |

ASA = American Society of Anesthesiologists, BMI = body mass index, CI = confidence interval, HR = hazard ratio, WIT = warm ischemic time.

paramount treatment option for patients with small renal masses. A satisfactory oncologic outcome is the cardinal goal of all surgical methods. Trifecta and penta-fecta have been proposed as measurement tools for the outcomes of PN. Recently published data reported that trifecta outcomes ranged from 31% to 78%.^[8,9,14,15] Because of the lack of a definition of trifecta for PN cases and variations in surgical approach and technique, PN studies using trifecta showed heterogeneous outcomes. We used complications (Clavien-Dindo complication grade ≥3), PSMs, and WIT ≥25 minutes to evaluate the trifecta achievement rate. Our achievement rate of trifecta was 43.8%, and this rate is at least partially similar to the results of previous studies.

Moreover, concerns regarding preservation of renal function emerged with the increasing demand for PN. Postoperative renal function after PN is affected by various factors, including decreasing WIT to reduce the ischemic injury, loss of normal parenchymal volume to prevent positive surgical margins, and ischemic damage in preserved tissue during the operation.^[16,17]

Table 4
Predictors of renal function recovery in patients with renal insufficiency after partial nephrectomy.

| | Univariate | | |
|-------------------------------------|------------|--------------|---------|
| | HR | 95% CI | P value |
| Age ≥ 60 y | 0.48 | 0.147–1.567 | .224 |
| Male | 0.74 | 0.203–2.717 | .653 |
| BMI | 0.95 | 0.792–1.132 | .552 |
| Diabetes mellitus | 0.94 | 0.079–11.143 | .962 |
| ASA ≥ 2 | 0.57 | 0.177–1.809 | .337 |
| Operative methods | | | |
| Open | Reference | | |
| Laparoscopy | 0.50 | 0.044–5.700 | .577 |
| Robot | 3.50 | 0.817–14.986 | .091 |
| Tumor size | 1.12 | 0.732–1.726 | .593 |
| Pedicle clamp | 0.87 | 0.255–2.994 | .830 |
| WIT > 25 minutes | 1.41 | 0.444–4.448 | .562 |
| Preoperative creatinine < 1.1 mg/ml | 4.375 | 1.203–15.911 | .025 |
| RENAL nephrometry score (≥8) | 1.27 | 0.393–4.117 | .687 |

ASA = American Society of Anesthesiologists, BMI = body mass index, CI = confidence interval, HR = hazard ratio, WIT = warm ischemic time.

The optimal threshold of WIT to prevent renal function deterioration remains controversial. In the trifecta standard, WIT <25 minutes was used, but recent papers proposed the optimal threshold of WIT to be within 20 minutes.^[18,19] Moreover, Thompson et al found that in patients with a solitary kidney, every minute of clamping has short- and long-term renal consequences.^[20] In this study, when analyzing WIT at each minute from 20 to 30 minutes, no significant cutoff point was found. However, WIT ≥25 minutes was a factor associated with prolonged recovery or no recovery of the preoperative eGFR.

Postoperative renal function was related to loss of normal parenchymal volume and ischemic damage in preserved tissue.^[21,22] Mir MC et al reported that renal function ultimately was primarily associated with parenchymal volume preservation, whereas ischemia played a role in the level of renal function present 4 to 12 months after PN.^[22] However, our multicenter analysis based on long-term follow-up demonstrated that the type of ischemia was not related to renal function. The results of this study are expected to be interpreted as a result of compensatory growth in the contralateral kidney and a more refined assessment of the effect of ischemia.

The role of various factors influencing the recovery of renal function after PN have been investigated, including diabetes, hypertension, hypercholesterolemia, male sex, old age, smoking, BMI ≥30 kg/m², vascular disease, use of anti-thrombolytic medication, remnant renal volume, and surgical methods and type.^[23] However, in our study, only the factor associated with preoperative renal function, creatinine <1.1 mg/ml, was found to affect renal function recovery.

The RENAL nephrometry score reflects the anatomic and surgical complexity. Several studies investigated the RENAL nephrometry score as a tool to predict factors associated with increased operative time, EBL, total renal volume loss, WIT, and complications.^[24,25] In the present study, as expected, WIT, complication rate, and positive surgical margins statistically increased with increasing RENAL nephrometry score. However, no study has predicted long term-renal function using the RENAL nephrometry score. Recently, Husain et al reported that healthy renal volume loss or non-neoplastic parenchymal volume was associated with RENAL nephrometry score.^[26] Simmons

MN reported that the RENAL nephrometry score was associated with changes in the percent of functional volume preservation and the perioperative functional decrease in an analysis of 237 patients who underwent PN from 2007 to 2010.^[24] We found that the eGFR at 2 years after PN, regardless of complexity, was not inferior to the preoperative eGFR. To our knowledge, this is the first study to show that the RENAL nephrometry score is not a prognostic factor predicting renal insufficiency and renal function recovery.

This study has several limitations in addition to its retrospective design. Heterogeneity of intraoperative management existed. Patient selection, multiple surgical methods, and techniques influenced by the physician's preference could account for the heterogeneity in our results. We do not have data on the number of cases that underwent radical nephrectomy because PN was not recommended for small renal mass. In addition, measurement of the RENAL nephrometry score and pathologic results were not centrally investigated but were assessed by urologists and pathologists at each institution. Nevertheless, we believe that this effect may reflect real-world clinical practice and is inherent in any retrospective study. Finally, we evaluated the parameters to predict renal function recovery; however, previous studies have investigated renal functional compensation as a phenomenon of renal compensatory adaptation.^[21,22] In the future, we plan to evaluate the contribution of ipsilateral atrophy or contralateral hypertrophy to renal functional recovery after partial nephrectomy.

5. Conclusion

Age ≥ 60 years and preoperative creatinine ≥ 1.1 mg/ml were risk factors for renal insufficiency following PN. Patients with renal insufficiency whose preoperative creatinine was < 1.1 mg/ml were more likely to have renal function recovery.

Author contributions

Conceptualization: Kyo Chul Koo.

Data curation: Dae Keun Kim, Kwang Hyun Kim, Woo Jin Bang, Hyung Joon Kim, Sung Yul Park, Koon Ho Rha, Byung Ha Chung, Jin Seon Cho.

Formal analysis: Kyo Chul Koo.

Writing – original draft: Kwang Suk Lee, Kyo Chul Koo.

References

- [1] Motzer RJ, Jonasch E, Agarwal N, et al. Kidney Cancer, Version 2.2017, NCCN clinical practice guidelines in oncology. *J Natl Compr Canc Netw* 2017;15:804–34.
- [2] Lee CT, Katz J, Shi W, et al. Surgical management of renal tumors 4 cm or less in a contemporary cohort. *J Urol* 2000;163:730–6.
- [3] Zini L, Perrotte P, Capitanio U, et al. Radical versus partial nephrectomy: effect on overall and noncancer mortality. *Cancer* 2009;115:1465–71.
- [4] Thompson RH, Boorjian SA, Lohse CM, et al. Radical nephrectomy for pT1a renal masses may be associated with decreased overall survival compared with partial nephrectomy. *J Urol* 2008;179:468–71. discussion 72–3.
- [5] Weight CJ, Larson BT, Gao T, et al. Elective partial nephrectomy in patients with clinical T1b renal tumors is associated with improved overall survival. *Urology* 2010;76:631–7.
- [6] MacLennan S, Imamura M, Lapitan MC, et al. Systematic review of oncological outcomes following surgical management of localised renal cancer. *Eur Urol* 2012;61:972–93.
- [7] Shin SJ, Ko KJ, Kim TS, et al. Trends in the use of nephron-sparing surgery over 7 Years: an analysis using the R.E.N.A.L. nephrometry scoring system. *PLoS One* 2015;10:e0141709.
- [8] Hung AJ, Cai J, Simmons MN, et al. Trifecta™ in partial nephrectomy. *J Urol* 2013;189:36–42.
- [9] Kim DK, Kim LH, Raheem AA, et al. Comparison of trifecta and pentafta outcomes between T1a and T1b renal masses following Robot-Assisted Partial Nephrectomy (RAPN) with minimum one year follow up: can RAPN for T1b renal masses be feasible? *PLoS One* 2016;11:e0151738.
- [10] Kutikov A, Uzzo RG. The R.E.N.A.L. nephrometry score: a comprehensive standardized system for quantitating renal tumor size, location and depth. *J Urol* 2009;182:844–53.
- [11] Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 2004;240:205–13.
- [12] Levey AS, Bosch JP, Lewis JB, et al. A more accurate method to estimate glomerular filtration rate from serum creatinine: a new prediction equation. Modification of diet in renal disease study group. *Ann Intern Med* 1999;130:461–70.
- [13] Chow WH, Devesa SS, Warren JL, et al. Rising incidence of renal cell cancer in the United States. *JAMA* 1999;281:1628–31.
- [14] Zargar H, Allaf ME, Bhayani S, et al. Trifecta and optimal perioperative outcomes of robotic and laparoscopic partial nephrectomy in surgical treatment of small renal masses: a multi-institutional study. *BJU Int* 2015;116:407–14.
- [15] Khalifeh A, Autorino R, Hillyer SP, et al. Comparative outcomes and assessment of trifecta in 500 robotic and laparoscopic partial nephrectomy cases: a single surgeon experience. *J Urol* 2013;189:1236–42.
- [16] Shin TY, Komninos C, Kim DW, et al. A novel mathematical model to predict the severity of postoperative functional reduction before partial nephrectomy: the importance of calculating resected and ischemic volume. *J Urol* 2015;193:423–9.
- [17] Komninos C, Shin TY, Tulliao P, et al. Renal function is the same 6 months after robot-assisted partial nephrectomy regardless of clamp technique: analysis of outcomes for off-clamp, selective arterial clamp and main artery clamp techniques, with a minimum follow-up of 1 year. *BJU Int* 2015;115:921–8.
- [18] Becker F, Van Poppel H, Hakenberg OW, et al. Assessing the impact of ischaemia time during partial nephrectomy. *Eur Urol* 2009;56:625–34.
- [19] Song C, Park S, Jeong IG, et al. Followup of unilateral renal function after laparoscopic partial nephrectomy. *J Urol* 2011;186:53–8.
- [20] Thompson RH, Lane BR, Lohse CM, et al. Every minute counts when the renal hilum is clamped during partial nephrectomy. *Eur Urol* 2010;58:340–5.
- [21] Simmons MN, Hillyer SP, Lee BH, et al. Functional recovery after partial nephrectomy: effects of volume loss and ischemic injury. *J Urol* 2012;187:1667–73.
- [22] Mir MC, Campbell RA, Sharma N, et al. Parenchymal volume preservation and ischemia during partial nephrectomy: functional and volumetric analysis. *Urology* 2013;82:263–8.
- [23] Kocher NJ, Rjepaj C, Robyak H, et al. Hypertension is the primary component of metabolic syndrome associated with pathologic features of kidney cancer. *World J Urol* 2017;35:67–72.
- [24] Simmons MN, Hillyer SP, Lee BH, et al. Nephrometry score is associated with volume loss and functional recovery after partial nephrectomy. *J Urol* 2012;188:39–44.
- [25] Meyer A, Woldu SL, Weinberg AC, et al. Predicting renal parenchymal loss after nephron sparing surgery. *J Urol* 2015;194:658–63.
- [26] Husain FZ, Rosen DC, Paulucci DJ, et al. R.E.N.A.L. nephrometry score predicts non-neoplastic parenchymal volume removed during robotic partial nephrectomy. *J Endourol* 2016;30:1099–104.