

## 터어키안 주변종양에 대한 감마나이프 방사선 수술\*

장종희 · 장진우 · 박용구 · 정상섭

= Abstract =

### Gamma Knife Radiosurgery for Juxtaseilar Tumors

Jong Hee Chang, M.D., Jin Woo Chang, M.D.,  
Yong Gou Park, M.D., Sang Sup Chung, M.D.

*Department of Neurosurgery, Brain Research Institute, Yonsei University College of Medicine, Seoul, Korea*

**O**bjective : Around the sellar area, there are many important structures. But, the optimal radiation dosage for minimal toxicity to surrounding neural tissue has not been firmly established. The purpose of this study is to evaluate the radiosurgical outcome of juxtaseilar tumors and to investigate the relationship between radiation dosage and toxicity to neural tissue.

**Method** : Between May 1992 and June 2000, we treated 65 juxtaseilar tumors by using the Leksell Gamma Knife. Among them, 52 patients who could be followed more than 1 year were included in this study. The radiosurgical dosage to the optic pathway, cavernous sinus, Meckel's cave, hypothalamus, pituitary gland and stalk, and brain stem was analyzed and correlated with clinical outcome. The mean follow-up period was 33.5 months (range 12.2 - 99.0 months).

**Result** : The clinical response rate was 69.2%. The volume response rate was 61.0% and the radiologic control rate was 92.7%. There were 4 complications (7.7%) of 2 trigeminal neuropathy, 1 abducens nerve palsy, and 1 trigeminal and transient abducens nerve palsy. The optic apparatus appeared to tolerate doses greater than 10Gy. The risk of cranial nerve complications in cavernous sinus seemed to be related to doses of more than 16Gy. In 3 of 4 patients who received more than 16Gy to cavernous sinus, the abducens or trigeminal neuropathy occurred. Also, one patient who received more than 15Gy to the Meckel's cave, trigeminal neuropathy developed. The hypothalamus, pituitary gland and stalk, and brain stem were relatively tolerable to radiation.

**Conclusion** : Gamma Knife radiosurgery seems to be an effective method to control the growth of juxtaseilar tumors. To avoid injury to surrounding important neural tissue, careful dose planning and further study for radiation toxicity to neural tissue were needed.

**KEY WORDS** : Radiosurgery · Gamma Knife · Juxtaseilar · Optic nerve · Cranial nerve.

서 론

ning)

(dose plan -

2)6-8)11)14)15).

, Meckel's cave,  
가

1999



27.1Gy(10.0 34.0Gy) .  
 40 90%  
 13.7Gy(5 17.5Gy) . 40  
 26.7Gy, 13.6  
 Gy, 12 25.4Gy, 12.5Gy ,  
 1 60Gy, 30  
 Gy

5. 임상적 반응

가 21 ,  
 가 15  
 (clinical response rate) 69.2% (Fig. 1).  
 13 65  
 가 12 , 4  
 1 (Table 3).

6. 방사선학적 반응

MRI CT 가 41 ,  
 25 가  
 (volume response rate) 61.0%




Fig. 1. Cumulative events curve by Kaplan-Meier survival analysis for clinical response of patients with juxtaseellar tumor since Gamma Knife radiosurgery.

Table 3. Clinical response of juxtaseellar tumors after Gamma Knife radiosurgery

Clinical results	No. of cases(%)
Improved**	21 (40.4%)
No new symptom*	15 (28.8%)
Sustained	12 (23.1%)
Aggravated	4 ( 7.7%)
Total	52 (100%)

\* : no newly developed symptom in asymptomatic patients  
 \*\* : clinical response, 36/52 patients(69.2%)

가 13 , 가 가 가 3 (Table 4),  
 (radiologic control rate) 92.7% (Fig. 2, 3).

7. 치료 실패

가 가 가 3  
 (clivus chordoma)  
 8.4  
 (sarcomatous change)  
 가 (cystic adenocar-  
 cinoma)  
 9.2 2  
 (nasopharyngeal carcinoma)

8. 합병증

52 4 (7.7%)  
 가 , 1  
 가

Table 4. Radiologic response of juxtaseellar tumors after Gamma Knife radiosurgery

Volume change	No. of cases(%)
Shrunken (<80%)*	25 (61.0%)
Static (80 - 120%)**	13 (31.7%)
Enlarged (>120%)	3 ( 7.3%)
Total	41 (100%***)

\* : volume response  
 \* , \*\* : radiologic control, 38/41 patients(92.7%)  
 \*\*\* : MRI follow-up in 41 patients(78.8%)





Fig. 2. Cumulative events curve by Kaplan-Meier survival analysis for radiologic control of patients with juxtaseellar tumor since Gamma Knife radiosurgery.



**Fig. 3.** A : Pre-radiosurgical MRI showing left cavernous sinus and Meckel's cave meningioma. B : Follow-up MRI(14 months after Gamma Knife radiosurgery) showing sh-rinked tumor.

가 2  
1 .

9. 시신경계의 방사선 조사량

5 47  
. 8Gy 가 가 40 , 8  
10Gy가 가 5 10Gy 가  
,  
13Gy 14Gy가 (Table  
5).  
, 10Gy 14.

9 40.5 .

10. 해면정맥동의 방사선 조사량

, 16Gy 가 48  
. 16Gy 16.2Gy  
17.4 , 16.5Gy  
6 가 . 18Gy  
18.5Gy가 2  
가 (Table 6).

11. Meckel's cave의 방사선 조사량  
Meckel's cave 1 15Gy 가  
, . 1

**Table 5.** Radiation dose to optic pathway (n = 47)\*

Dose range(Gy)	No. of cases
<8	40
8 - 10	5
10 - 12	0
12 - 14	2

\* : no visual complication

**Table 6.** Radiation dose to cavernous sinus (n = 52)

Dose (Gy)*	No. of cases	Complications
<15	46	none
16.0	2	none
16.2	1	none
16.5	1	V **
18.0	1	VI ***
18.5	1	VI (transient), V


\* : radiation dose to the lateral wall of cavernous sinus

\*\* : decreased facial sensation

\*\*\* : limitation on lateral gaze

Meckel's cave 17.5Gy가  
, 가 6  
, 14

12. 기타 주요구조물의 방사선 조사량  
, , 13Gy,  
21.15Gy, 14.25Gy가



**Fig. 4.** A : Preoperative MRI showing suprasellar craniopharyngioma. B : Postoperative and pre-radiosurgical MRI showing residual tumor near hypothalamus and optic pathway. C-F : Serial follow-up MRI (4, 13, 24, and 35 months after Gamma Knife radiosurgery, respectively) showing gradually decreased tumor remnant.

kel's cave,

13. 종양에 적정량이하의 방사선이 조사된 경우 11)

12Gy 가 가 9 5  
 MRI 5Gy 9.5Gy가  
 12 35  
 가 , 8 8.5Gy가 가  
 3 가 가 가  
 (Fig. 4). 가 (Schwann's cell) 10)

고 찰

가 가 10Gy 가 가 8  
 2)11), 가 가  
 3)6)8-10)13)14) Leber 7)8)

50 10Gy , 10 15Gy,

Mec -

15Gy 가 0%, 26.7%, 77.8% 가 10Gy 가 16Gy . Meckel's cave 15Gy , Me-

가 , Morita <sup>11)</sup> 가 8Gy 가 Meckel's cave 17.5Gy가 1 , dose) 88 59 8Gy ( 16Gy ) 12 16Gy가 (9 12mm 21Gy, 13Gy, 14Gy 가 가 10Gy . 2 52 14Gy가 1 13Gy, 1 41 , 가 38 (92.7%) 52 14.9 40.5 , 2 36 (69.2%) , 69.2% 가 10Gy 가 (27/39) , 39 50% <sup>1)</sup>, 30Gy 5 3, 4, 5, 6 가 가 가 <sup>7)</sup>. Tishler <sup>14)</sup> 12Gy <sup>4)</sup>. 가 가 9 5 MRI 3 8 8.5Gy 가 3, 4, 6 4.5 30Gy가 , 5 5 가 Leber <sup>7)</sup> 가 가 20Gy가 , Morita <sup>11)</sup> Meckel's cave 19Gy <sup>2)6)12)</sup>. 5Gy가 1 9.5Gy가 가 가 16Gy 가 48 . Iwai <sup>5)</sup> 24 . 16Gy 16.2Gy 17.4 , 8 15Gy (median, 10.6Gy) 16.5Gy, 18Gy, 18.5Gy가 3

가

## 결 론

dose plan  
가

- : 2000 8 16
  - : 2000 10 12
  - :  
120 - 752 134
- : 02) 361 - 5625, : 02) 393 - 9979  
E - mail : ygpark@yumc.yonsei.ac.kr

## References

- 1) Chang JW, Kim SH, Huh R, Park YG, Chung SS : *The effects of stereotactic radiosurgery on secondary facial pain. Stereotact Funct Neurosurg 72 (suppl 1) : 29-37, 1999*
- 2) Chung WY, Pan HC, Guo WY, Shiau CY, Wang LW, Wu HM, et al : *Protection of visual pathway in Gamma Knife radiosurgery for craniopharyngiomas. Stereotact Funct Neurosurg 70 (suppl 1) : 139-151, 1998*
- 3) Duma C, Lunsford LD, Kondziolka D, Harsh GR IV, Flickinger JC : *Stereotactic radiosurgery of cavernous sinus meningiomas as an addition or alternative to microsurgery. Neurosurgery 32 : 699-704, 1993*
- 4) Ganz JC, Backlund EO, Thorsen FA : *The results of Gamma Knife surgery of meningiomas, related to size of tumor and dose. Stereotact Funct Neurosurg 61 (suppl 1) : 23-29, 1993*
- 5) Iwai Y, Yamanaka K, Yasui T, Komiyama M, Nishikawa M, Nakajima H, et al : *Gamma Knife surgery for skull base meningiomas. The effectiveness of low-dose treatment. Surg Neurol 52 : 40-45, 1999*
- 6) Kobayashi T, Tanaka T, Kida Y : *Stereotactic Gamma radiosurgery of craniopharyngiomas. Pediatr Neurosurg 21 (suppl 1) : 69-74, 1994*
- 7) Leber KA, Bergloff J, Langmann G, Mokry M, Schrottner O, Pendl G : *Radiation sensitivity of visual and oculomotor pathways. Stereotact Funct Neurosurg 64 (suppl 1) : 233-238, 1995*
- 8) Leber KA, Bergloff J, Pendl G : *Dose-response tolerance of the visual pathways and cranial nerves of the cavernous sinus to stereotactic radiosurgery. J Neurosurg 88 : 43-50, 1998*
- 9) Liscak R, Simonova G, Vymazal J, Janouskova L, Vladyka V : *Gamma Knife radiosurgery of meningiomas in the cavernous sinus region. Acta Neurochir (Wien) 141 : 473-480, 1999*
- 10) Mehta MP, Kinsella TJ : *Cavernous sinus cranial neuropathies : is there a dose-response relationship following radiosurgery? Int J Radiat Oncol Biol Phys 27 : 477-480, 1993 (editorial)*
- 11) Morita A, Coffey RJ, Foote RL, Schiff D, Gorman D : *Risk of injury to cranial nerves after gamma knife radiosurgery for skull base meningiomas : experience in 88 patients. J Neurosurg 90 : 42-49, 1999*
- 12) Prasad D, Steiner M, Steiner L : *Gamma Knife surgery for craniopharyngioma. Acta Neurochir (Wien) 134 : 167-176, 1995*
- 13) Shaw EG, Coffey RJ, Dinapoli RP : *Neurotoxicity of radiosurgery. Semin Radiat Oncol 5 : 235-245, 1995*
- 14) Tishler RB, Loeffler JS, Lunsford LD, Duma C, Alexander E III, Kooy HM, et al : *Tolerance of cranial nerves of the cavernous sinus to radiosurgery. Int J Radiat Oncol Biol Phys 27 : 215-221, 1993*
- 15) Urie MM, Fullerton B, Tatsuzaki H, Birnbaum S, Suit HD, Phil D, et al : *A dose response analysis of injury to cranial nerves and/or nuclei following proton beam radiation therapy. Int J Radiat Oncol Biol Phys 23 : 27-39, 1992*