

Nitric Oxide

= Abstract =

Nitric Oxide Production of Rat Osteoblast Cells by Pulsed Electromagnetic Field Radiation

Seong-Hwan Moon, M.D., Jin-Woo Lee, M.D., Jun-Seop Jahng, M.D.

Department of Orthopaedic Surgery Yonsei University College of Medicine, Seoul, Korea

Experimental study was conducted to prove the effect of pulsed electromagnetic field(PEMF) on the production of Nitric oxide(NO) from the cultured rat osteoblast-like cells. Calvarium of thirty Sprague-Dawley rats was digested by sequential collagenase and cultured in-vitro. The osteoblast cell phenotype was confirmed by expression of osteocalcin by immunohistochemistry. PEMF was generated and applied to cultured osteoblast cells. Production of NO was measured by Greiss reaction. Ten minute exposure of PEMF to osteoblast cell showed increased NO content at 24 and 48 hours(p<0.05). Cultures with different duration of PEMF exposure(10, 20, 30 60 minutes) demonstrated similar responses. In conclusion. this study proved that NO can be generated with PEMF which support the notion that NO can be a possible mediator of PEMF on bone metabolism.

Key Words : Osteoblast, Pulsed electromagnetic field, Nitric oxide

Nitric Oxide(NO) Free radical NO가
 가

17) , 4,8)

134

Tel : 02) 361-5649, Fax : 02) 363-1139
E-mail : shmoon@yumc.yonsei.ac.kr

* 1996

4)
 (osteoblast) inter-
 leukin-1, TNF- α , interferon-
 NO²¹⁾, dexamethasone, 1.
 cyclohexamide, PTH, Vitamine D, TGF- β
 가^{17,21)}. Estrogen 1 Sprague-Dawley rat 30
 NO²⁶⁾
 calcitonin cAMP 가가 inducible Phosphate buffered
 nitric oxide synthase(NOS) mRNA saline(PBS) 1-2mm³
 가 NO 가 가 PBS
 NO¹⁷⁾ . 0.1% collagenase, 0.05%
 NO²²⁾ trypsin, 136mM NaCl, 2.6mM KCl,
 0.36mM NaH₂PO₄, 5mM EDTA
 37° 20
 7) . 4 . 3, 4
 nylon mesh
 6) Dulbeccó's
 modified eagle's medium (DMEM) 10cc
 2 .
 10% Fetal calf serum(FCS)
 1,2,9,11,12) DMEM 가 5×10⁶/M ϕ
 tissue culture flask(25cm²)
 CO₂ incubator (37°C, 5%
 4 . 24 , 3 , 7
 3), BMP-2, BMP- CO₂). 7 100%
 lysozyme
 19), 13), DNA .
 가¹⁸⁾ .
 NO가¹⁶⁾ 2. (Osteocalcin)
 NO 7 4% paraformaldehyde
 NO 4 °C protein blocking agent
 가 5 . Rat osteocalcin anti-
 가 NO serum(1:1000) 4 °C
 NO . biotin-strepta
 NO 가 . vidin-peroxidase (Lipshaw, Pittsburgh,
 NO . PA) immunoreactivity
 가 NO 3.
 McLeod¹⁵⁾

30cm polyethylene tube 12cm NO (normalization)

3 650 NO 3

1.8millitelsa(mT) magnetic flux density (Walker Mode MG2A gaussmeter) 30Hz sinusoidal 1.8mT NO t-test

30Hz sinusoidal 1.8mT NO (ANOVA) p<0.05

4.

1) (10) NO 1.

(50μl) 10 CO₂ incubator 48 6 osteoclastin 90% (Fig. 1).

2) NO 10, 20, 30, 60 CO₂ incubator 가 2. (10) NO 10 6

48 80.

5. NO

NO nitrite(NO₂) Greiss metric assay NO Greiss NO

NO Nitrite spectrophoto NO₂- NO₃- NO₂- 24).




Fig. 1. Immunohistochemical staining for osteocalcin in cultured osteoblast-like cells from calvarium of Sprague-Dawley rat(X 200)

tor, microfilament

Integrin recep-

NO

23,25)

NO

(depolarization)
가
NOS가 가 NO가
가
NOS

REFERENCES

- 1) **Adam BD, Frykman GK and Taleisnik J** : Treatment of scaphoid nonunion with casting and pulsed electromagnetic fields: A study continuation. *J Hand Surg*, 17:910-914, 1992.
- 2) **Basset CAI, Pawluk RJ and Pilla AA** : Augmentation of bone repair by inductively coupled electromagnetic fields. *Science*, 184:575-577, 1974.
- 3) **Bodamyali T, Bhatt B, Hughes FJ, Winrow VR, Kanczler JM, Simon B, Abbott J, Blake DR and Stevens CR** : Pulsed electromagnetic fields simultaneously induce osteogenesis and upregulate transcription of bone morphogenic proteins 2 and 4 in rat osteoblasts in vitro. *Biochem Biophys Res Comm*, 250:458-461, 1998.
- 4) **Collin-Osdoby P, Allen Nickols G and Osdoby P** : Bone cell function, regulation, and communication: A role for nitric oxide. *J Cell Biochem*, 57:399-408, 1995.
- 5) **Conti P, Gigante GE, Cifone MG, Alesse E, Ianni G, Reale M and Angeletti PU** : Reduced mitogenic stimulation of human lymphocytes by extremely low frequency electromagnetic fields. *FEBS letters*, 162:156-160, 1983.
- 6) **Enzler MA, Summer-Smith G, Waelchli-Suter C and Perren SM** : Treatment of nonuniting osteotomies with pulsating electromagnetic fields. A controlled animal experiment. *Clin Orthop*, 187:272-276, 1984.

NO

NO

inducible NOS

constitutive NOS

NO

Miura ¹⁶⁾
NO

가
(
5)

1,2,9,11,12),
NO

14),

NO

cytokine

- 7) **Evans DM and Ralston SH** : Nitric oxide and bone. *J Bone Miner Res*, 11:300-305, 1996.
- 8) **Evans CH, Stefanovic-Racic M and Lancaster J** : Nitric Oxide and its role in orthopaedic disease. *Clin Orthop*, 312:275-294, 1995.
- 9) **Glazer PA, Heilmann MR, Lotz JC and Bradford DS** : Use of electromagnetic fields in a spinal fusion. A Rabbit model. *Spine*, 22:2351-2356, 1997
- 10) **Green LC, Wagner DA, Glogowski J, Skipper PL, Wishnok JS and Tannenbaum SR** : Analysis of nitrite and [15N] nitrate in biological fluid. *Ann Biochem*, 126:131-138, 1982.
- 11) **Heckman JD, Ingram AJ, Lyod RD, Luck JV and Mayer PW** : Nonunion treatment with pulsed electromagnetic fields. *Clin Orthop*, 161:58-66, 1981.
- 12) **Holmes GB** : Treatment of delayed unions and nonunions of the proximal fifth metatarsal with pulsed electromagnetic fields. *Foot Ankle Int*, 15:552-556, 1994.
- 13) **Luben RA** : Effects of low-energy electromagnetic fields on membrane signal transduction process in biological system. *Heath Physics*, 61:15-28, 1991.
- 14) **Lyle DB, Schechter P, Adley WR and Lundak RL** : Suppression of T-lymphocyte cytotoxicity following exposure to sinusoidal amplitude-modulated fields. *Bio - electromag*, 4:281-292, 1983.
- 15) **McLeod KJ, Donahue HJ, Levin PE, Fontaine MA and Rubin CT** : Electric Fields modualte bone cell function in a density-dependent manner. *J Bone Miner Res*, 8:977-984, 1993.
- 16) **Miura M, Takayama K and Okada J** : Increased nitric oxide and cyclic GMP of rat cerebellum by radio frequency burst type electromagnetic field radiation. *J Physiol*, 461:513-524, 1993.
- 17) **Moncada S, Palmer RMJ and Higgs EA** : Nitric oxide: Physiology, pathophysiology, and pharmacology. *Pharm Rev*, 40:109-142, 1991.
- 18) **Noda M, Johnston DE, Chiabrera A and Rodan GA** : Effect of electric currents on DNA synthesis in rat osteosarcoma cells: dependence on conditions that influence cell growth. *J Orthop Res*, 5:253-260, 1987.
- 19) **Norton LA** : Effects of a pulsed electromagnetic field on a mixed chondroblastic tissue culture. *Clin Orthop*, 167:280-290, 1982.
- 20) **Ralston SH, Ho L-P, Helfrich MH, Grabowski PS, Johnston PW and Benjamin N** : Nitric oxide: A cytokine-induced regulator of bone resorption. *J Bone Miner Res*, 10:1040-1049, 1995.
- 21) **Ralston SH, Todd D, Helfrich M, Benjamin N and Grabowski PS** : Human osteoblast-like cells produce nitric oxide and express inducible nitric oxide synthase. *Endocrinol*, 135:330-336, 1994.
- 22) **Riancho JA, Salas E, Zarrabeitia T, Olmos JM, Amado JA, Fernades-luna JL and Gonzales-Macias J** : Expression and functional role of nitric oxide synthase in osteoblast-like cells. *J Bone Miner Res*, 10:439-446, 1995.
- 23) **Salter DM, Robb JE and Wright MO** : Electrophysiological responses of human bone cells to mechanical stimulation: evidence for specific integrin function in mechanotransduction. *J Bone Miner Res*, 12:1133-1141, 1997.
- 24) **Stefanovic-Racic M, Stadler J and Evans CH** : Nitric oxide and arthritis. *Arthritis Rheum*, 36:1036-1044, 1993.
- 25) **Toma CD, Ashkar S, Gray ML, Schaffer JL and Gerstenfeld LC** : Signal transduction of mechanical stimuli is dependent on microfilament integrity: Identification of osteopontin as a mechanically induced gene in osteoblasts. *J Bone Miner Res*, 12:1626-1636, 1997.
- 26) **Wimalawansa SJ, De Marco G, Gangula P and Yallampalli C** : Nitric oxide alleviates ovariectomy induced one loss. *Bone*, 18:301-304, 1996.