



# Cohort Profile: Firefighter Research on the Enhancement of Safety and Health (FRESH), a Prospective Cohort Study on Korean Firefighters

Yun Tae Kim<sup>1\*</sup>, Woo Jin Kim<sup>2\*</sup>, Jee Eun Choi<sup>1</sup>, Mun-joo Bae<sup>3</sup>, Heeseon Jang<sup>1</sup>, Chan Joo Lee<sup>4</sup>, Hye-Jeong Lee<sup>5</sup>, Dong Jin Im<sup>5</sup>, Byoung Seok Ye<sup>6</sup>, Mi-Ji Kim<sup>7</sup>, Yeoju Jeong<sup>7</sup>, Sung Soo Oh<sup>8</sup>, Young-Chul Jung<sup>9</sup>, Eun Seok Kang<sup>10</sup>, Sungha Park<sup>4</sup>, Seung Koo Lee<sup>5</sup>, Ki Soo Park<sup>7</sup>, Sang Baek Koh<sup>8</sup>, and Changsoo Kim<sup>2,11</sup>

<sup>1</sup>Department of Public Health, Yonsei University College of Medicine, Seoul;

<sup>2</sup>Department of Preventive Medicine, Yonsei University College of Medicine, Seoul;

<sup>3</sup>Department of Occupational and Environmental Health, Yonsei University Graduate School of Public Health, Seoul;

<sup>4</sup>Division of Cardiology, Yonsei Cardiovascular Hospital, Yonsei University College of Medicine, Seoul;

<sup>5</sup>Department of Radiology, Research Institute of Radiological Science, Yonsei University College of Medicine, Seoul;

<sup>6</sup>Department of Neurology, Yonsei University College of Medicine, Seoul;

<sup>7</sup>Department of Preventive Medicine and Institute of Health Science, Gyeongsang National University College of Medicine, Jinju;

<sup>8</sup>Department of Occupational and Environmental Medicine, Wonju Severance Christian Hospital, Yonsei University Wonju College of Medicine, Wonju;

<sup>9</sup>Department of Psychiatry, Yonsei University College of Medicine, Seoul;

<sup>10</sup>Division of Endocrinology and Metabolism, Department of Internal Medicine, Yonsei University College of Medicine, Seoul;

<sup>11</sup>Institute of Human Complexity and Systems Science, Yonsei University, Songdo, Korea.

Firefighters have a high risk of developing cardiovascular and mental disorders due to their physical and chemical environments. However, in Korea, few studies have been conducted on environmental risk of firefighters. The Firefighter Research on the Enhancement of Safety and Health (FRESH) study aimed to discover the risk factors for cardiovascular disease and mental disorders among firefighters. Former and current firefighters were recruited from three university hospitals. A total of 1022 participants completed baseline health examinations from 2016 to 2017. All participants were scheduled for follow-ups every 2 years. Baseline health survey, laboratory testing of blood and urine samples, blood heavy metal concentration, urine polycyclic aromatic hydrocarbons (PAHs) metabolites, stress-related hormone test, natural killer cell activity, as well as physical and mental health examinations that focused on cardiovascular and mental disorders, were conducted. In addition, 3 Tesla (3T) brain magnetic resonance imaging (MRI) and neuropsychological tests were also performed to investigate structural and functional changes in the brains of 352 firefighters aged >40 years or new hires with less than 1 year of service.

**Key Words:** Firefighters, cohort studies, cardiovascular diseases, mental disorders, magnetic resonance imaging, Republic of Korea

**Received:** September 6, 2019 **Revised:** December 2, 2019

**Accepted:** December 2, 2019

**Corresponding author:** Changsoo Kim, MD, PhD, Department of Preventive Medicine, Yonsei University College of Medicine, 50-1 Yonsei-ro, Seodaemun-gu, Seoul 03722, Korea.

Tel: 82-2-2228-1860, Fax: 82-2-392-8133, E-mail: preman@yuhs.ac

\*Yun Tae Kim and Woo Jin Kim contributed equally to this work.

•The authors have no potential conflicts of interest to disclose.

© Copyright: Yonsei University College of Medicine 2020

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Firefighters are considered to have one of the most dangerous jobs. In the Republic of Korea (ROK), the number of fire service casualties has been reported to reach 2058 over the past 5 years.<sup>1</sup> During fire suppression and first aid rescue, firefighters are exposed to hazardous environments, making them vulnerable to various physical and mental disorders.<sup>2,3</sup> Extremely high temperatures and smoke, which contains gaseous pollutants and particulate toxins, at fire sites increase the risk of cardiovascular disease in firefighters.<sup>4-6</sup> There is also a risk to the mental health of firefighters, as they often encounter injuries or deaths.<sup>7,8</sup>

Several cohort studies have investigated the risk or protective factors for physical and mental disorders among firefighters. Firefighter Obesity Research: Work-place Assessment to Reduce Disease (FORWARD) was a representative large-scale study that aimed to prevent long-term diseases caused by obesity in the United States.<sup>7</sup> Another cohort study investigated cancer risk among approximately 30000 firefighters working in three U.S. cities.<sup>8</sup> Representative large-scale studies have also been conducted in Europe. Using data from the Nordic occupational cancer project conducted in five Nordic countries (Denmark, Finland, Iceland, Sweden, and Norway), 16422 male firefighters were tracked for 45 years to identify their cancer patterns.<sup>9</sup> Moreover, in Finland, a 13-year follow-up study was conducted on the association between musculoskeletal pain and other symptoms in firefighters.<sup>10-12</sup> Most studies have shown that firefighters are at high risk of developing occupational diseases.<sup>13-17</sup>

In Korea, a cohort of 33416 male firefighters, based on data from the Korea National Central Cancer Registry, was built to investigate cancer incidence from 1996 to 2007.<sup>18</sup> A study conducted by the Korean National Fire Agency (NFA) investigated work-related incidents over a 5-year period.<sup>19</sup> Although a number of similar studies have been conducted in the ROK, most of those studies used registry data or National Health Insurance claims data. Cohort studies focusing on the physical and mental health of firefighters, while considering the work and environment of firefighters, have not yet been conducted.<sup>20</sup> The current study's protocols were approved by the institutional review boards of Severance Hospital, Yonsei University Health System in Seoul, Korea (4-2016-0187), Wonju Severance Christian Hospital in Wonju, Korea (CR316014-002), and Gyeongsang National University Hospital in Jinju, Korea (GNUH 2016-04-015-006). In this prospective cohort study, we aimed to identify the risk and protective factors for cardiovascular disease and mental health in firefighters.<sup>4,5</sup>

We recruited firefighters, fire service academy trainees, and retirees from the NFA in Korea. Baseline health examinations were conducted at three university hospitals: Severance Hospital recruited firefighters from Seoul, Gyeonggi, Chungcheongnam-do, Daejeon, and Jeollabuk-do regions; Wonju Severance Christian Hospital recruited firefighters from Gangwon and Chungcheongbuk-do regions; and Gyeongsang National University Hospital recruited firefighters from Gyeongsangbuk-do, Gyeongsangnam-do, and Jeollanam-do regions. A total of 1022 firefighters were recruited from 2016 to 2017. The participants consisted of 100 fire service academy students, 448 fire-control workers, 213 paramedics and rescue workers, 185 office administrators, and 76 retirees. According to recruiting institutions, 401 participants were recruited from Severance Hospital, 307 from Wonju Severance Christian Hospital, and 314 from Gyeongsang National University Hospital. Among these participants, 978 (95.70%) were male and 44 (4.30%) were female. Firefighters who participated in the baseline examina-

tion were followed-up every 2 years. Before starting the research, a standardized protocol was prepared and distributed to all nurses who conducted the examination. They received standardized training on screening schedules, process verification, and examination simulations.

The topics and items in the questionnaires used in FRESH cohort are listed in Table 1. All participants were interviewed by trained nurses. Questionnaires at baseline covered information on anthropometric measures and medical history, and evaluated factors related to the participants' occupational environment, such as work shifts and experience of psychological trauma.

Musculoskeletal symptoms were surveyed using the Musculoskeletal Discomfort questionnaire of the Korean Occupational Safety and Health Agency (KOSHA), which is based on the musculoskeletal symptom criteria established by the National Institute for Occupational Safety and Health in the United States.<sup>21</sup> Ocular surface disease index (OSDI) survey was used to assess dry eye symptoms.<sup>22</sup> Currently, the regular health examination program performed by the NFA includes slit-lamp examination, fundus photography, ocular tonometry, and eyesight test; therefore, considering the redundancy of the examination, we only included xerophthalmia survey.<sup>23</sup>

To measure mental health status, we included questionnaires on post-traumatic stress disorder (PTSD), sleep disorder, depression, and anxiety disorder. PTSD was measured using the PTSD Checklist-Specific (PCL-S).<sup>24,25</sup> Sleep disorders were measured using the Pittsburgh Sleep Quality Index (PSQI), and depression was characterized by a shortened Center for Epidemiologic Studies Depression scale (CES-D).<sup>26,27</sup> The Alcohol Use Disorders Identification Test in Korea (AUDIT-K) was used to evaluate alcohol consumption; finally, the Beck Anxiety Inventory (BAI) was used to measure anxiety disorders as follows: severe anxiety (BAI score  $\geq 32$ ), moderate anxiety (27-31), mild anxiety (22-26), and normal (0-21).<sup>28,29</sup> The following blood indices were measured: biochemical indicators, diagnostic blood test, electrolytes lipid markers, myocardial marker, hepatobiliary function test, renal function test, blood glucose-related test, and high-sensitivity C-reactive protein (hsCRP).

Firefighters are often exposed to various health hazards, especially heavy metals and persistent organic pollutants. The concentrations of two heavy metals (lead, cadmium) and four polycyclic aromatic hydrocarbons (PAHs; 2-naphthol, 2-hydroxyfluorene, 1-hydroxyphenanthrene, 1-hydroxypyrene) were measured in the blood and urine, respectively.

For stress-related hormone measurements, the level of serum cortisol, salivary cortisol, and cortisone were measured, and hair was collected to perform long-term stress hormone analysis. Natural killer (NK) cell activity was measured to examine the relationship between immune function and stress.

Among the 1022 participants, 352 further underwent 3 Tesla (3T) brain magnetic resonance imaging (MRI), as firefighters are exposed to particulate matter and carbon monoxides and

**Table 1.** Elements Evaluated at Baseline in FRESH Specialized Cohort

Classification	Contents	Method
Physical measurements	Height, weight, body mass index, vision, waist circumference, hip circumference, thigh circumference, arm circumference, resting blood pressure, pulse rate	Questionnaire
Health survey		Questionnaire
Demographic characteristics	Age, sex, date of birth (legal and actual), educational level, marriage status, household income, number of work shifts (per month), number of movements (per month) Medical history of stroke, transient ischemic attack, cardiovascular disease, chronic renal disease, hypertension, dyslipidemia, diabetes mellitus, thyroid disease, liver disease, asthma or chronic obstructive pulmonary disease, osteoporosis, arthritis, mental health disease, autoimmune disease, neoplasm Family (confined to immediate family) history of myocardial infarction, hypertension, stroke, diabetes mellitus, mental health disease, neoplasm Current medication and reproductive health information (menarche, pregnancy, gestational diabetes and hypertension, oral contraceptive use, female hormone use)	
Health-related behaviors	Smoking, drinking, sleep duration, obstructive sleep apnea risk, snoring, physical activity	
Mental health questionnaire	Stressful life event, PTSD, sleep disorder, alcohol consumption (AUDIT-K), depression (CES-D), anxiety disorder (BAI)	Questionnaire
Musculoskeletal symptom questionnaires	Musculoskeletal discomfort questionnaires of the Korean Occupational Safety and Health Agency (KOSHA Code H-30-2003) (pain level, frequency and duration of pain in the neck, shoulder, elbow, arm, hand, wrist, waist, leg, foot)	Questionnaire
Biochemical indicators		Laboratory analysis
Diagnostic blood test (blood)	WBC count, WBC differential count, lymphocyte, monocyte, eosinophil, basophil, RBC count, hemoglobin, hematocrit, MCV, MCH, MCHC, platelet count, MPV, PDW, RDW	
Electrolytes (blood)	Ca, Cl, P, K, Na	
Lipid markers (blood)	Total cholesterol, triglyceride, HDL cholesterol, LDL cholesterol	
Myocardial marker (blood)	Hs Troponin-T, NT-proBNP	
Hepatobiliary function test (blood)	Total protein, albumin, total bilirubin, ALP, AST, GGT, ALT	
Renal function test (urine)	BUN, microalbumin/creatinine ratio, microalbumin, creatinine, uric acid	
Other test (blood)	HbA1c, fasting blood glucose, hsCRP	
Environmental pollutants		Laboratory analysis
Blood heavy metal concentration	Cd, Pb	
Urine PAH metabolites	2-OHF, 1-OHPHE, 1-OHP, 2-naphthol	
Stress hormone test	Serum cortisol, salivary cortisol, salivary cortisone, NK cell activity	Laboratory analysis
Cardiovascular examination	Electrocardiogram, cardiopulmonary exercise test, carotid ultrasonography*, echocardiography <sup>†</sup> , PWV test*, ABI test*, cardiac CT, pulmonary CT*, 24-hour Holter monitoring <sup>‡</sup>	Examination
Brain MRI <sup>§</sup>	Resting functional MRI, dDTI axial, faDTI axial, isoDTI axial, 3D T1WI non-contrast coronal, 3D T1WI non-contrast axial, 3D T1WI non-contrast sagittal, T2WI axial, 3D T2 fluid-attenuated inversion-recovery	Examination
Neuropsychological examination (SNSB) <sup>§</sup>	Memory function (SVLT recognition, SVLT free/delayed recalls, RCFT free/delayed recalls), Language and related function (K-BNT), Visuospatial function (RCFT copy), Frontal/Executive function [Stroop test-color reading, Trail making test, COWAT, Category fluency test, COWAT Letter (phonemic) fluency test], Attention (Digit span test)	Examination

FRESH, Firefighter Research on the Enhancement of Safety and Health; PTSD, post-traumatic stress disorder; AUDIT-K, Alcohol Use Disorders Identification Test in Korea; CES-D, Center for Epidemiologic Studies Depression scale; BAI, Beck Anxiety Inventory; WBC, white blood cell; RBC, red blood cell; MCV, mean corpuscular volume; MCH, mean corpuscular hemoglobin; MCHC, mean corpuscular hemoglobin concentration; MPV, mean platelet volume; PDW, platelet distribution width; RDW, red cell distribution width; Ca, Calcium; Cl, chloride; P, phosphorus; K, potassium; Na, sodium; HDL, high-density lipoprotein; LDL, low-density lipoprotein; NT-proBNP, N-terminal-pro-brain natriuretic peptide; ALP, alkaline phosphatase; AST, aminotransferase; GGT,  $\gamma$ -glutamyl transferase; ALT, alanine aminotransferase; BUN, blood urea nitrogen; hsCRP, high-sensitivity C-reactive protein; Cd, Cadmium; Pb, lead; PAH, polycyclic aromatic hydrocarbon; 2-OHF, 2-hydroxyfluorene; 1-OHPHE, 1-hydroxyphenanthrene; 1-OHP, 1-hydroxypyrene; NK cell, natural killer cell; PWV, pulse wave velocity; ABI, ankle brachial index; SNSB, Seoul Neuropsychological Screening Battery; SVLT, Seoul Verbal Learning Test; RCFT, Ray Complex Figure Test; K-BNT, Korean-Boston Naming Test; COWAT, Controlled Oral Word Association Test.

\*Conducted at Severance Hospital and Gyeongsang National University Hospital; <sup>†</sup>Conducted at Severance Hospital; <sup>‡</sup>Applicants Only; <sup>§</sup>Approximately 350 participants aged >40 years or fire service academy students, applicants only.

are at risk of mild traumatizing brain injury; these could result in structural and functional abnormalities in the brain. The brain MRI protocols included the following: dTI axial, faDTI axial, isoDTI axial, 3D T1WI non-contrast coronal, 3D T1WI non-contrast axial, 3D T1WI non-contrast sagittal, T2WI axial, resting functional MRI, and 3D T2 fluid-attenuated inversion-recovery. All three hospitals used the same brain MRI protocols. Brain volume and cortical thickness were estimated using Inbrain® software (<https://www.inbrain.co.kr/>).<sup>30-33</sup> While brain MRI can identify structural changes, neuropsychological examinations are required to identify functional abnormalities. Neuropsychological examinations not only indicate abnormalities in the functional areas of the brain, but also compare them to the average level. Participants undergoing brain MRI underwent neuropsychological testing using the Seoul Neuropsychological Screening Battery (SNSB).<sup>34</sup>

Baseline characteristics are shown in Tables 2 and 3. Distribution and statistical significance of the variables by duty were verified, while assuming that work environment could vary greatly depending on the duty, and that the duty would have a significant effect on cardiovascular and mental health. Systolic blood pressure was the lowest in paramedics and res-

cue workers, and the highest in fire-control workers ( $p<0.001$ ). The proportion of participants with high-risk PTSD was 111 (10.86%). Among firefighters, the proportion of participants with high-risk PTSD (14.08%) was the highest among paramedics and rescue workers. The risk of severe depression, according to CES-D test, was the highest in retiree group. Severe anxiety was identified in eight participants (0.78%). PSQI tests showed that 486 out of 1022 participants had normal sleep quality (47.55%), whereas 536 had poor sleep quality (52.45%). SNSB test was performed in 350 participants, which included 16 fire service academy students, 177 fire-control workers, 62 paramedics and rescue workers, 59 office administrators, and 36 retirees.

Of the 352 brain MRI tests, 255 showed negative findings in brain. Thirteen and 11 subjects showed old infarct lesions and small vessel disease, respectively, while no participant showed acute cerebral infarction or hemorrhagic lesions. Twelve participants with neoplastic lesions had benign tumors, and no malignant tumor was observed. In addition, 60, two, and four subjects showed nonspecific blight object, atrophic changes, and normal variants (mastoid effusion, cavum septum pellucidum, cavum vergae), respectively.

**Table 2.** General Characteristics of Firefighters in FRESH Cohort at Baseline

	Total (n=1022)	School educators (n=100)	Fire-control workers (n=448)	Paramedics and rescue workers (n=213)	Office administrators (n=185)	Retirees (n=76)	p value
Age (yr)	41.77±10.59	29.46±4.93	42.20±9.44	39.10±8.34	43.17±9.49	59.62±2.18	<0.001
Sex							<0.001
Male	978 (95.70)	96 (96.00)	443 (98.88)	190 (89.20)	174 (94.05)	76 (100.00)	
Female	44 (4.30)	4 (4.00)	5 (1.12)	23 (10.80)	11 (5.95)	0 (0.00)	
Height (cm)	172.64±5.83	174.64±6.12	172.60±7.22	172.57±6.77	172.22±6.04	170.09±4.36	<0.001
Weight (kg)	74.04±9.63	74.16±8.69	74.78±10.32	73.79±10.19	74.30±10.60	71.07±8.79	0.059
Body mass index (kg/m <sup>2</sup> )	24.83±2.65	24.32±2.50	24.98±2.62	24.60±3.03	25.02±2.95	24.56±2.59	0.571
Waist circumference (cm)	86.08±7.40	83.93±6.89	88.08±39.61	85.30±7.07	87.04±7.85	87.67±7.18	0.011
Hip circumference (cm)	97.93±5.76	97.81±4.54	98.11±5.01	97.80±7.93	98.57±5.29	95.83±4.76	<0.001
Thigh circumference (cm)	30.28±3.01	50.46±3.98	50.47±4.16	51.08±4.50	50.37±4.92	46.90±4.25	<0.001
Arm circumference (cm)	50.31±4.47	30.18±2.54	30.00±3.73	31.10±3.44	30.46±3.15	29.35±3.08	0.022
Systolic blood pressure (mm Hg)	127.13±12.88	127.00±9.88	128.33±12.12	124.86±12.14	126.86±14.33	127.28±12.82	<0.001
Diastolic blood pressure (mm Hg)	80.31±9.69	77.23±8.42	81.49±9.95	78.15±9.57	80.77±9.54	82.28±8.52	0.341
Triglyceride (mg/dL)	139.73±87.21	115.46±74.83	141.08±87.74	137.76±86.05	148.81±94.15	147.05±80.70	0.032
Fasting blood sugar (mg/dL)	90.13±13.81	87.16±12.63	89.74±14.27	88.03±10.90	91.82±14.31	98.08±15.41	<0.001
HbA1c (mg/dL)	5.49±0.53	5.30±0.24	5.48±0.57	5.43±0.41	5.53±0.54	5.90±0.55	<0.001
Cadmium (mg/dL)*	0.70 (0.04–4.75)	0.56 (0.13–1.62)	0.65 (0.04–4.75)	0.74 (0.11–2.33)	0.77 (0.12–2.2)	0.89 (0.28–2.41)	<0.001
Lead (mg/dL)*	1.79 (0.41–7.00)	1.48 (0.67–4.27)	1.77 (0.41–5.97)	1.75 (0.71–5.85)	1.89 (0.65–3.84)	2.41 (0.86–7)	<0.001
2-naphthol (µg/L) <sup>†</sup>	2.66 (0–74.16)	3.41 (0–57.6)	2.69 (0–53.76)	2.93 (0–74.16)	2.63 (0–50.62)	2 (0–57.19)	<0.001
2-hydroxyfluorene (µg/L) <sup>†</sup>	0.19 (0–5.67)	0.2 (0.06–2.75)	0.19 (0–2.13)	0.18 (0–5.67)	0.18 (0–3.45)	0.18 (0.02–1.88)	<0.001
1-hydroxyphenanthrene (µg/L) <sup>†</sup>	0.19 (0–3.63)	0.24 (0.03–1.77)	0.2 (0–2.7)	0.17 (0.02–3.63)	0.17 (0–3.16)	0.22 (0.08–0.59)	<0.001
1-hydroxypyrene (µg/L) <sup>†</sup>	0.2 (0–5.81)	0.2 (0–2.45)	0.2 (0.02–5.81)	0.19 (0.03–2.37)	0.2 (0.02–1.72)	0.2 (0.07–2.94)	<0.001

FRESH, Firefighter Research on the Enhancement of Safety and Health.

Values are presented as mean±standard deviation or n (%) unless otherwise indicated.

\*Kruskal-Wallis test was conducted; median (min-max) values are suggested instead of mean±standard deviation; <sup>†</sup>Measured as urinary concentration; median (min-max) values are suggested instead of mean±standard deviation.

**Table 3.** Mental Health Status of Firefighters in FRESH Cohort at Baseline

	Total (n=1022)	School educators (n=100)	Fire-control workers (n=448)	Paramedics and rescue workers (n=213)	Office administrators (n=185)	Retirees (n=76)	p value
PTSD							<0.001
High risk (<11)	111 (10.86)	6 (6.00)	49 (10.94)	30 (14.08)	17 (9.19)	9 (11.84)	
Low risk (≥11)	911 (89.14)	94 (94.00)	399 (89.06)	183 (85.92)	168 (90.81)	67 (88.16)	
CES-D							0.188
Severe depression (≥23)	19 (1.86)	1 (1.00)	8 (1.79)	4 (1.88)	3 (1.62)	3 (3.95)	
Moderate depression (19–22)	18 (1.76)	4 (4.00)	7 (1.56)	3 (1.41)	3 (1.62)	1 (1.32)	
Mild depression (14–18)	35 (3.42)	2 (2.00)	17 (3.79)	7 (3.29)	8 (4.32)	1 (1.32)	
Normal (0–13)	950 (92.96)	93 (93.00)	416 (92.86)	199 (93.42)	171 (92.44)	71 (93.41)	
AUDIT-K							<0.001
High alcohol dependence (≥26)	21 (2.05)	4 (4.00)	7 (1.56)	4 (1.88)	5 (2.70)	1 (1.32)	
Probable alcohol dependence (15–25)	44 (4.31)	0 (0.00)	23 (5.13)	9 (4.23)	8 (4.32)	4 (5.26)	
Hazardous drinking (12–14)	257 (25.15)	26 (26.00)	118 (26.34)	42 (19.72)	52 (28.11)	19 (25.00)	
Normal (0–11)	700 (68.49)	70 (70.00)	300 (66.97)	158 (74.17)	120 (64.87)	52 (68.42)	
BAI							0.243
Severe anxiety disorder (≥32)	8 (0.78)	0 (0.00)	3 (0.67)	1 (0.47)	2 (1.08)	2 (2.63)	
Moderate anxiety disorder (27–31)	11 (1.08)	1 (1.00)	4 (0.89)	1 (0.47)	3 (1.62)	2 (2.63)	
Mild anxiety disorder (22–26)	20 (1.96)	4 (4.00)	7 (1.56)	3 (1.41)	4 (2.16)	2 (2.63)	
Normal (0–21)	983 (96.18)	95 (95.00)	434 (96.88)	208 (97.65)	176 (95.14)	70 (92.11)	
PSQI							0.002
Poor sleep quality (≥6)	536 (52.45)	50 (50.00)	241 (53.79)	113 (53.05)	104 (56.22)	28 (36.84)	
Normal (<6)	486 (47.55)	50 (50.00)	207 (46.21)	100 (46.95)	81 (43.78)	48 (63.16)	

FRESH, Firefighter Research on the Enhancement of Safety and Health; PTSD, post-traumatic stress disorder; CES-D, Center for Epidemiologic Studies Depression scale; AUDIT-K, Alcohol Use Disorders Identification Test in Korea; BAI, Beck Anxiety Inventory; PSQI, Pittsburgh Sleep Quality Index.

FRESH cohort study was able to identify the risk and protective factors for cardiovascular disease and mental disorders in more than 1000 Korean firefighters. In order to consider the exposure variations in occupational environments, we recruited firefighters from three different regions in Korea. In addition to different variables, exposure to heavy metals and PAHs can also be used to confirm the association of occupational exposure with cardiovascular and mental disorders. Although firefighters are part of an occupational group that is vulnerable to a wide variety of mental health hazards and toxic substances, previous studies have only examined the association between mental disorders, such as anxiety disorders and PTSD, and the results of questionnaire surveys.<sup>23,24</sup> Since 3T brain MRI and neuropsychological tests were performed in this study, it was possible to identify both functional and structural changes in the brain.

FRESH cohort offered several advantages. This was the first study to investigate cardiovascular disease and cognitive function of firefighters in Korea. While previous cohort studies on firefighters were mainly retrospective studies using secondary data, this study examined factors that affect cardiovascular disease, mental disease, and cognitive dysfunction using various clinical tests as a prospective study. In addition, inspections were conducted for firefighters nationwide using standardized screening protocols. In the study design process, standard-

ization protocols were prepared and distributed to all researchers, and standardization training was provided to the personnel who conducted the examinations to minimize measurement errors at different institutions.

Nevertheless, our research also had some limitations. First of all, selection bias may have resulted from only including subjects who applied for the study, rather than randomly selecting the participants. For example, people who are usually interested in health-related issues may have been more likely to participate in this study, which could lead to the healthy worker effect. In addition, some of the items that should be investigated specifically for firefighters were not included in our questionnaire. For example, years of service and recent exposure to trauma can be important variables in analyzing the relationship between disease and work characteristics. To compensate for this, starting in 2018, when baseline participants began their first follow-up, we added questions about the date of entry, date of retirement (if leaving), and trauma experienced within the last 6 months.

FRESH study is available to those who are interested in collaboration. FRESH research committee requests a short research proposal, including background information, research questions, methods, and authorship. FRESH research committee is responsible for the distribution and control of data. Researchers who are interested in collaborative research can con-

tact the FRESH investigator at freshcohort@gmail.com.

## ACKNOWLEDGEMENTS

This research was supported by the Fire Fighting Safety & 119 Rescue Technology Research and Development Program funded by the National Fire Agency (“MPSS-Fire Safety-2015-80”). The funders had no role in the study design, data collection, analysis, decision to publish, or preparation of this manuscript.

## AUTHOR CONTRIBUTIONS

**Conceptualization:** Yun Tae Kim, Woo Jin Kim, Jee Eun Choi, Mun-joo Bae, Heeseon Jang, Sungha Park, Chan Joo Lee, Seung Koo Lee, Hye-Jeong Lee, Dong Jin Im, Eun Seok Kang, Young-Chul Jung, Byoung Seok Ye, Sang Baek Koh, Sung Soo Oh, Ki Soo Park, Mi-Ji Kim, and Changsoo Kim. **Data curation:** Yun Tae Kim, Woo Jin Kim, Jee Eun Choi, Heeseon Jang, Sung Soo Oh, Mi-Ji Kim, Yeouju Jeong. **Formal analysis:** Yun Tae Kim and Woo Jin Kim. **Funding acquisition:** Ki Soo Park, Sang Baek Koh, and Changsoo Kim. **Investigation:** Woo Jin Kim, Sung Soo Oh, and Yeouju Jeong. **Methodology:** Yun Tae Kim and Woo Jin Kim. **Project administration:** Ki Soo Park, Sang Baek Koh, and Changsoo Kim. **Resource:** Ki Soo Park, Sang Baek Koh, and Changsoo Kim. **Software:** Yun Tae Kim. **Supervision:** Ki Soo Park, Sang Baek Koh, and Changsoo Kim. **Validation:** Woo Jin Kim. **Visualization:** Yun Tae Kim. **Writing—original draft:** Yun Tae Kim and Woo Jin Kim. **Writing—review & editing:** Yun Tae Kim, Woo Jin Kim, and Changsoo Kim.

## ORCID iDs

Yun Tae Kim	<a href="https://orcid.org/0000-0001-5171-2962">https://orcid.org/0000-0001-5171-2962</a>
Woo Jin Kim	<a href="https://orcid.org/0000-0001-5520-4228">https://orcid.org/0000-0001-5520-4228</a>
Jee Eun Choi	<a href="https://orcid.org/0000-0001-8290-9522">https://orcid.org/0000-0001-8290-9522</a>
Mun-joo Bae	<a href="https://orcid.org/0000-0002-0242-9826">https://orcid.org/0000-0002-0242-9826</a>
Heeseon Jang	<a href="https://orcid.org/0000-0002-9737-7220">https://orcid.org/0000-0002-9737-7220</a>
Chan Joo Lee	<a href="https://orcid.org/0000-0002-8756-409X">https://orcid.org/0000-0002-8756-409X</a>
Hye-Jeong Lee	<a href="https://orcid.org/0000-0003-4349-9174">https://orcid.org/0000-0003-4349-9174</a>
Dong Jin Im	<a href="https://orcid.org/0000-0001-8139-5646">https://orcid.org/0000-0001-8139-5646</a>
Byoung Seok Ye	<a href="https://orcid.org/0000-0003-0187-8440">https://orcid.org/0000-0003-0187-8440</a>
Mi-Ji Kim	<a href="https://orcid.org/0000-0002-8646-832X">https://orcid.org/0000-0002-8646-832X</a>
Yeouju Jeong	<a href="https://orcid.org/0000-0002-6926-1124">https://orcid.org/0000-0002-6926-1124</a>
Sung Soo Oh	<a href="https://orcid.org/0000-0003-0801-3052">https://orcid.org/0000-0003-0801-3052</a>
Young-Chul Jung	<a href="https://orcid.org/0000-0002-0578-2510">https://orcid.org/0000-0002-0578-2510</a>
Eun Seok Kang	<a href="https://orcid.org/0000-0002-0364-4675">https://orcid.org/0000-0002-0364-4675</a>
Sungha Park	<a href="https://orcid.org/0000-0001-5362-478X">https://orcid.org/0000-0001-5362-478X</a>
Seung Koo Lee	<a href="https://orcid.org/0000-0001-5646-4072">https://orcid.org/0000-0001-5646-4072</a>
Ki Soo Park	<a href="https://orcid.org/0000-0001-5571-3639">https://orcid.org/0000-0001-5571-3639</a>
Sang Baek Koh	<a href="https://orcid.org/0000-0001-5609-6521">https://orcid.org/0000-0001-5609-6521</a>
Changsoo Kim	<a href="https://orcid.org/0000-0002-5940-5649">https://orcid.org/0000-0002-5940-5649</a>

## REFERENCES

- National Fire Agency (NFA). Statistical Yearbook (2018) [Internet]. Sejong: NFA; c2018 [accessed on 2019 August 22]. Available at: [https://www.nfa.go.kr/nfa/releaseinformation/statisticalinformation/main/?boardId=bbs\\_000000000000019&mode=view&cntId=16&category=&pageIdx=](https://www.nfa.go.kr/nfa/releaseinformation/statisticalinformation/main/?boardId=bbs_000000000000019&mode=view&cntId=16&category=&pageIdx=)
- Brandt-Rauf PW, Fallon LF, Tarantini T, Idema C, Andrews L. Health hazards of fire fighters: exposure assessment. *J Occup Environ Med* 1988;45:606-12.

- Kim JM, Lee HJ. Hazards exposed to firefighters in fire-physical, chemical, and biologic factors. *J Korean Med Assoc* 2008;51:1072-7.
- Kales SN, Soteriades ES, Christophi CA, Christiani DC. Emergency duties and deaths from heart disease among firefighters in the United States. *N Engl J Med* 2007;356:1207-15.
- Soteriades ES, Smith DL, Tsismenakis AJ, Baur DM, Kales SN. Cardiovascular disease in US firefighters: a systematic review. *Cardiol Rev* 2011;19:202-15.
- Li K, Lipsey T, Leach HJ, Nelson TL. Cardiac health and fitness of Colorado male/female firefighters. *Occup Med (Lond)* 2017;67:268-73.
- Dobson M, Choi B, Schnall PL, Wigger E, Garcia-Rivas J, Israel L, et al. Exploring occupational and health behavioral causes of firefighter obesity: a qualitative study. *Am J Ind Med* 2013;56:776-90.
- Daniels RD, Kubale TL, Yiin JH, Dahm MM, Hales TR, Baris D, et al. Mortality and cancer incidence in a pooled cohort of US firefighters from San Francisco, Chicago and Philadelphia (1950-2009). *Occup Environ Med* 2014;71:388-97.
- Pukkala E, Martinsen JI, Weiderpass E, Kjaerheim K, Lynge E, Trygvadottir L, et al. Cancer incidence among firefighters: 45 years of follow-up in five Nordic countries. *Occup Environ Med* 2014;71:398-404.
- Airila A, Hakanen JJ, Luukkonen R, Lusa S, Punakallio A, Leino-Arjas P. Developmental trajectories of multisite musculoskeletal pain and depressive symptoms—the effects of job demands and resources and individual factors. *Psychol Health* 2014;29:1421-41
- Lusa S, Miranda H, Luukkonen R, Punakallio A. Sleep disturbances predict long-term changes in low back pain among Finnish firefighters: 13-year follow-up study. *Int Arch Occup Environ Health* 2015;88:369-79.
- Punakallio A, Lusa S, Luukkonen R, Airila A, Leino-Arjas P. Musculoskeletal pain and depressive symptoms as predictors of trajectories in work ability among Finnish firefighters at 13-year follow-up. *J Occup Environ Med* 2014;56:367-75.
- Lourel M, Abdellaoui S, Chevalyre S, Paltrier M, Gana K. Relationships between psychological job demands, job control and burnout among firefighters. *N Am J Psychol* 2008;10:489-96.
- Ângelo RP, Chambel MJ. The reciprocal relationship between work characteristics and employee burnout and engagement: a longitudinal study of firefighters. *Stress Health* 2015;31:106-14.
- Morren M, Yzermans CJ, van Nispen RM, Wevers SJ. The health of volunteer firefighters three years after a technological disaster. *J Occup Health* 2005;47:523-32.
- Huynh JY, Xanthopoulou D, Winefield AH. Social support moderates the impact of demands on burnout and organizational connectedness: a two-wave study of volunteer firefighters. *J Occup Health Psychol* 2013;18:9-15.
- Schermer TR, Malbon W, Adams R, Morgan M, Smith M, Crockett AJ. Change in lung function over time in male metropolitan firefighters and general population controls: a 3-year follow-up study. *J Occup Health* 2013;55:267-75.
- Ahn YS, Jeong KS, Kim KS. Cancer morbidity of professional emergency responders in Korea. *Am J Ind Med* 2012;55:768-78.
- Kim HD, An YS, Kim DH, Jeong KS, Ahn YS. An overview of compensated work-related injuries among Korean firefighters from 2010 to 2015. *Ann Occup Environ Med* 2018;30:57.
- Kim YK, Ahn YS, Kim K, Yoon JH, Roh J. Association between job stress and occupational injuries among Korean firefighters: a nationwide cross-sectional study. *BMJ Open* 2016;6:e012002.
- Lee DK, Kim JH. A study on the prevention system of musculoskeletal disorders in Korea and other countries. *J Ergon Soc Korea* 2010;29:423-33.

22. Schiffman RM, Christianson MD, Jacobsen G, Hirsch JD, Reis BL. Reliability and validity of the ocular surface disease index. *Arch Ophthalmol* 2000;118:615-21.
23. National Fire Agency (NFA). Regulations for the health and safety management of fire officials national safety and control ordinance. No. 1 Annex 4 (2015) [Internet]. Sejong: NFA; c2015 [accessed on 2019 August 22]. Available at: <http://law.go.kr/LSW/admRulLsInfoP.do?admRulSeq=2100000013941>.
24. Wilkins KC, Lang AJ, Norman SB. Synthesis of the psychometric properties of the PTSD checklist (PCL) military, civilian, and specific versions. *Depress Anxiety* 2011;28:596-606.
25. Blanchard EB, Jones-Alexander J, Buckley TC, Forneris CA. Psychometric properties of the PTSD Checklist (PCL). *Behav Res Ther* 1996;34:669-73.
26. Buysse DJ, Reynolds CF 3rd, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res* 1989;28:193-213.
27. Radloff LS. The CES-D scale: a self-report depression scale for research in the general population. *Appl Psychol Meas* 1977;1:385-401.
28. Beck AT, Epstein N, Brown G, Steer RA. An inventory for measuring clinical anxiety: psychometric properties. *J Consult Clin Psychol* 1988;56:893-7.
29. Saunders JB, Aasland OG, Babor TF, de la Fuente JR, Grant M. Development of the alcohol use disorders identification test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption--II. *Addiction* 1993;88:791-804.
30. Lee JS, Kim C, Shin JH, Cho H, Shin DS, Kim N, et al. Machine learning-based individual assessment of cortical atrophy pattern in Alzheimer's disease spectrum: development of the classifier and longitudinal evaluation. *Sci Rep* 2018;8:4161.
31. Zhu J, Jin Y, Wang K, Zhou Y, Feng Y, Yu M, et al. Frequency-dependent changes in the regional amplitude and synchronization of resting-state functional MRI in stroke. *PLoS One* 2015;10:e0123850.
32. Zou QH, Zhu CZ, Yang Y, Zuo XN, Long XY, Cao QJ, et al. An improved approach to detection of amplitude of low-frequency fluctuation (ALFF) for resting-state fMRI: fractional ALFF. *J Neurosci Methods* 2008;172:137-41.
33. Zuo XN, Di Martino A, Kelly C, Shehzad ZE, Gee DG, Klein DF, et al. The oscillating brain: complex and reliable. *Neuroimage* 2010;49:1432-45.
34. Jahng SM, Na DL, Kang YW. Constructing a composite score for the Seoul neuropsychological screening battery-core. *Dement Neurocogn Disord* 2015;14:137-42.