



저작자표시-비영리-변경금지 2.0 대한민국

이용자는 아래의 조건을 따르는 경우에 한하여 자유롭게

- 이 저작물을 복제, 배포, 전송, 전시, 공연 및 방송할 수 있습니다.

다음과 같은 조건을 따라야 합니다:



저작자표시. 귀하는 원저작자를 표시하여야 합니다.



비영리. 귀하는 이 저작물을 영리 목적으로 이용할 수 없습니다.



변경금지. 귀하는 이 저작물을 개작, 변형 또는 가공할 수 없습니다.

- 귀하는, 이 저작물의 재이용이나 배포의 경우, 이 저작물에 적용된 이용허락조건을 명확하게 나타내어야 합니다.
- 저작권자로부터 별도의 허가를 받으면 이러한 조건들은 적용되지 않습니다.

저작권법에 따른 이용자의 권리는 위의 내용에 의하여 영향을 받지 않습니다.

이것은 [이용허락규약\(Legal Code\)](#)을 이해하기 쉽게 요약한 것입니다.

[Disclaimer](#)

Effect of pregabalin administration
upon reperfusion in a rat model of
hyperglycemic stroke:
mechanistic insights associated with
High-mobility group box 1

Young Song

Department of Medicine

The Graduate School, Yonsei University

Effect of pregabalin administration
upon reperfusion in a rat model of
hyperglycemic stroke:
mechanistic insights associated with
High-mobility group box 1

Directed by Professor Jae-Kwang Shim

The Doctoral Dissertation
submitted to the Department of Medicine,
the Graduate School of Yonsei University
in partial fulfillment of the requirements
for the degree of Doctor of Philosophy

Young Song

June 2017

This certifies that the Doctoral Dissertation
of Young Song is approved.

Thesis Supervisor: Jae-Kwang Shim

Thesis Committee Member #1: Young Lan Kwak

Thesis Committee Member #2: Jeon-Soo Shin

Thesis Committee Member #3: Sak Lee

Thesis Committee Member #4: Sung Yong Park

The Graduate School
Yonsei University

June 2017

ACKNOWLEDEMENTS

First of all, I would like to express the deepest appreciation to my thesis supervisor, Professor Jae-Kwang Shim, who has encouraged me throughout the whole process. This dissertation would not have been possible without his guidance and persistent help.

I would like to thank my committee members, Professor Young Lan Kwak, Jeon-Soo Shin, Sak Lee, and Sung Yong Park, whose expert, sincere and valuable advice and guidance truly enriched this doctoral thesis. I also wish to thank Ji Hae Jun and Eun Jung Shin, who have lent their helping hand in this venture.

I take this opportunity to record my sincere thanks to all of my colleagues at the Department of Anesthesia and Pain Medicine for their help and encouragement.

Finally, I dedicate this work to my loving family – their unconditional love and constant support is what made all of this possible.

Young Song

< TABLE OF CONTENTS >

ABSTRACT.....	1
I. INTRODUCTION.....	3
II. MATERIALS AND METHODS	
1. Animals preparation.....	5
2. MCAO models and study groups.....	5
3. Neurological deficits.....	6
3. Infarct size.....	7
4. Apoptosis.....	7
5. Immunoblot analysis.....	8
6. Statistical analysis.....	8
III. RESULTS	
1. Functional and histological neuroprotection.....	9
2. HMGB1/TLR-4/p-NF- κ B and pro-inflammatory cytokines.....	12
3. Nitric oxide synthases activity.....	14
4. Bcl-2, Bax, cytochrome C, and caspase-3 activity.....	15
IV. DISCUSSION.....	17
V. CONCLUSION.....	22
REFERENCES.....	23
ABSTRACT (IN KOREAN)	27

LIST OF FIGURES

Figure 1. Determination of the dose range for pregabalin treatment	10
Figure 2. Effect of pregabalin treatment upon reperfusion on apoptosis in the brain of hyperglycemic stroke rat	11
Figure 3. Pregabalin treatment attenuated expressions of HMGB1/TLR-4/p-NF- κ B/IL-1 β and TNF- α in the brain of hyperglycemic stroke rat	13
Figure 4. Pregabalin treatment favorably modulated expressions of NOS in the brain of hyperglycemic stroke rat.....	14
Figure 5. Pregabalin treatment favorably modulated expressions of apoptosis-related molecules in the brain of hyperglycemic stroke rat	16

ABSTRACT

Effect of pregabalin administration upon reperfusion in a rat model of hyperglycemic stroke: mechanistic insights associated with HMGB1

Young Song

*Department of Medicine
The Graduate School, Yonsei University*

(Directed by Professor Jae-Kwang Shim)

Hyperglycemia, which reduces the efficacy of treatments and worsens clinical outcomes, is common in stroke. Ability of pregabalin to reduce neuroexcitotoxicity may provide protection against stroke, even under hyperglycemia. We investigated its protective effect against hyperglycemic stroke and its possible molecular mechanisms. Male Wistar rats administered dextrose to cause hyperglycemia, underwent middle cerebral artery occlusion for 1 h and subsequent reperfusion. Rats were treated with an intraperitoneal injection of 30 mg/kg pregabalin or an equal amount of normal saline at the onset of reperfusion (n = 16 per group). At 24 h after reperfusion, neurological deficit, infarct volume, and apoptotic cell count were assessed. Western blot analysis was performed to determine protein expression of high-mobility group

box 1 (HMGB1), toll-like receptor-4 (TLR-4), phosphorylated nuclear factor-kappa B (p-NF- κ B), interleukin-1beta (IL-1 β), tumor necrosis factor-alpha (TNF- α), phosphorylated inducible and endothelial nitric oxide synthase (p-iNOS, p-eNOS), Bcl-2, Bax, Cytochrome C, and caspase-3 in the brain. Pregabalin-treated rats showed significantly improved neurological function (31% decrease in score), reduced infarct size (by 33%), fewer apoptotic cells (by 63%), and lower expression levels of HMGB1, TLR4, p-NF- κ B, IL-1 β , and TNF- α , compared with control rats. Decreased p-iNOS and increased p-eNOS expressions were also observed. Expression of Bax, Cytochrome C, and cleaved caspase-3/caspase3 was significantly downregulated, while Bcl-2 expression was increased by pregabalin treatment. Pregabalin administration upon reperfusion decreased neuronal death and improved neurological function in hyperglycemic stroke rats. Cogent mechanisms would include attenuation of HMGB1/TLR-4-mediated inflammation and favorable modulation of the NOS.

Key words: stroke, hyperglycemia, pregabalin, high-mobility group box 1, nitric oxide synthase

Effect of pregabalin administration up reperfusion in a rat model of hyperglycemic stroke: mechanistic insights associated with HMGB1

Young Song

*Department of Medicine
The Graduate School, Yonsei University*

(Directed by Professor Jae-Kwang Shim)

I. INTRODUCTION

Irrespective of a history of diabetes, approximately 30–40% of patients that present with acute ischemic stroke exhibit hyperglycemia, which is known to exacerbate clinical outcomes.¹ Unfortunately, the application of intensive glycemic control does not improve outcomes leaving clinicians with an additional burden, whilst already being confronted with limited therapeutic options against stroke in general.^{2, 3}

The adverse influence of acute hyperglycemia has also been confirmed in animal models of middle cerebral artery occlusion (MCAO).⁴ After energy depletion, ischemic injury universally starts with presynaptic neuronal discharge leading to activation of voltage-gated calcium channels (VGCC) and release of excitatory neurotransmitters in the ischemic core.⁵

This excitotoxicity is followed by delayed inflammatory responses in the penumbra, with high-mobility group box 1 (HMGB1) recently identified as the key pro-inflammatory molecule linking these two successive events.⁶ In the context of acute hyperglycemia, accumulating evidence suggests that

intensification of these pathologic processes leads to increased cerebral injury.

⁷⁻⁹ In addition, hyperglycemia has also been shown to abolish the experimentally proven protective effects of certain agents, such as volatile anesthetic, against cerebral ischemia-reperfusion (I-R) injury.^{10,11}

The excitotoxicity persists for hours, even after reperfusion, providing an estimated therapeutic window of up to 10–12 hours.¹² Therefore, we hypothesized that therapies aimed at this initial event would successfully ameliorate its downstream complex biochemical events leading to neuronal loss, and retain their protective effects against cerebral I-R injury even in acute hyperglycemic condition.

Pregabalin, a widely used drug for neuropathic pain, robustly binds to the $\alpha 2\text{-}\delta$ subunit of the VGCC reducing Ca^{2+} influx and release of excitotoxic neurotransmitters at presynaptic nerve endings.¹³ Pregabalin's neuroprotective effect has been evaluated in terms of spinal cord injury¹⁴ and cerebral I-R injury induced by deep hypothermic circulatory arrest¹⁵ or normoglycemic MCAO¹⁶, providing promising results. However, evidence regarding its neuroprotective effects and related mechanisms against stroke is lacking in the context of hyperglycemia, which deserves a high priority considering its prevalence and clinical impact on the outcome.

Therefore, the aim of this present study was to investigate the neuroprotective effects of pregabalin in a rat model of hyperglycemic stroke and its related key molecular mechanisms associated with HMGB1.

II. MATERIALS AND METHODS

1. Animal preparation

All animal procedures were approved by the committee for the Care and Use of Laboratory Animals, Yonsei University College of Medicine, and were performed in accordance with the Guide for the Care and Use of Laboratory Animals published by the US National Institutes of Health. Rats were fasted except for water for 8 h before surgery, and allowed free access to food and water after surgery. All rats received dextrose (1.2 g/kg) 1 h before MCAO via the tail vein. A blood glucose concentration >11.1 mmol/L was considered as hyperglycemia.¹⁷ The blood glucose concentration was determined at baseline, before MCAO, upon reperfusion, and 24 h thereafter.

2. MCAO models and study groups

Male Wistar rats (8–10 wk old) weighing 270–300 g were anesthetised with xylazine (Rompun, Vial Korea, 10 mg/kg) and tiletamine/zolazepam (Zoletil 50, Virbac Korea, 30 mg/kg). To minimize potential suffering from the procedure, supplemental analgesia with local lidocaine infiltration was provided if there was sudden movement or changes in vital sign of animals. The tail artery was cannulated to monitor mean arterial pressure (MAP) and collect blood. The heart rate (HR) was monitored by subcutaneous stainless steel electrodes connected to the power lab system (ML845 PowerLab with ML132; AD

Instruments, Colorado Springs, CO). The body temperature was continuously monitored and maintained around 37°C using a heating pad.

The experimental MCAO model was generated as previously described by Longa et al.¹⁸ Briefly, the left common carotid artery (CCA), external carotid artery (ECA), and internal carotid artery (ICA) were exposed through a midline neck incision. The ECA was dissected further distally and coagulated along with the terminal lingual and maxillary artery branches, which were then divided. A 4-0 monofilament suture (Doccol Corporation, MA, USA) with silicone rubber coated-rounded head was inserted through the proximal ECA stump into the ICA. After 60 minutes of occlusion, the suture was withdrawn, and the skin was sutured with an infiltration of 0.5% bupivacaine for analgesia. For sham operations, all procedures were identical except that the suture was not inserted. The rats were randomly assigned to three groups: 1) Hyperglycemia-Sham group (n = 8), normal saline 2 ml intraperitoneally (IP) after 60 min of ICA exposure without MCAO; 2) Hyperglycemia-untreated MCAO control group (n = 16), normal saline 2 ml IP after 60 min of MCAO, followed by reperfusion, 3) Hyperglycemia-pregabalin treated MCAO group (n = 16), pregabalin 30 mg/kg in normal saline 2 ml IP after 60 min of MCAO, followed by reperfusion. The dose of pregabalin was selected from previous studies of spinal cord injury and deep hypothermic circulatory arrest,^{14,15} and also from our preliminary study, in which rats treated with 30 mg/kg demonstrated a smaller infarct size, better neurological function, and lower apoptotic cell count than those treated with 10

mg/kg (Fig. 1). The comparative dose of 10 mg/kg was selected from the previous MCAO study in mice under normoglycemia.¹⁶ The dose of 30 mg/kg was chosen for all subsequent experiments. Physical condition of all experimental animals were monitored throughout the procedure and then, every 12 h until the euthanasia was performed with IP injection of an overdose (60mg/kg) of sodium pentobarbital. None of the animals became ill or die prior to the experimental endpoint.

3. Neurological deficits

Neurobehavioral assessment was performed assessing four different functions (general status, simple motor deficit, complex motor deficit, and sensory deficit) by combining elements from several previously described scoring systems.¹⁹ The score given in each test (by a researcher blinded to the treatment group) was the sum of all four individual scores: 0 (best) – 48 (worst) score.

4. Infarct size

At the end of 24 h reperfusion, brains were then perfusion-fixed and 2 mm coronal sections were cut with a tissue cutter. The infarct area of the brain section was determined using 2% 2,3,5-triphenyltetrazolium chloride staining as described previously.¹⁶ The borders of the infarct in each brain slice were outlined and the area quantified using Image J software. For each brain section, the infarct volume was calculated by subtracting the area of the non-infarcted

ipsilateral hemisphere from that of the intact contralateral hemisphere. The percentage of infarct volume was determined by dividing the sum for the area of infarction by the total of that of contralateral hemisphere to avoid the influence of edema.²⁰

5. Apoptosis

The detection of apoptosis on brain paraffin sections was evaluated using the terminal deoxynucleotidyl transferase-mediated dUTP nick-end labeling (TUNEL) staining according to the manufacturer's instructions (Roche Diagnostics). The number of TUNEL-positive cells was counted in the cortex and the striatum by an observer blinded to the treatment groups using an Olympus microscope at $\times 400$.

6. Immunoblot analysis

The tissue samples from the striatum and cortex of the ipsilateral hemisphere were homogenized in a buffer consisting of 20 mM Tris (pH 7.5), 150 mM NaCl, 1% Triton, 1 mM Na₂EDTA, 1 mM EGTA, 2.5 mM sodium pyrophosphate, and protease inhibitors. Equal amounts of total protein from each group underwent immunoblotting assays. Proteins were separated on sodium dodecyl sulfate-polyacrylamide gel electrophoresis, and immunoblotting with anti-HMGB1, anti-phospho-inducible nitric oxide synthase (iNOS), anti-iNOS, anti-phospho-endothelial NOS (eNOS), anti-eNOS

(Abcam, Cambridge), anti-Toll-like receptor-4 (TLR-4), anti-Actin (Santa Cruz, CA), anti-Bcl-2, anti-Bax, anti-cytochrome C, anti-cleaved caspase-3, anti-caspase-3, anti-phospho-nuclear factor-kappa B (NF- κ B), and anti-NF- κ B, (all Cell Signaling Technology, Beverly, MA) was performed at least three times each.

7. Statistical analysis

All results are expressed as mean (SD). Serum glucose concentrations were analyzed using a repeated-measures analysis of variance (ANOVA). Other variables were analyzed using a two-way ANOVA. The P-values of post hoc tests were adjusted using Bonferroni's method. Values of $P < 0.05$ were considered statistically significant.

III. RESULTS

1. Functional and histological neuroprotection

Dextrose administration increased blood glucose concentrations >11.1 mmol/L until 24 h after reperfusion in all rats without any intergroup differences (S1 Fig). Hemodynamic data, including MAP and HR, also showed no intergroup differences (data not shown). No infarctions were observed in the sham group, while extensive infarct lesions developed in the striatum and lateral cortex of the control group. Compared with the control group, the infarct volume was significantly reduced in all rats in the pregabalin (30 mg/kg) group [$38.4 \pm$

5.5 % vs. 25.9 ± 3.5 %, $P < 0.05$] (Fig. 1a and 1b). The neurologic deficit score of the pregabalin group was significantly lower (better) compared with the control group [11 ± 3 vs. 16 ± 3 , $P < 0.05$] (Fig. 1c).




Figure 1. Determination of the dose range for pregabalin treatment.

Rats were treated with the indicated dose (mg/kg) of pregabalin (PGB). Representative pictures of TTC-stained brain sections from rat in the different group (A); Infarct volumes expressed as a percentage of area at risk (B); Neurological deficit scores (C). Values are mean \pm SD from 8 independent rats.

* $P < 0.05$. Scale bar: 1 cm.

There were more TUNEL-positive apoptotic cells in the penumbra cortex of the control group than in the pregabalin group [$63.9 \pm 5.6\%$ vs. $23.8 \pm 6.1\%$, $P < 0.05$] (Fig. 2).




Figure 2. Effect of pregabalin treatment upon reperfusion on apoptosis in the brain of hyperglycemic stroke rat.

Pregabalin reduced the number of TUNEL-positive cells compared with the control group. Quantitative representation of TUNEL-positive cells was determined by random counting of four fields per section. Values are mean \pm SD from 8 independent rats. * $P < 0.05$. Scale bar: 50 μm .

2. HMGB1/TLR-4/p-NF- κ B and pro-inflammatory cytokines

The expression level of HMGB1 was comparable between the sham group and the pregabalin group. Compared to that, it was significantly higher in the control group. Expression of TLR-4 was significantly increased in both the control and pregabalin groups compared with that in the sham group, while it was significantly lower in the pregabalin group than in the control group. Expression of p-NF- κ B was significantly increased in both the control and pregabalin groups compared with that in the sham group, while it was significantly lower in the pregabalin group than in the control group (Fig. 3). Compared with the sham group, expressions of IL-1 β and TNF- α were significantly increased both in the control and pregabalin groups. Their expression levels were significantly lower in the pregabalin group than in the control group.




Figure 3. Pregabalin treatment attenuated expressions of HMGB1/TLR-4/p-NF-κB/IL-1β and TNF-α in the brain of hyperglycemic stroke rat.

The protein levels of HMGB1, TLR-4, P-NF-κB, NF-κB, IL-1β and TNF-α were detected in the ipsilateral hemisphere of rat brain in each group using western blot analysis. Right panels were quantified signal by scanning densitometry. Values are mean \pm SD from 5 independent rats. *P < 0.05.

3. Nitric oxide synthases activity

The expression of p-iNOS could only be weakly detected in the sham group while it was significantly upregulated both in the control and pregabalin groups compared with that in the sham group. Its expression was significantly blunted in the pregabalin group compared with that in the control group. Levels of p-eNOS were decreased in the control group compared with those in the sham group, while its expression was retained in the pregabalin group (Fig. 4).




Figure 4. Pregabalin treatment favorably modulated expressions of NOS in the brain of hyperglycemic stroke rat.

The western blot analysis performed. The protein levels of P-iNOS and iNOS were detected in the ipsilateral hemisphere of rat brain in each group. Right panel was quantified signal by scanning densitometry. Values are mean \pm SD from 5 independent experiments. *P < 0.05.

4. Bcl-2, Bax, cytochrome C, and caspase-3 activity

Compared with the sham group, both the control and pregabalin groups exhibited a significant reduction in Bcl-2 expression, while it was significantly higher in the pregabalin group than in the control group. On the contrary, expressions of Bax and cytochrome C were significantly increased in both the control and pregabalin groups compared with those in the sham group, while their increases were significantly smaller in the pregabalin group compared with those in the control group (Fig. 5A).

With regard to cleaved caspase-3, its protein expression level was also significantly increased in both the control and pregabalin groups than in the sham group, while the increase was significantly smaller in the pregabalin group than in the control group (Fig. 5B).




Figure 5. Pregabalin treatment favorably modulated expressions of apoptosis-related molecules in the brain of hyperglycemic stroke rat.

The protein levels of Bcl-2, Bax, Cytochrome C (A), Cleaved caspase-3 and Caspase-3 (B) were detected in the ipsilateral hemisphere of rat brain in each group using western blot analysis. Right panels were quantified signal by scanning densitometry. Values are mean \pm SD from 5 independent rats. *P < 0.05.

IV. DISCUSSION

In the present study, we observed the functional and histological neuroprotective effects of pregabalin administration upon reperfusion after 1 h of MCAO in a rat model of hyperglycemic stroke. It reduced infarct size by 33%, improved neurological deficits (31% decrease in score), presented with fewer apoptotic cells (by 63%), and demonstrated favorable expression of apoptosis-regulatory molecules in the brain tissue. Activation of HMGB1/TLR-4 signaling and NF- κ B-mediated pro-inflammatory cascades was attenuated by pregabalin treatment. Favorable modulation of NOS isoforms was also revealed.

Hyperglycemia has been observed to be present in one-third of all ischemic stroke patients and is known to be associated with poorer prognosis regardless of diabetes history.¹ Of note, a previous meta-analysis reported even higher mortality of stroke in non-diabetic hyperglycemic patients compared with pre-existing diabetic patients.²¹ In conjunction, experimental studies have revealed that hyperglycemia exacerbates metabolic deterioration, intensifies the inflammatory response and oxidative stress, augments blood-brain barrier (BBB) disruption, and lowers the availability of NO, leading to increased infarct volume, edema, and hemorrhagic transformation.^{7,8,22,23} Even worse, several therapeutic modalities that have shown consistent efficacy in animal studies including the volatile anesthetic post-conditioning lost their neuroprotective effects or even enhanced cerebral damage in hyperglycemic states.^{10,11} The

results of intensive glycemic control in an attempt to attenuate exacerbation of stroke have also been largely disappointing.^{2,3}

One of the most prominent and initial metabolic issues related to ischemic stroke is distressed Ca^{2+} homeostasis and bursts of excitatory amino acids, which ultimately result in neuronal cell death[5]. Several N-methyl-D-aspartate (NMDA) receptor antagonists and calcium channel blockers (mostly L-type) have shown protective effects against stroke-related damage in a number of experimental models.²⁴ However, few of them have achieved successful clinical translations mainly because of their adverse psychotomimetic side effects or unwanted influence on the systemic vasculature lowering the perfusion pressure.^{25,26} In consideration of the exaggerated Ca^{2+} imbalance and neuronal excitability,⁷ hyperglycemic patients may require even higher doses, which may not become applicable on account of these safety issues.

By binding to VGCC, pregabalin suppresses Ca^{2+} influx and has an effect more upstream of the neuronal excitotoxicity, thus blocking the release of not only glutamate but also other excitatory amino acids and providing more comprehensive coverage compared with NMDA antagonists.¹³ Indeed, its protective effects have already been demonstrated in experimental models of normoglycemic MCAO and deep hypothermic circulatory arrest with cardiopulmonary bypass.^{15,16} Thus, we hypothesized that blocking this initial insult, neuroexcitotoxicity, may ameliorate its downstream events leading to neuronal death and convey neuroprotection against ischemic stroke, even under

hyperglycemic condition.

We found in the current hyperglycemic stroke experiment that pregabalin has potent anti-inflammatory properties via downregulation of the HMGB1/TLR-4 signaling pathway and attenuation of p-NF- κ B expression. HMGB1, a ubiquitous non-histone nuclear protein, is a key mediator of the immune mechanism in stroke. After extracellular release early after ischemia, it activates immune cells and induces inflammatory cytokine expression via binding to cell surface receptors even hours to days after reperfusion in the ischemic penumbra.⁶ Hyperglycemia enhances extracellular release of HMGB1, which has been demonstrated as an important mechanism for worsened ischemic damage.⁹ Expressions of its receptors implicated in stroke, including TLRs, have also been demonstrated to be increased by hyperglycemia.²⁷ Of note, experimental evidences indicate that HMGB1 may participate in further excitotoxicity in the penumbra after stroke by stimulating NMDA receptors and the release of excitatory amino acids.^{28,29} It is likely that pregabalin may have sufficient protective capacity to interrupt such a continuous damaging process in the hyperglycemic states.

One of the most detrimental effects of hyperglycemia in stroke is the more pronounced decrease in blood flow in the penumbra related to NO dysfunction, which may persist late into reperfusion.³⁰ eNOS activity is critical in maintaining adequate tissue perfusion and BBB integrity under central nervous system (CNS) trauma or inflammation, as well as ischemic stroke.^{31,32}

Accordingly, eNOS knockout mice exhibited larger infarction following ischemic stroke.³³ Hyperglycemia has been shown to decrease the activity of eNOS through downregulation of the phosphatidylinositol 3-kinase/Akt pathway.³⁴ In the present study, pregabalin preserved eNOS activity in hyperglycemic stroke rats, which might also have played a role in its neuroprotective effects.

On the other hand, we observed decreased phosphorylation of iNOS following pregabalin treatment. iNOS has been regarded as a major contributor to the initiation and exacerbation of inflammatory and degenerative conditions of the CNS.³⁵ It has been associated with superoxide and peroxynitrite generation and nitrosylation of mitochondrial metabolic enzymes, leading to neuronal cell death.^{35,36} The essential molecule for the gene transcription and mRNA stability of iNOS during stroke is NF- κ B, which is activated by oxidative stress, glutamate, pro-inflammatory cytokines, and by hyperglycemia *per se*.³⁷ Downregulation of iNOS in our hyperglycemic stroke rats by pregabalin might have been attributable to inhibition of the NF- κ B transcription pathway, cytokine production, and oxidative stress.

Although complexly interrelated, the overall hierarchy of neuronal injury after oxygen deprivation starts with excitotoxicity and necrosis, which triggers an inflammatory response that causes further neuronal loss by apoptosis.⁵ An important downstream mediator of the neuro-excitotoxicity linked with inflammation was found to be HMGB1, which subsequently influences a

well-known key inflammatory regulator, NF- κ B.⁶ In that context, our results provide primary evidence that blocking the foremost and universal event after neuronal energy depletion, excitotoxicity, with pregabalin may improve outcome after ischemic stroke under hyperglycemic conditions. The current study confirms that pregabalin's beneficial influences are present in all of the above mentioned key inflammatory regulators, NOS, and the anti-apoptotic pathways of neuronal cell death after I-R. An additional strength of the current study is that it firstly addressed the neuroprotective effect of a drug with known clinical safety and reliable pharmacokinetics³⁸ against hyperglycemic stroke. Considering the high prevalence of hyperglycemia with stroke and its indisputable association with adverse outcome while hindering potentially effective treatments, our results should be of clinical relevance that merits further clinical trials.

This study is subject to the following limitations. First, we cannot draw any conclusion regarding the potential direct influence of pregabalin on each of the specific downstream pathways of neuronal cell death as there are no specific antagonists to pregabalin and it blocks more than just the release of glutamate, including other excitatory neurotransmitters. Second, we did not make any direct comparisons with normoglycemic conditions as they were already reported in a previous study of a mouse MCAO model, which indirectly investigated its calcium-related mechanisms.¹⁶ In that study, although not completely identical in terms of the methods, pregabalin resulted in up to 70%

reduction in infarct size as opposed to 33% in our study implicating the adverse influence of hyperglycemia. Moreover, mechanisms for hyperglycemia-induced exacerbation of the stroke pathophysiology including excitotoxicity and HMGB1 cascades have been already well established.^{7-9,22,23} While avoiding redundancies, we have provided evidence that neuroprotection against stroke can be achieved in the hyperglycemic states in which existing therapeutic modalities have lost their efficacies.^{39,40} Third, although we could clearly delineate pregabalin's neuroprotective effects at 24 h after reperfusion, delayed neurologic injury may further ensue 24-72 h after reperfusion, which merits a further study.

V. CONCLUSION

Pregabalin administration upon reperfusion conveyed significant functional and histological neuroprotection after 1 h of MCAO under hyperglycemia. These beneficial influences were observed to be associated with major regulators of inflammation, HMGB1, and subsequently, NF- κ B and anti-apoptotic pathways, as well as NOS.

REFERENCES

1. Luitse MJ, Biessels GJ, Rutten GE, Kappelle LJ. Diabetes, hyperglycaemia, and acute ischaemic stroke. *Lancet Neurol.* 2012; 11: 261-71.
2. Gray CS, Hildreth AJ, Sandercock PA, O'Connell JE, Johnston DE, Cartledge NE, et al.; GIST Trialists Collaboration. Glucose-potassium-insulin infusions in the management of post-stroke hyperglycaemia: the UK Glucose Insulin in Stroke Trial (GIST-UK). *Lancet Neurol.* 2007; 6: 397-406.
3. Bruno A, Kent TA, Coull BM, Shankar RR, Saha C, Becker KJ, et al. Treatment of hyperglycemia in ischemic stroke (THIS): a randomized pilot trial. *Stroke.* 2008; 39: 384-9.
4. MacDougall NJ, Muir KW. Hyperglycaemia and infarct size in animal models of middle cerebral artery occlusion: systematic review and meta-analysis. *J Cereb Blood Flow Metab.* 2011; 31: 807-18.
5. Lipton P. Ischemic cell death in brain neurons. *Physiol Rev.* 1999; 79: 1431-568.
6. Kim JB, Sig Choi J, Yu YM, Nam K, Piao CS, Kim SW, et al. HMGB1, a novel cytokine-like mediator linking acute neuronal death and delayed neuroinflammation in the postischemic brain. *J Neurosci.* 2006; 26: 6413-21.
7. Li PA, Shuaib A, Miyashita H, He QP, Siesjo BK, Warner DS. Hyperglycemia enhances extracellular glutamate accumulation in rats subjected to forebrain ischemia. *Stroke.* 2000; 31: 183-92.
8. Suh SW, Shin BS, Ma H, Van Hoecke M, Brennan AM, Yenari MA, et al. Glucose and NADPH oxidase drive neuronal superoxide formation in stroke. *Ann Neurol.* 2008; 64: 654-63.
9. Huang J, Liu B, Yang C, Chen H, Eunice D, Yuan Z. Acute hyperglycemia worsens ischemic stroke-induced brain damage via high mobility group box-1 in rats. *Brain Res.* 2013; 1535: 148-55.
10. Hu Q, Ma Q, Zhan Y, He Z, Tang J, Zhou C, et al. Isoflurane enhanced hemorrhagic transformation by impairing antioxidant enzymes in hyperglycemic rats with middle cerebral artery occlusion. *Stroke.* 2011; 42: 1750-6.
11. Huang M, Qian Y, Guan T, Huang L, Tang X, Li Y. Different neuroprotective responses of Ginkgolide B and bilobalide, the two Ginkgo components, in ischemic rats with hyperglycemia. *Eur J Pharmacol.* 2012; 677: 71-6.
12. Davalos A, Castillo J, Serena J, Noya M. Duration of glutamate release after acute ischemic stroke. *Stroke.* 1997; 28: 708-10.

13. Joshi I, Taylor CP. Pregabalin action at a model synapse: Binding to presynaptic calcium channel $\alpha(2)$ -delta subunit reduces neurotransmission in mice. *Eur J Pharmacol.* 2006; 553: 82-8.
14. Ha KY, Carragee E, Cheng I, Kwon SE, Kim YH. Pregabalin as a Neuroprotector after Spinal Cord Injury in Rats: Biochemical Analysis and Effect on Glial Cells. *J Korean Med Sci.* 2011; 26: 404-11.
15. Shim JK, Ma Q, Zhang Z, Podgoreanu MV, Mackensen GB. Effect of pregabalin on cerebral outcome after cardiopulmonary bypass with deep hypothermic circulatory arrest in rats. *J Thorac Cardiovasc Surg.* 2014; 148: 298-303.
16. Yoon JS, Lee JH, Son TG, Mughal MR, Greig NH, Mattson MP. Pregabalin suppresses calcium-mediated proteolysis and improves stroke outcome. *Neurobiol Dis.* 2011; 41: 624-9.
17. Hirose R, Xu F, Dang K, Liu T, Behrends M, Brakeman PR, et al. Transient hyperglycemia affects the extent of ischemia-reperfusion-induced renal injury in rats. *Anesthesiology.* 2008; 108: 402-14.
18. Longa EZ, Weinstein PR, Carlson S, Cummins R. Reversible Middle Cerebral-Artery Occlusion without Craniectomy in Rats. *Stroke.* 1989; 20: 84-91.
19. Yokoo N, Sheng HX, Mixco J, Homi HM, Pearlstein RD, Warner DS. Intraischemic nitrous oxide alters neither neurologic nor histologic outcome: A comparison with dizocilpine. *Anesth Analg.* 2004; 99: 896-903.
20. Swanson RA, Sharp FR. Infarct measurement methodology. *J Cereb Blood Flow Metab.* 1994; 14: 697-8.
21. Capes SE, Hunt D, Malmberg K, Pathak P, Gerstein HC. Stress hyperglycemia and prognosis of stroke in nondiabetic and diabetic patients: a systematic overview. *Stroke.* 2001; 32: 2426-32.
22. Morigi M, Angioletti S, Imberti B, Donadelli R, Micheletti G, Figliuzzi M, et al. Leukocyte-endothelial interaction is augmented by high glucose concentrations and hyperglycemia in a NF- κ B-dependent fashion. *J Clin Invest.* 1998; 101: 1905-15.
23. Fabian RH, Kent TA. Hyperglycemia Accentuates Persistent "Functional Uncoupling" of Cerebral Microvascular Nitric Oxide and Superoxide Following Focal Ischemia/Reperfusion in Rats. *Transl Stroke Res.* 2012; 3: 482-90.
24. Kaur H, Prakash A, Medhi B. Drug therapy in stroke: from preclinical to clinical studies. *Pharmacology.* 2013; 92: 324-34.
25. Davis SM, Lees KR, Albers GW, Diener HC, Markabi S, Karlsson G, et al. Selfotel in acute ischemic stroke - Possible neurotoxic

- effects of an NMDA antagonist. *Stroke*. 2000; 31: 347-54.
26. Kaste M, Fogelholm R, Erila T, Palomäki H, Murros K, Rissanen A, et al. A randomized, double-blind, placebo-controlled trial of nimodipine in acute ischemic hemispheric stroke. *Stroke*. 1994; 25: 1348-53.
27. Carrillo-Sepulveda MA, Spitler K, Pandey D, Berkowitz DE, Matsumoto T. Inhibition of TLR4 attenuates vascular dysfunction and oxidative stress in diabetic rats. *J Mol Med (Berl)*. 2015; 93: 1341-54.
28. Pedrazzi M, Raiteri L, Bonanno G, Patrone M, Ledda S, Passalacqua M, et al. Stimulation of excitatory amino acid release from adult mouse brain glia subcellular particles by high mobility group box 1 protein. *J Neurochem*. 2006; 99: 827-38.
29. Balosso S, Liu J, Bianchi ME, Vezzani A. Disulfide-containing high mobility group box-1 promotes N-methyl-D-aspartate receptor function and excitotoxicity by activating Toll-like receptor 4-dependent signaling in hippocampal neurons. *Antioxid Redox Signal*. 2014; 21: 1726-40.
30. Kawai N, Keep RF, Betz AL. Hyperglycemia and the vascular effects of cerebral ischemia. *Acta Neurochir Suppl*. 1997; 70: 27-9.
31. Villalba N, Sackheim AM, Nunez IA, Hill-Eubanks DC, Nelson MT, Wellman GC, et al. Traumatic Brain Injury Causes Endothelial Dysfunction in the Systemic Microcirculation through Arginase-1-Dependent Uncoupling of Endothelial Nitric Oxide Synthase. *J Neurotrauma*. 2016 Apr 1.
32. Austin SA, Santhanam AV, Hinton DJ, Choi DS, Katusic ZS. Endothelial nitric oxide deficiency promotes Alzheimer's disease pathology. *J Neurochem*. 2013; 127: 691-700.
33. Huang Z, Huang PL, Ma J, Meng W, Ayata C, Fishman MC, et al. Enlarged infarcts in endothelial nitric oxide synthase knockout mice are attenuated by nitro-L-arginine. *J Cereb Blood Flow Metab*. 1996; 16: 981-7.
34. Du XL, Edelstein D, Dimmeler S, Ju QD, Sui C, Brownlee M. Hyperglycemia inhibits endothelial nitric oxide synthase activity by posttranslational modification at the Akt site. *J Clinical Invest*. 2001; 108: 1341-8.
35. Hickey MJ, Granger DN, Kubes P. Inducible nitric oxide synthase (iNOS) and regulation of leucocyte/endothelial cell interactions: studies in iNOS-deficient mice. *Acta Physiol Scand*. 2001; 173: 119-26.
36. Pannu R, Singh I. Pharmacological strategies for the regulation of inducible nitric oxide synthase: Neurodegenerative versus

- neuroprotective mechanisms. *Neurochem Int.* 2006; 49: 170-82.
37. Bemeur C, Ste-Marie L, Montgomery J. Increased oxidative stress during hyperglycemic cerebral ischemia. *Neurochem Int.* 2007; 50: 890-904.
38. Toth C. Pregabalin: latest safety evidence and clinical implications for the management of neuropathic pain. *Ther Adv Drug Safe.* 2014; 5: 38-56.
39. Won SJ, Tang XN, Suh SW, Yenari MA, Swanson RA. Hyperglycemia promotes tissue plasminogen activator-induced hemorrhage by Increasing superoxide production. *Ann Neurol.* 2011; 70: 583-90.
40. Masrur S, Cox M, Bhatt DL, Smith EE, Ellrodt G, Fonarow GC, et al. Association of Acute and Chronic Hyperglycemia With Acute Ischemic Stroke Outcomes Post-Thrombolysis: Findings From Get With The Guidelines-Stroke. *J Am Heart Assoc.* 2015; 4: e002193.

ABSTRACT (IN KOREAN)

고혈당 뇌졸중 유도 백서에서 재관류 시 프레가발린 투여가 미치는 영향과 High-mobility group box 1 관련 기전

<지도교수 심재광>

연세대학교 대학원 의학과

송영

고혈당은 뇌졸중에서 흔하게 발생하고, 치료 효과를 저해하며 예후를 악화시킨다. Pregabalin의 신경 흥분 독성에 대한 억제 작용이 고혈당 조건에서도 뇌졸중에 대한 보호 효과를 나타내는지 알아보고 관련된 분자 기전을 규명하고자 하였다.

Dextrose 투여로 고혈당을 유도한 백서에 1시간의 중대뇌동맥 폐색과 재관류를 시행하였다. 재관류 시작과 동시에 각각 16마리의 pregabalin 군과 대조군에 30 mg/kg의 pregabalin 또는 같은 양의 생리 식염수를 복강 내 주사하였다. 재관류 24 시간 후, 신경학적 장애 척도, 뇌 경색 면적 및 자멸사 세포 수 측정을 시행하였다. 백서의 뇌조직 내 High-mobility group box 1 (HMGB1), Toll-like receptor-4 (TLR-4), phosphorylated nuclear factor-kappa B (p-NF- κ B), interleukin-1 β (IL-1 β), tumor necrosis factor-alpha (TNF- α), inducible nitric oxide synthase (iNOS), endothelial nitric oxide synthase (eNOS), Bcl-2, Bax, Cytochrome C, 그리고

Caspase-3의 단백질 발현을 알아보기 위하여 Western blot 분석을 시행하였다.

대조군과 비교하여 pregabalin군에서 현저히 신경학적 장애가 적었고 뇌경색 면적은 33% 감소하였으며 자멸사 세포 수는 63% 적게 관찰되었다. 뇌조직 내 HMGB1, TLR-4, p-NF- κ B, IL-1 β , 그리고 TNF- α 의 발현은 대조군에 비해 유의하게 감소되어 있었고, p-iNOS 발현의 감소 및 p-eNOS 발현의 증가 소견이 관찰되었다. 또한 pregabalin 군에서 Bax, Cytochrome C, 및 cleaved caspase-3/caspase-3의 발현은 유의하게 감소되어 있었고, Bcl-2의 발현은 증가되어 있었다.

고혈당 뇌졸중 유도 백서에서 재관류 시점의 pregabalin 투여는 세포사를 줄이고 신경학적 기능을 향상시켰다. 이는 HMGB1/TLR4 매개 염증 반응에 대한 억제 및 NOS에 대한 유익한 조절 작용 기전과 관련된 것으로 생각된다.

핵심되는 말: 뇌졸중, 고혈당, pregabalin, high-mobility group box 1, toll-like receptor-4, nitric oxide synthase