



저작자표시-비영리-변경금지 2.0 대한민국

이용자는 아래의 조건을 따르는 경우에 한하여 자유롭게

- 이 저작물을 복제, 배포, 전송, 전시, 공연 및 방송할 수 있습니다.

다음과 같은 조건을 따라야 합니다:



저작자표시. 귀하는 원저작자를 표시하여야 합니다.



비영리. 귀하는 이 저작물을 영리 목적으로 이용할 수 없습니다.



변경금지. 귀하는 이 저작물을 개작, 변형 또는 가공할 수 없습니다.

- 귀하는, 이 저작물의 재이용이나 배포의 경우, 이 저작물에 적용된 이용허락조건을 명확하게 나타내어야 합니다.
- 저작권자로부터 별도의 허가를 받으면 이러한 조건들은 적용되지 않습니다.

저작권법에 따른 이용자의 권리는 위의 내용에 의하여 영향을 받지 않습니다.

이것은 [이용허락규약\(Legal Code\)](#)을 이해하기 쉽게 요약한 것입니다.

[Disclaimer](#)

Validation of pediatric index of  
mortality 3 for predicting mortality  
among pediatric patients admitted to an  
intensive care unit in Korea

Jae Hwa Jung

Department of Medicine

The Graduate School, Yonsei University

Validation of pediatric index of  
mortality 3 for predicting mortality  
among pediatric patients admitted to an  
intensive care unit in Korea

Directed by Professor Myung Hyun Sohn

The Master's Thesis  
submitted to the Department of Medicine,  
the Graduate School of Yonsei University  
in partial fulfillment of the requirements for the degree  
of Master of Medical Science

Jae Hwa Jung

June 2017

This certifies that the Master's Thesis of  
Jae Hwa Jung is approved.

-----  
Thesis Supervisor: Myung Hyun Sohn

-----  
Thesis Committee Member#1: Kyung Won Kim

-----  
Thesis Committee Member#2: Moo Suk Park

The Graduate School  
Yonsei University

June 2017

## ACKNOWLEDGEMENTS

I am deeply grateful to my beloved family, friends, and esteemed professor. This paper is dedicated to those who have given me a lot of attention and encouragement.

## <TABLE OF CONTENTS>

ABSTRACT .....	1
I. INTRODUCTION .....	3
II. MATERIALS AND METHODS .....	5
1. Study design .....	5
2. Patient selection .....	5
3. Data collection .....	5
4. Statistical analysis .....	6
5. Ethics statement .....	7
III. RESULTS .....	8
1. Clinical characteristics of patients .....	8
2. PIM 3 between survivor and non-survivor group .....	10
3. PIM 3 performance .....	11
IV. DISCUSSION .....	15
V. CONCLUSION .....	18
REFERENCES .....	19
ABSTRACT (IN KOREAN) .....	21

## LIST OF FIGURES

Figure 1. Area under the receiver operating characteristic curve (AUC) of PIM 2, PIM 3, PRISM III .....	11
Figure 2. Calibration curves for observed mortality against predicted risk of death for the PIM 2, PIM 3 models .....	14

## LIST OF TABLES

Table 1. Demographic and clinical characteristics of the study population .....	9
Table 2. Characteristics of variables in PIM 3 between survivor and non-survivor .....	10
Table 3. Hosmer-Lemeshow test for deciles of risk .....	13

## ABSTRACT

Validation of pediatric index of mortality 3 for predicting mortality among pediatric patients admitted to an intensive care unit in Korea

Jae Hwa Jung

*Department of Medicine  
The Graduate School, Yonsei University*

(Directed by Professor Myung Hyun Sohn)

**Background:** For advancement in pediatric critical care medicine, the precise prediction of survival is considered imperative. The pediatric index of mortality (PIM) 3 was developed in 2013; it is a representative and simple model consisting of 10 variables that are measured at the time of admission to intensive care units (ICUs).

**Objectives:** The objectives of this study were to evaluate the usefulness of the PIM 3 for predicting mortality and validating PIM 3 in Korean children admitted to a single intensive care unit.

**Subjects and Methods:** We enrolled children at least 1 month old but less than 18 years of age who were admitted to the Severance Hospital Medical ICU between March 2009 and February 2015. Performances of the pediatric risk of mortality (PRISM) III, PIM 2, and PIM 3 were evaluated by assessing the area under the receiver operating characteristic (ROC) curve, conducting the

Hosmer-Lemeshow test, and calculating the standardized mortality ratio (SMR).

**Results:** In total, 503 children were enrolled; the areas under the ROC curve for PRISM III, PIM 2, and PIM 3 were 0.775, 0.796, and 0.826, respectively. The area under the ROC curve was significantly greater for PIM 3 than for PIM 2 ( $P < 0.001$ ) and PRISM III ( $P = 0.016$ ). There were no significant differences in the Hosmer-Lemeshow test results for PRISM III ( $P = 0.498$ ), PIM 2 ( $P = 0.249$ ), and PIM 3 ( $P = 0.337$ ). The SMR calculated using PIM 3 (1.11) was closer to 1 than that calculated using PIM 2 (0.84).

**Conclusion:** PIM 3 showed better prediction for the risk of mortality than PIM 2 for the Korean pediatric population admitted in the ICU.

---

Key words: PIM 3, children, intensive care unit, risk of mortality, validation study

Validation of pediatric index of mortality 3 for predicting mortality  
among pediatric patients admitted to an intensive care unit in Korea

Jae Hwa Jung

*Department of Medicine*  
*The Graduate School, Yonsei University*

(Directed by Professor Myung Hyun Sohn)

## I. INTRODUCTION

For advancement in pediatric critical care medicine, the precise prediction of survival is considered imperative<sup>1</sup>. However, there are many challenges in pediatric survival prediction<sup>2,3</sup>. For example, accurate assessment of consciousness in non-verbal children is limited<sup>4</sup>, and normal ranges of blood pressure and physiological parameters for children will depend on their age, weight, and height<sup>5</sup>. In addition, there is difficulty in performing invasive procedures to measure prediction parameters.

Since the early 1980s, various scoring systems have been used in pediatric intensive care units (ICU) to evaluate disease severity in patients<sup>6-10</sup>. These scoring systems assist in the prediction of patient mortality and allow comparison of standards of care in different ICUs. Among the scoring systems, the pediatric index of mortality (PIM) has been considered a representative model. The PIM, developed from data collected from pediatric ICUs in New Zealand, the United Kingdom, and Australia between 1994 and 1996 by Shann *et al.*<sup>8</sup>, is a simple model that consists of 10 variables measured at the time of admission to the ICU<sup>9</sup>. In addition to mortality prediction, the PIM had a function for evaluating the quality of medical care and optimizing the employment of resources<sup>11,12</sup>.

Another widely used prediction model, the pediatric risk of mortality (PRISM)

model, was developed using data collected from pediatric ICUs in the United States between 1980 and 1985 by Pollack *et al.*<sup>7</sup>; this was later updated in 1996<sup>10</sup>. The PRISM III score is calculated from the patient's physiologic data, descriptive information, and diagnosis. The PRISM III predicts the risk for mortality during that admission.

The PIM models were assessed by comparing them with prediction models, such as PRISM III, in order to determine their ability to predict prognostic scores<sup>13-15</sup>. The results of the first study investigating the PIM showed that it was a good model for the prediction and classification of mortality in groups of children and adolescents in the ICU<sup>8</sup>. One advantage of the PIM model over the PRISM model is that it is based on 10 variables only, all of which are collected at the point of admission; the PRISM requires 24 hours of intensive care management before data can be collected<sup>16</sup>. Several other studies have reported that the previous PIM models have performed well in the prediction of mortality<sup>14,17,18</sup>.

The various mortality prediction models need to be validated before they can be applied in environments that differ from those in which they were developed. In 2003, the PIM study group published a revised version of the PIM. Compared with the original version, PIM 2 showed better calibration and had better adjustment for varying diagnostic groups<sup>19</sup>. In Korea, PIM 2 had better results than PRISM III in mortality prediction<sup>20</sup>. The PIM was again updated in 2013 and was called the PIM 3; however, this new version has not yet been evaluated in Korea.

The objectives of this study were to evaluate the usefulness of the PIM 3 for predicting mortality and to validate PIM 3 in Korean children admitted to a single intensive care unit. Additionally, we aimed to determine other factors strongly correlated with the predicted mortality rate.

## II. MATERIALS AND METHODS

### 1. Study Design

This was a retrospective study of children admitted to the Severance Hospital Medical ICU at Yonsei University between March 2009 and February 2015.

### 2. Patient Selection

All patients at least 1 month old but less than 18 years of age who were admitted to the Severance Hospital Medical ICU during the study period were included. Patients who were discharged or who died within 24 hours of ICU admission were excluded. Additionally, patients were excluded from the study if data entry was incomplete. Neonates in the neonatal ICU, children in post-operative recovery in the surgical ICU, and children with cardiac problems in the cardiac ICU were not included.

### 3. Data Collection

This was a retrospective study in which data were collected from medical records. Demographic data were collected from all study participants, including age, sex, diagnosis, admission route, and length of stay in the ICU. Patients were divided into age groups (infants, preschoolers, schoolchildren, and adolescents). The admission route was divided into the general ward, emergency room, and operating room. According to the original studies, 17 physiological data and 8 additional risk factors were collected for the PRISM III, and 10 physiological data were collected for the PIM 2 and PIM 3. The outcome of interest was survival; a non-survivor was defined as a patient who died in the ICU, who discharged for palliative care, or who died within 24 hours of discharge. Experienced nurses and ICU doctors reviewed all the data.

The PRISM III, PIM 2, and PIM 3 scores were calculated using the formulas available in published findings<sup>9,10</sup>. The PRISM III score was calculated using 17 physiological parameters and additional risk factors. Physiologic data included

the most abnormal values among 17 variables in the first 24 hours of the ICU stay. The PIM 3 score was calculated using 10 physiological variables collected within the first hour of admission to the ICU.

#### 4. Statistical analysis

Parametrically distributed values were expressed as means  $\pm$  standard deviations (SDs). Demographic and physiologic data were described using Student's t-tests for continuous variables and the chi-square tests for categorical variables. The performance of each scoring system was evaluated by assessing area under the receiver operating characteristic (ROC) curve, conducting the Hosmer-Lemeshow test, and calculating the standardized mortality ratio (SMR).

The capacity for discrimination between a survivor and non-survivor was made using the area under the ROC curve. An area under the ROC curve of 0.75 or more was considered statistically significant<sup>21</sup>. The areas under the ROC curve for PRISM III, PIM 2, and PIM 3 were compared using the chi-square test.

Statistical analysis of the calibration of each model was conducted using the Hosmer-Lemeshow test. The Hosmer-Lemeshow test is a statistical test that compares the observed and expected mortality risk for goodness of fit<sup>22</sup>. To assess the calibration of the ICU scoring systems, the expected and observed numbers of survivors and mortalities in deciles of risk were calculated. A large, significant difference in the observed and expected values signifies that the model did not meet the goodness of fit test. In this statistical analysis, the model was judged appropriate when the *P* was 0.05 or more<sup>23</sup>.

The overall prediction for the risk of mortality was assessed by calculating the expected and observed numbers of survivors and mortalities using the SMR calculation. The SMR with a 95% confidence interval (CI) was calculated by dividing the actual mortality rate by the cumulative predicted mortality rate for the study population<sup>24</sup>. An SMR close to 1 signifies better prediction; an SMR

significantly less or more than 1 could be interpreted as an overestimation or underestimation of mortality in the ICU.

Statistical analysis and data management were performed using Excel (Excel 2010; Microsoft) and the Statistical Package for the Social Sciences (SPSS Inc.) version 23.0. *P*-values less than 0.05 were considered statistically significant.

#### 5. Ethics statement

All data were collected and analyzed retrospectively in this study. This study was approved by the Institutional Review Board of Severance Hospital (Seoul, Korea, IRB No. 4-2015-1194).

### III. RESULTS

#### 1. Clinical characteristics of patients

In total, 583 patients were included in the study; however, 29 were excluded because they died or were discharged within 24 hours of admission, and 51 were excluded because data were incomplete. Therefore, 503 patients were actually enrolled in the study.

Table 1 shows the clinical and demographic characteristics of the patients in the study population. There were 297 (59.0%) male patients and 206 (41.0%) female patients. The majority of patients were preschoolers at ICU admission ( $n = 215$ , 42.7%). Diagnostic categories at ICU admission included respiratory problems, such as respiratory infections or airway obstructions ( $n = 267$ , 53.1%); neurologic problems, such as status epilepticus or central nervous system infections ( $n = 96$ , 19.1%); postoperative care ( $n = 41$ , 8.2%); hematology-oncologic problems, such as neutropenic fever during chemotherapy or tumor lysis syndrome ( $n = 34$ , 6.8%); nephrology problems, such as renal failure or need for renal replacement therapy ( $n = 26$ , 5.2%); gastrointestinal problems, such as intestinal bleeding or fulminant hepatitis ( $n = 20$ , 4.0%); and others, such as cardiac arrest of unknown origin ( $n = 19$ , 3.8%). In total, 51.1% ( $n = 257$ ) of patients were admitted from the general ward. The overall ICU mortality rate was 19.9% ( $n = 100$ ). According to the admission route, the mortality rates were 26.8%, 12.3%, and 14.7% for general ward, emergency room, and operating room, respectively.

There were no differences in sex ( $P = 0.369$ ) and time to ICU discharge ( $P = 0.068$ ) between the survivor and non-survivor group. Patients were older in the non-survivor group than in the survivor group ( $P = 0.003$ ).

Table 1. Demographic and clinical characteristics of the study population

	Survivor (n=403)	Non-survivor (n=100)	Total (n=503)
Male	234 (58.1)	63 (63.0)	297 (59.0)
Age (year)	4.5±4.5	6.2±5.0	4.8±4.6
ICU stay duration (days)	14.7±18.7	26.8±65.1	17.1±34.5
Class of age			
Infants (1-11 months)	104 (90.4)	11 (9.6)	115 (22.9)
Preschoolers (1-5 years)	169 (78.6)	46 (21.4)	215 (42.7)
Schoolchildren (6-11 years)	95 (77.9)	27 (22.1)	122 (24.3)
Adolescents (12-17 years)	35 (68.6)	16 (31.4)	51 (10.1)
Main problem			
Pulmonology	216 (80.9)	51 (19.1)	267 (53.1)
Neurology	82 (85.4)	14 (14.6)	96 (19.1)
Post operation/procedure	39 (95.1)	2 (4.9)	41 (8.2)
Hemato-oncology	17 (50.0)	17 (50.0)	34 (6.8)
Nephrology	20 (76.9)	6 (23.1)	26 (5.2)
Gastroenterology	13 (65.0)	7 (35.0)	20 (4.0)
Others	16 (84.2)	3 (15.8)	19 (3.8)
Admission route			
General ward	188 (73.2)	69 (26.8)	257 (51.1)
Emergency room	186 (87.7)	26 (12.3)	212 (42.1)
Operating room	29 (85.3)	5 (14.7)	34 (6.8)

Abbreviation: ICU, Intensive Care Unit

Data are expressed as n (%) for discrete variable or as otherwise indicated.

## 2. PIM 3 between the survivor and non-survivor groups

There were significant differences in the PIM 3 scoring parameters between survivors and non-survivors, as shown in Table 2.

Table 2. Characteristics of variables in PIM 3 between the survivor and non-survivor

Parameters	Survivor (n=403)	Non-survivor (n=100)	<i>P</i> value
<b>Systolic BP (mmHg)</b>			
Infants (1-11 months)	84.4±17.9	70.3±18.2	0.004
Preschoolers (1-5 years)	87.7±17.3	72.3±24.2	<0.001
Schoolchildren (6-11 years)	95.2±21.2	83.3±25.6	0.017
Adolescents (12-17 years)	93.6±19.7	80.6±27.0	0.033
No pupil light reflex	9 (2.2)	20 (20.0)	<0.001
FiO <sub>2</sub> x 100/PaO <sub>2</sub>	0.396±0.297	0.734±0.614	<0.001
Base Excess	-2.1±7.0	-4.7±9.7	0.002
Mechanical ventilation	362 (89.8)	97 (97.0)	0.013
Elective admission	33 (8.1)	2 (1.8)	0.019
Recovery	38 (9.3)	3 (2.7)	0.027
<b>Risk assessment</b>			
Low-risk group	26 (6.5)	0 (0.0)	0.009
High-risk group	115 (28.5)	21 (21.0)	0.129
Very high-risk group	42 (10.5)	59 (59.0)	<0.001

Abbreviation: PIM 3, Pediatric Index of Mortality 3; BP, blood pressure; FiO<sub>2</sub>, Fraction of inspired Oxygen; PaO<sub>2</sub>, Arterial blood oxygen partial pressure

Very-high risk group: Cardiac arrest, Severe combined immune deficiency, Leukemia or lymphoma after first induction, Bone marrow transplant recipient, Liver failure.

The mortality rates were 15.4% and 58.4% for the high-risk and very high-risk groups, respectively. More patients in the survivor group recovered from surgery and/or the other procedures investigated in this study ( $P = 0.027$ ). When compared with survivors, the non-survivors were more likely to be treated with a ventilator at the time of ICU admission ( $P = 0.013$ ) or categorized as being at very high risk for mortality ( $P < 0.001$ ).

### 3. PIM 3 performance

The areas under the ROC curve were 0.775 (95% CI 0.736-0.811), 0.796 (95% CI 0.758-0.831), and 0.826 (95% CI 0.790-0.858) for PRISM III, PIM 2, and PIM 3, respectively. The area under the ROC curve was significantly greater for PIM 3 than for PIM 2 ( $P < 0.001$ ) and PRISM III ( $P = 0.016$ ) (Fig. 1).




Fig. 1. Area under the receiver operating characteristic curve (AUC) of PIM 2, PIM 3, PRISM III

Area under ROC of PIM 3 was significantly greater than PIM 2 ( $P < 0.001$ ) and PRISM III ( $P = 0.016$ ).

Table 3 shows the overall calibration of the models across the deciles of mortality risk by using the Hosmer-Lemeshow test. There were no significant differences observed in expected deaths (PRISM III  $P = 0.498$ , PIM 2  $P = 0.249$ , PIM 3  $P = 0.337$ ). All three models showed goodness of fit.

**Table 3.** Hosmer-Lemeshow test for deciles of risk

PRISM III				PIM 2				PIM 3			
Values	n	O	E	Values	n	O	E	Values	n	O	E
0-0	83	2	4.78	0.005-0.030	50	2	1.53	0.002-0.016	50	2	1.12
1-2	41	1	2.91	0.030-0.052	50	4	2.92	0.016-0.038	51	3	2.93
3-4	37	6	3.08	0.052-0.070	50	4	3.78	0.038-0.053	50	3	4.04
5-5	48	5	4.90	0.070-0.092	50	0	4.61	0.053-0.065	50	3	4.80
6-7	44	7	5.42	0.092-0.127	50	5	5.47	0.065-0.086	50	3	5.64
8-9	50	9	7.41	0.127-0.238	50	7	7.81	0.086-0.138	51	7	7.79
10-11	44	8	8.02	0.238-0.303	50	15	10.44	0.138-0.177	50	6	9.55
12-14	48	12	11.07	0.303-0.365	50	9	12.37	0.177-0.251	50	13	11.52
15-20	52	17	17.75	0.365-0.674	50	19	16.27	0.251-0.514	50	23	16.46
21-51	56	33	34.65	0.674-0.996	53	35	34.81	0.514-0.999	51	37	36.15
	503	100			503	100			503	100	
$\chi^2(8) = 7.36, P = 0.498$				$\chi^2(8) = 10.24, P = 0.249$				$\chi^2(8) = 9.06, P = 0.337$			

Abbreviation: PRISM, Pediatric Risk of Mortality; PIM 2, Pediatric Index of Mortality 2; PIM 3, Pediatric Index of Mortality 3; O = observed deaths, E = expected deaths

The SMRs were 0.84 (95% CI 0.68-1.02) and 1.11 (95% CI 0.91-1.35) for PIM scores 2 and 3, respectively. The SMR calculated using PIM 3 was closer to 1 than that calculated using PIM 2 (Fig. 2.).




Fig. 2. Calibration curves for observed mortality against predicted risk of death for the PIM 2, PIM 3 models.

Abbreviation: PIM 2, Pediatric Index of Mortality 2; PIM 3, Pediatric Index of Mortality 3; SMR, Standardized Mortality Ratio

Filled squares, PIM 3 SMR; open triangles, PIM 2 SMR; dotted line, line for equal observed and expected deaths (SMR = 1)

#### IV. DISCUSSION

This is the first study that assessed the new version of PIM 3 in Korean children, in an environment substantially different from the environment in which PIM 3 was generated. We compared the performances of PRISM III, PIM 2, and PIM 3 at a single institution using three statistical methods: area under the ROC curve, the Hosmer-Lemeshow test, and SMR. On the basis of the results of these three statistical methods, we conclude that PIM 3 performs better for predicting mortality than PIM 2 or PRISM III, and it is suitable for use in the Korean children.

There are major changes in the grouping system in the PIM 3 scoring model compared with PIM 2<sup>9</sup>. First, there are three new risk indicator fields in PIM 3 categorized as very high-risk, high-risk, and low-risk groups. The very high-risk category included patients with cardiac arrest preceding ICU admission, severe combined immune deficiencies, leukemia or lymphoma after first induction therapy, patients who were bone marrow transplant recipients, and patients with liver failure. All these patients were previously categorized as being at high-risk according to PIM 2. In our study, 237 patients who were categorized as being at high-risk according to PIM 2 were subdivided into high-risk ( $n = 136$ ) and very high-risk ( $n = 101$ ) categories according to PIM 3. Another 19 patients were newly classified into the low-risk category using PIM 3; most of them had seizures, which are not categorized into a risk classification using PIM 2. Second, only the most severe risk factor could be used in the calculation for PIM 3 validation. However, both high-risk and low-risk categories can be used for the calculation of PIM 2 validation. These changes are a significant reason why the PIM 3 is suitable for use.

PIM 3 is the most recently updated model for predicting mortality among patients admitted to the ICU, but it has been criticized for predicting risk of mortality among neonates and adolescents<sup>25</sup>. As previously discussed<sup>1</sup>, the

PIM score has a possible limitation in measuring blood pressure in neonates. Normal blood pressure in neonates is very different depending on body weight. Our study did not include neonates less than 1 month old; however, we categorized patients based on age: infants, preschoolers, schoolchildren, and adolescents. The SMRs were 0.32, 0.99, 1.08, and 0.89 for infants, preschoolers, schoolchildren, and adolescents, respectively. The mortality prediction was less accurate in infants than among the other age categories. These results imply that PIM 3 may be insufficient for predicting mortality not only in neonates but also in infants.

Another possible factor that could affect the precise mortality prediction is the admission route to the ICU: general ward, emergency room, and operating room. Depending on the admission route, the medical environment experienced by the patients before admission is very different. The PIM 3 scores were calculated after ICU admission; therefore, we assumed that the admission route could affect the PIM 3 score. Patients admitted from the general ward may have appeared to be more stable than they actually were because they had received prior medical treatment. Conversely, patients admitted from the emergency room may have appeared to have more severe disease than they actually did. We investigated patients' admission pathways and compared their SMRs. The SMRs were 1.22 and 0.58 for those admitted from the general ward and from the emergency room, respectively. There were more deaths than expected among those admitted from the general ward, and there were fewer deaths than expected among those admitted from the emergency room. The treatment environment according to the hospitalization route may be different, and this could potentially affect PIM scoring.

Our study confirmed the suitability of using PIM 3 through multidimensional statistical analysis. Additionally, we investigated other possible factors that could have affected mortality prediction. Despite the strengths of this study, there were several limitations. First, there was insufficient patient enrollment;

we did not include neonatal, surgical, and cardiac ICU patients. Additionally, this study was conducted at a single center. Multicenter studies that include neonatal, surgical, and cardiac ICU patients will be needed in the future. Second, our study was conducted in a retrospective manner. The authors did their best, but there were patients who were excluded from the study because of incomplete data. There is a need for prospective studies on this topic.

## V. CONCLUSION

PIM 3 is a simple and useful tool for predicting mortality among children admitted to the ICU. Our study showed that PIM 3 showed better prediction of the risk of mortality in children admitted to the ICU in Korea, when compared to PIM 2. Depending on the medical environment, the admission route and age of the patient may affect the PIM 3 scoring system.

## REFERENCES

1. Marcin JP, Wetzel RC. The parsimonious pediatric index of mortality\*. *Pediatr Crit Care Med* 2013;14:718-9.
2. Knaus WA, Wagner DP, Draper EA, Zimmerman JE, Bergner M, Bastos PG, et al. The APACHE III prognostic system. Risk prediction of hospital mortality for critically ill hospitalized adults. *Chest* 1991;100:1619-36.
3. Hargrove J, Nguyen HB. Bench-to-bedside review: outcome predictions for critically ill patients in the emergency department. *Crit Care* 2005;9:376-83.
4. Kirkham FJ, Newton CR, Whitehouse W. Paediatric coma scales. *Dev Med Child Neurol* 2008;50:267-74.
5. Pomerance HH. Nelson Textbook of Pediatrics. *Arch Pediatr Adolesc Med* 2015;151:324.
6. Pollack MM, Yeh TS, Ruttiman UE, Holbrook PR, Fields AI. Evaluation of pediatric intensive care. *Crit Care Med* 1984;12:376-83.
7. Pollack MM, Ruttimann UE, Getson PR. Pediatric risk of mortality (PRISM) score. *Crit Care Med* 1988;16:1110-6.
8. Shann F, Pearson G, Slater A, Wilkinson K. Paediatric index of mortality (PIM): a mortality prediction model for children in intensive care. *Intensive Care Med* 1997;23:201-7.
9. Straney L, Clements A, Parslow RC, Pearson G, Shann F, Alexander J, et al. Paediatric index of mortality 3: an updated model for predicting mortality in pediatric intensive care\*. *Pediatr Crit Care Med* 2013;14:673-81.
10. Pollack MM, Patel KM, Ruttimann UE. PRISM III: an updated Pediatric Risk of Mortality score. *Crit Care Med* 1996;24:743-52.
11. Pollack MM, Cuedon TT, Patel KM, Ruttimann UE, Getson PR, Levetown M. Impact of quality-of-care factors on pediatric intensive care unit mortality. *JAMA* 1994;272:941-6.
12. Slater A, Shann F, Group APS. The suitability of the Pediatric Index of Mortality (PIM), PIM2, the Pediatric Risk of Mortality (PRISM), and PRISM III for monitoring the quality of pediatric intensive care in Australia and New Zealand. *Pediatr Crit Care Med* 2004;5:447-54.
13. Martha VF, Garcia PC, Piva JP, Einloft PR, Bruno F, Rampon V. [Comparison of two prognostic scores (PRISM and PIM) at a pediatric intensive care unit]. *J Pediatr (Rio J)* 2005;81:259-64.
14. Thukral A, Lodha R, Irshad M, Arora NK. Performance of Pediatric Risk of Mortality (PRISM), Pediatric Index of Mortality (PIM), and PIM2 in a pediatric intensive care unit in a developing

- country. *Pediatr Crit Care Med* 2006;7:356-61.
15. Wolfler A, Osello R, Gualino J, Calderini E, Vigna G, Santuz P, et al. The Importance of Mortality Risk Assessment: Validation of the Pediatric Index of Mortality 3 Score. *Pediatr Crit Care Med* 2016;17:251-6.
  16. Jones GD, Thorburn K, Tigg A, Murdoch IA. Preliminary data: PIM vs PRISM in infants and children post cardiac surgery in a UK PICU. *Intensive Care Med* 2000;26:145.
  17. Netto AL, Muniz VM, Zandonade E, Maciel EL, Bortolozzo RN, Costa NF, et al. [Performance of the Pediatric Index of Mortality 2 in a pediatric intensive care unit]. *Rev Bras Ter Intensiva* 2014;26:44-50.
  18. Ciofi degli Atti ML, Cuttini M, Rava L, Rinaldi S, Brusco C, Cogo P, et al. Performance of the pediatric index of mortality 2 (PIM-2) in cardiac and mixed intensive care units in a tertiary children's referral hospital in Italy. *BMC Pediatr* 2013;13:100.
  19. Slater A, Shann F, Pearson G, Paediatric Index of Mortality Study G. PIM2: a revised version of the Paediatric Index of Mortality. *Intensive Care Med* 2003;29:278-85.
  20. Hwang HS, Lee NY, Han SB, Kwak GY, Lee SY, Chung SY, et al. Performance effectiveness of pediatric index of mortality 2 (PIM2) and pediatric risk of mortality III (PRISM III) in pediatric patients with intensive care in single institution: Retrospective study. *Korean Journal of Pediatrics* 2008;51:1158-64.
  21. Zou KH, O'Malley AJ, Mauri L. Receiver-operating characteristic analysis for evaluating diagnostic tests and predictive models. *Circulation* 2007;115:654-7.
  22. Lemeshow S, Hosmer DW, Jr. A review of goodness of fit statistics for use in the development of logistic regression models. *Am J Epidemiol* 1982;115:92-106.
  23. Rapoport J, Teres D, Lemeshow S, Gehlbach S. A method for assessing the clinical performance and cost-effectiveness of intensive care units: a multicenter inception cohort study. *Crit Care Med* 1994;22:1385-91.
  24. Pearson GA, Stickley J, Shann F. Calibration of the paediatric index of mortality in UK paediatric intensive care units. *Arch Dis Child* 2001;84:125-8.
  25. Choi KM, Ng DK, Wong SF, Kwok KL, Chow PY, Chan CH, et al. Assessment of the Pediatric Index of Mortality (PIM) and the Pediatric Risk of Mortality (PRISM) III score for prediction of mortality in a paediatric intensive care unit in Hong Kong. *Hong Kong Med J* 2005;11:97-103.

## ABSTRACT (IN KOREAN)

중환자실에 입실한 한국 소아 환자의 사망률 예측을 위한 Pediatric Index of Mortality 3 의 유효성 평가

<지도교수 손 명 현>

연세대학교 대학원 의학과

정 재 화

**배경:** 소아 중환자 의학이 발달함에 따라 정확한 사망률을 예측하는 것이 중요시 되고 있다. 여러 사망률 예측 모형 중 Pediatric Index of mortality (PIM) 3 은 간편하면서도 대표적인 모형으로, 중환자실 입실시의 10가지 지표를 통해 계산된다.

**연구 목적:** 중환자실에 입실한 한국 소아 환자들의 사망률 예측을 위한 PIM 3 모형의 정확성과 유효성을 평가한다.

**연구 방법:** 2009년 3월에서 2015년 2월까지 세브란스 병원 내과 계 중환자실에 입실한 1개월 이상 18세 미만 소아 환자들을 대상으로 연구를 진행하였다. 수용자 작업특성곡선 (receiver operating characteristic curve, ROC)과 Hosmer-Lemeshow 적합도 검사, 표준화 사망비 (standardized mortality ratio, SMR) 검사를 통해 Pediatric risk of mortality (PRISM) III, PIM 2, PIM 3 모형을 비교하였다.

**결과:** 총 503명의 환자를 분석하였고 각 모형의 ROC 곡선하면적 (area under curve)는 PRISM III = 0.775, PIM 2 = 0.796, PIM 3 = 0.826이었다. Hosmer-Lemeshow 검사 결과 세가지 모형 모두 표본집단에서 적합하였다. PIM 2와 PIM 3모형의 SMR은 각각 0.84과 1.11이었다.

**결론:** 한국 중환자실 소아 환자에서 PIM 3 모형의 유효성을 확인하였고, PIM 2에 비해 더 정확 하였다.

---

핵심되는 말 : PIM 3, 소아, 중환자실, 사망률 예측, 유효성 평가