

Images in
Cardiovascular Medicine



OPEN ACCESS

Received: Jan 20, 2020
Revised: Feb 19, 2020
Accepted: Mar 11, 2020

Correspondence to

Byeong-Keuk Kim, MD, PhD

Division of Cardiology, Department of Internal Medicine, Severance Cardiovascular Hospital, Yonsei University College of Medicine, 250 Seongsan-ro, Seodaemun-gu, Seoul 03722, Korea.

E-mail: kimbk@yuhs.ac

Copyright © 2020. The Korean Society of Cardiology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Yongcheol Kim

<https://orcid.org/0000-0001-5568-4161>

Byeong-Keuk Kim

<https://orcid.org/0000-0003-2493-066X>

Sung-Jin Hong

<https://orcid.org/0000-0003-4893-039X>

Jung-Sun Kim

<https://orcid.org/0000-0003-2263-3274>

Deok-Kyu Cho

<https://orcid.org/0000-0002-3881-411X>

Donghoon Choi

<https://orcid.org/0000-0002-2009-9760>

Myeong-Ki Hong

<https://orcid.org/0000-0002-2090-2031>

Yangsoo Jang

<https://orcid.org/0000-0002-2169-3112>

<https://e-kcj.org>

Successful Culotte Stenting for Unprotected Left Main Trifurcation Disease: Insights from Optical Coherence Tomography

Yongcheol Kim , MD¹, Byeong-Keuk Kim , MD, PhD², Sung-Jin Hong , MD, PhD², Jung-Sun Kim , MD, PhD², Deok-Kyu Cho , MD, PhD¹, Donghoon Choi , MD, PhD¹, Myeong-Ki Hong , MD, PhD², and Yangsoo Jang , MD, PhD²

¹Division of Cardiology, Department of Internal Medicine, Yonsei University College of Medicine and Cardiovascular Center, Yongin Severance Hospital, Yongin, Korea

²Division of Cardiology, Department of Internal Medicine, Severance Cardiovascular Hospital, Yonsei University College of Medicine, Seoul, Korea

A 41-year-old man presenting with new-onset effort angina was referred to our clinic. Angiography revealed a left main trifurcation lesion including significant stenosis in the unprotected left main coronary artery (ULMCA), ostial left anterior descending artery (LAD), ostial left circumflex artery (LCX) with grade 3 collateral flow from the right coronary artery, and diffuse ramus intermedius artery (RI) (**Supplementary Video 1**). After an 8-French extra backup guiding catheter with a side hole was engaged into the left coronary artery via the right femoral approach, plain old balloon angioplasty (POBA) was performed from the ULMCA to LAD, LCX, and RI, respectively. Optical coherence tomography (OCT) demonstrated successful POBA for ostial LCX; thus, percutaneous coronary intervention (PCI) was planned with the 2-stent culotte technique from the ULMCA to the LAD and RI, rather than the crush technique, because 3 strut layers should be avoided on the ostial LCX (**Figure 1A**). A 3.0×38 mm everolimus-eluting stent (EES; Xience Sierra[®], Abbott Vascular) was implanted from the ULMCA to the RI. Subsequently, a 3.5×18 mm EES implantation from the ULMCA to the LAD was achieved with the culotte technique. After stent optimization with kissing balloon inflation and the proximal optimization technique, the final angiography showed no residual stenosis (**Figure 1B** and **Supplementary Video 2**), and post-stenting OCT demonstrated a minimized neo-carina between the ostial LAD and RI (**Figure 1C**) and two strut layers from the distal left main trunk to the shaft including the ostial LCX (**Figure 1D-J**).

The high resolution (10 μm) of OCT can provide detailed post-PCI information, including stent apposition and edge dissections, but there are limited data regarding the OCT-guided PCI with two-stent technique for the bifurcation lesion, especially ULMCA.¹⁾²⁾ This case highlights the feasibility of OCT-guided complex PCI and OCT images of successful culotte technique.




Figure 1. (A) The concept of culotte stenting (line with dots: 2 layers of struts). (B) Final coronary angiography after stenting using the culotte technique. (C) Left main bifurcation area between the ostial LAD and RI. (D, E) Two layers of struts from the RI to LCX. (F) Left main bifurcation area with the ostial LCX (asterisk: guidewire coming from the LCX). (G-J) Two layers of struts in the ULMCA shaft. (arrows: struts of the implanted stent from the ULMCA to RI; arrowheads: struts of implanted stent from the ULMCA to LAD) LAD = left anterior descending artery; LCX = left circumflex artery; ULMCA = unprotected left main coronary artery; RI = ramus intermedius artery.

Conflict of Interest

The authors have no financial conflicts of interest.

Author Contributions

Conceptualization: Kim Y; Data curation: Kim Y, Kim BK; Formal analysis: Kim Y; Investigation: Kim Y; Methodology: Kim BK; Project administration: Kim BK; Supervision: Kim BK, Hong SJ, Kim JS, Cho DK; Visualization: Kim Y; Writing - original draft: Kim Y; Writing - review & editing: Kim BK, Hong SJ, Kim JS, Cho DK, Choi D, Hong MK, Jang Y.

SUPPLEMENTARY MATERIALS

Supplementary Video 1

Initial angiography demonstrating left main trifurcation disease.

[Click here to view](#)

Supplementary Video 2

Final angiography after successful percutaneous coronary intervention with the 2-stent culotte technique.

[Click here to view](#)

REFERENCES

1. Lee CH, Hur SH. Optimization of percutaneous coronary intervention using optical coherence tomography. *Korean Circ J* 2019;49:771-93.
[PUBMED](#) | [CROSSREF](#)
2. Kim Y, Johnson TW, Park SH, et al. Optical coherence tomography findings of non-ST elevation myocardial infarction with multivessel disease. *Korean Circ J* 2020;50:88-90.
[PUBMED](#) | [CROSSREF](#)