

Original Article



Dose-Response Association between Smoking Cessation and Arterial Stiffness: The Cardiovascular and Metabolic Diseases Etiology Research Center (CMERC) Cohort

Ga Bin Lee , BA¹, Jee Seon Shim , PhD², and Hyeon Chang Kim , MD, PhD^{2,3}

¹Department of Public Health, Yonsei University Graduate School, Seoul, Korea

²Cardiovascular and Metabolic Diseases Etiology Research Center, Yonsei University College of Medicine, Seoul, Korea

³Department of Preventive Medicine, Yonsei University College of Medicine, Seoul, Korea



Received: Aug 21, 2019

Revised: Oct 31, 2019

Accepted: Nov 27, 2019

Correspondence to

Hyeon Chang Kim, MD, PhD

Department of Preventive Medicine, Yonsei University College of Medicine, 50 Yonsei-ro, Seodaemun-gu, Seoul 03722, Korea.
E-mail: HCKIM@yuhs.ac

Copyright © 2020. The Korean Society of Cardiology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Ga Bin Lee 

<https://orcid.org/0000-0001-6574-1414>

Jee Seon Shim 

<https://orcid.org/0000-0002-8671-3153>

Hyeon Chang Kim 

<https://orcid.org/0000-0001-7867-1240>

Funding

This work supported by a grant from the Korea Health Technology R&D Project funded by the Ministry of Health and Welfare, Republic of Korea (grant number HI13CO715).

ABSTRACT

Background and Objectives: Cigarette smoking has been reported to be associated with arterial stiffness. However, the clinical relevance of smoking cessation on arterial stiffness is debatable. Thus, we evaluated whether smoking cessation is associated with arterial stiffness.

Methods: A total of 1,169 male participants aged 30–64 years with absence of cardiovascular diseases in 2013–2017 were selected from the Cardiovascular and Metabolic Disease Etiology Research Center cohort study. The subjects were classified into the following 5 groups based on self-reporting: non-smokers, former smokers (<1, 1 to <10, and ≥10, years after cessation), and current smokers. Arterial stiffness was assessed using the augmentation index (AIx). The radial artery AIx was obtained from the peripheral artery waveform. The association was explored cross-sectionally.

Results: The AIx of former smokers did not differ from that of non-smokers ($p=0.089$). However, after former smokers were stratified by duration of smoking cessation, we noted a linear trend according to the smoking status ($p<0.001$). Men who quit smoking <1 year ago showed an elevated AIx ($\beta=3.94$, standard error=1.54, $p=0.011$) as much as ones of current smokers ($\beta=4.39$, standard error=0.74, $p<0.001$), while those who quit more than a decade ago showed an AIx similar to that of non-smokers ($\beta=0.35$, standard error=0.82, $p=0.670$) after controlling covariates.

Conclusions: A dose-response association between smoking cessation and AIx was revealed, which implies the possibility of a reversible effect of smoking cessation on arterial stiffness. Therefore, our findings may motivate current smokers to modify their smoking habits to delay or reverse disease progression.

Keywords: Smoking cessation; Tobacco smoking; Vascular stiffness; Radial artery; Health behavior

Conflict of Interest

The authors have no financial conflicts of interest.

Author Contributions

Conceptualization: Lee GB; Data curation: Shim JS, Kim HC; Formal analysis: Lee GB; Funding acquisition: Kim HC; Investigation: Shim JS, Kim HC; Methodology: Lee GB, Kim HC; Project administration: Lee GB, Shim JS, Kim HC; Resources: Kim HC; Supervision: Shim JS, Kim HC; Validation: Shim JS, Kim HC; Visualization: Lee GB; Writing - original draft: Lee GB; Writing - review & editing: Lee GB, Kim HC.

INTRODUCTION

Tobacco use remains a common habit with increasing numbers worldwide. Although there has been a steady global reduction in the prevalence of daily smoking since 1980, a significant elevation in the number of smokers has been reported concurrently due to population growth.¹⁾ In particular, Korea is a country with a high prevalence of daily smoking and high cigarette consumption.¹⁾ Cigarette smoking is a well-known risk factor of the progression of cardiovascular diseases (CVDs),²⁾ and arterial stiffness may play a underlying pathophysiological mechanism.³⁾

Smoking is a modifiable lifestyle risk factor for CVD unlike the other risk factors, such as age, sex, and ethnicity.⁴⁾ Cigarette smoking is related with arterial stiffness,³⁾ however, the effect of smoking cessation on arterial stiffness has not been clarified. One UK study reported a significant reduction in augmentation index (AIx) in successful quitters 4 weeks after smoking cessation compared to relapsed smokers.⁵⁾ In contrast, no significant difference was investigated in brachial-ankle pulse wave velocity between baseline and 12 months later in a Chinese primary care clinical study.⁶⁾ Another Netherland study reported 2 years of smoking cessation did not significantly affect to a lower progression in intima-media thickness nor to an improved results of dynamic vessel wall properties.⁷⁾ It remains unclear whether smoking cessation is protective for the development of the arterial stiffness.

Therefore, this study aimed to investigate whether there is an association between smoking cessation and arterial stiffness assessed using radial AIx, an estimate of vascular wave reflection and provides a comprehensive assessment of arterial stiffness.

METHODS

The Cardiovascular and Metabolic Diseases Etiology Research Center (CMERC) study constructed 2 prospective cohorts, a cohort consisting of a general population (the CMERC cohort) and high-risk patients (CMERC-HI cohort). This cross-sectional study obtained baseline data from the CMERC cohort. Korean adults aged 30–64 years who were free of myocardial infarction, heart failure, or stroke participated in this study. All participants completed health questionnaires and examinations. The sampling and measurement procedures in the CMERC cohort have been described in detail previously.⁸⁾ Data were collected from 4,060 participants (1,426 men, 2,634 women) between the 2013 and 2017 baseline survey and final analyses were available for 1,169 men. Women were excluded due to the small number of smoking history (n=2,634). Participants taking antihypertensive drugs (n=254) due to its effect on blood pressure and those for whom key variables were missing (n=3) were also excluded.

Information on demographic health-related behaviors and medical history were obtained based on participants' self-reports through a face-to-face interview. Physical activity was evaluated using the International Physical Activity Questionnaire-Short Form - Korean version,⁹⁾ which consists of walking frequency and duration, moderate-intensity activity, vigorous activity, and sedentary activity during the past 7-day. Participants were stratified into 3 groups based on sedentary time (<4, ≥4 and <7, ≥7 hours/day). Participants were classified as current regular alcohol drinkers, former drinkers, or non-drinkers. History of diseases such as diabetes and hypertension was identified through self-reporting of a physician's

diagnosis. Current medication information was also obtained from the participants. People using anti-hypertensive medications were not included for the present study. Physical examinations including measurements of body size, composition, and blood pressure were conducted by trained research personnel. Participants rested for ≥ 5 minutes before the blood pressure measurement, and systolic blood pressure (SBP) and diastolic blood pressure (DBP) were repeatedly measured 3 times at 2-minute intervals in the right arm using an automated oscillometric device (HEM-7080; Omron Health, Matsusaka, Japan). The mean blood pressure of second and third-time measurements was used in the analysis. Pulse pressure (PP) was attained from the difference between SBP and DBP. Biochemical analyses of a fasting blood sample were assessed in the baseline CMERC cohort.

All participants were interviewed using standardized questionnaires to obtain information regarding cigarette smoking. Trained interviewers conducted face-to-face interviews and questionnaire surveys according to the protocol. Details about lifetime smoking experience, current smoking, and smoking amount and duration were obtained from individuals. Smoking status was self-reported as current, former, or non-smoker. Former smokers were categorized into 3 subgroups according to years since smoking cessation: < 1 , 1 to < 10 , and ≥ 10 . The cumulative smoking exposure (i.e., smoking pack-years [SPY]) was calculated from self-reports of cigarette smoking from the questionnaire. Cumulative smoking exposure was estimated as daily packs smoked multiplied by the number of years smoked.

Arterial stiffness was assessed by radial AIx, which was calculated using a non-invasive device (HEM-9000AI; Omron Healthcare Co., Ltd., Kyoto, Japan). The participants were asked to assume a seated position after taking at least a 5-minute rest and remain still and relaxed during all measurements. Radial AIx was calculated as follows: $(\text{Second peak SBP-DBP}) / (\text{First peak SBP-DBP}) \times 100$ (%).¹⁰⁾

Since radial AIx was affected by heart rate, the index was corrected to 75 beats/minute. Waveform data were confirmed by well-trained technicians.

Statistical analyses

We used the analysis of variance for normally distributed continuous variables, the Kruskal-Wallis test for skewed distributed variables, and the χ^2 test for categorical variables to compare the demographic characteristic of individuals according to 5 smoking status groups. Adjusted means of AIx was attained from using analysis of covariance. First, the associations between smoking status (non-smokers [reference group], former smokers, and current smokers) and arterial stiffness, AIx, was investigated by multivariate linear regression. Adjustments were made for age, height, weight, SBP, DBP, diabetes diagnosis, low-density lipoprotein cholesterol, alcohol intake, and sedentary time. Second, subgroup analyses stratified by smoking status according to the duration of smoking cessation (non-smokers [reference group], former smokers [< 1 , 1 to < 10 , ≥ 10 years], and current smokers) were conducted. Third, whether there were multiplicative interactions between former smoking status and cumulative smoking exposure (SPY, continuously) or age (< 55 , ≥ 55) was investigated, and linear trends according to smoking status, from non-smokers to current smokers, were also examined.

We conducted several sensitivity analyses. First, we performed subgroup analyses to investigate cumulative smoking exposure. Former smokers were stratified into 3 groups according to tertile value of smoking pack-years (≤ 8 , > 8 and < 20 , and ≥ 20). Second, to

examine whether our findings differed by the duration of smoking cessation, we applied various cutoffs. Third, to examine whether our findings differed from the participants who took anti-hypertensive medication, we included those participants in the analyses. Fourth, we conducted the analysis also for the female participants (n=2,257) after excluding the ones with taking anti-hypertensive medication. All statistical analyses were performed using SAS software version 9.4 (SAS Institute, Inc., Cary, NC, USA).

Ethics

Informed consent was obtained from all participants. The study protocol was approved by the Institutional Review Board of Severance Hospital (YUIRB-4-2013-0661).

RESULTS

The general characteristics of the study participants stratified by smoking status are presented in **Table 1**. The mean age of the former smokers (52.1 years) was higher than that of the non-smokers (47.8 years) and current smokers (45.7 years). Former smokers tended to have higher SBP, DBP, PP, and AIx values than the current and non-smokers. In addition, there was a linear trend of AIx according to smoking status.

Table 1. Descriptive characteristics of study participants

Variables	Smoking status					p value	p for trend
	Non-smoker (n=271)	Former smokers, duration of cessation (years)			Current smoker (n=420)		
		<1 (n=42)	1 to <10 (n=157)	≥10 (n=279)			
Age (years)	47.8±11.1	45.1±10.6	48.0±9.8	55.4±7.3	45.7±9.8	<0.001	0.873
BMI (kg/m ²)	24.4±2.8	25.6±3.1	24.9±2.6	24.3±2.5	24.9±3.2	0.004	0.139
Waist (cm)	84.6±7.4	88.3±8.0	86.8±7.5	85.7±6.9	86.9±8.5	0.001	0.003
SBP (mmHg)*	122.4±12.7	125.8±14.5	125.4±12.0	126.3±14.8	123.8±13.9	0.010	0.136
DBP (mmHg)*	79.4±9.8	82.1±11.0	81.4±9.7	81.3±9.6	80.2±10.8	0.095	0.346
Pule pressure (mmHg)	43.0±6.6	43.7±7.5	43.9±6.9	45.0±9.8	43.6±7.9	0.059	0.176
AIx75 (%)	72.2±12.8	75.1±11.5	74.5±11.4	77.1±11.1	75.1±11.2	<0.001	<0.001
Total cholesterol (mg/dL)	194.5±35.1	195.3±34.6	205.2±32.2	196.7±35.9	198.2±33.9	0.036	0.299
HDL cholesterol (mg/dL)	51.9±12.4	50.6±14.4	52.6±12.5	52.7±12.6	50.7±13.1	0.249	0.356
LDL cholesterol (mg/dL)	119.1±32.1	120.0±32.5	125.8±30.7	121.5±32.2	121.6±33.4	0.376	0.425
Triglyceride (mg/dL)	112 (78–162)	133 (71–132)	139 (72–120)	114 (73–135)	146 (99–212)	<0.001	0.001
HbA1c (%)	5.5 (5.2–5.7)	5.5 (5.3–5.8)	5.5 (5.2–5.6)	5.6 (5.1–5.6)	5.5 (5.3–5.8)	0.007	0.001
Fasting insulin (uIU/mL)	8.3 (6.3–10.0)	10.0 (6.3–9.8)	9.3 (6.4–9.6)	8.1 (6.5–9.3)	8.5 (6.5–11.0)	0.001	0.660
Fasting glucose (mg/dL)	88 (83–94)	89 (81–92)	92 (80–91)	92 (79–90)	90 (83–98)	<0.001	0.001
Diabetes mellitus (%)	8 (3)	1 (2)	5 (3)	10 (4)	9 (2)	0.797	0.639
Hypertension [†] (%)	12 (4)	2 (5)	6 (4)	22 (8)	28 (7)	0.325	0.150
Physical activity (days/week)						0.394	
None	176 (65)	33 (80)	105 (67)	176 (63)	295 (70)		
≤3	71 (26)	6 (15)	37 (24)	69 (25)	90 (21)		
>3	24 (9)	3 (7)	15 (10)	34 (12)	35 (8)		
Drinking						<0.001	
Never	43 (16)	4 (10)	14 (9)	21 (8)	22 (5)		
Formerly	22 (8)	1 (2)	4 (3)	33 (12)	4 (1)		
Currently	206 (76)	37 (90)	139 (89)	225 (81)	394 (94)		
Sedentary time (hours/day)						0.854	
≤4	77 (28)	13 (32)	41 (26)	81 (29)	132 (31)		
4–7	72 (27)	11 (27)	52 (33)	79 (28)	113 (27)		
>7	122 (45)	18 (44)	64 (41)	119 (43)	174 (41)		

Values are presented as mean±standard deviation or number (%).

AIx75 = augmentation index adjusted for 75 beat/minute; BMI = body mass index; DBP = diastolic blood pressure; HbA1c = hemoglobin A1c; HDL = high density lipoprotein; LDL = low density lipoprotein; SBP = systolic blood pressure.

*Obtained from upper-arm blood pressure; [†]Hypertensive patients but not taking anti-hypertensive medication.

Table 2 presents the association between smoking status and arterial stiffness. Current smokers showed significantly higher AIx ($\beta=4.34$, standard error=0.74, $p<0.001$) compared to non-smokers, but not in former smokers ($\beta=1.23$, standard error=0.72, $p=0.089$). After considering duration of smoking cessation, the altered association was investigated between former cigarette smoking and arterial stiffness. A significant increase in AIx in individuals who quit smoking less than 1-year prior ($\beta=3.94$, standard error=1.54, $p=0.011$) and those with a duration of 1–9 years ($\beta=1.87$, standard error=0.94, $p=0.047$) compared to non-smokers. In addition, a linear trend of AIx according to smoking status considering the duration of smoking cessation was found ($p<0.001$). **Figure 1** shows the adjusted mean AIx values according to the smoking status. There was a dose-response elevation of AIx that those who quit smoking for a shorter duration showed higher values. There were no significant multiplicative interactions between former smoking status and smoking pack-years ($p=0.101$) or age ($p=0.366$). In sensitivity analyses using cumulative smoking exposure, different durations of smoking cessation categorized by former smoking status, and including subjects taking anti-hypertensive medication, similar results were obtained (**Supplementary Figure 1**). In addition, the results of the female participants mostly aligned with those of the primary analyses except that former smokers less than a year showed a lower AIx value compared to non-smokers (**Supplementary Table 1**).

Table 2. Association between smoking status and augmentation index

Smoking status	Number	Age adjusted model		Age and BP adjusted model		Fully adjusted model*	
		β (SE)	p value	β (SE)	p value	β (SE)	p value
Non	271	Reference		Reference		Reference	
Former	478	1.65 (0.80)	0.039	0.99 (0.76)	0.191	1.23 (0.72)	0.089
Current	420	4.05 (0.81)	<0.001	3.80 (0.77)	<0.001	4.34 (0.74)	<0.001
Non	271	Reference		Reference		Reference	
Former, ≥ 10 years quitting	279	0.82 (0.91)	0.374	0.21 (0.87)	0.810	0.35 (0.82)	0.670
Former, 1 to <10 years quitting	157	2.21 (1.04)	0.034	1.54 (0.98)	0.119	1.87 (0.94)	0.047
Former, <1 year quitting	42	4.27 (1.72)	0.011	3.37 (1.63)	0.039	3.94 (1.54)	0.011
Current	420	4.08 (0.81)	<0.001	3.83 (0.77)	<0.001	4.39 (0.74)	<0.001
p for linear trend		<0.001		<0.001		<0.001	

BP = blood pressure; SE = standard error.

*Fully adjusted model was adjusted for age, systolic blood pressure, diastolic blood pressure, height, weight, diabetes diagnosis, low-density lipoprotein cholesterol, alcohol intake, and sedentary time.




Figure 1. Adjusted means of augmentation index according to smoking status. Adjusted for age, systolic blood pressure, diastolic blood pressure, height, weight, diabetes diagnosis, low-density lipoprotein cholesterol, alcohol intake, and sedentary time.

AIx75 = augmentation index adjusted for 75 beat/minute.

*Statistically significant compared to non-smokers ($p<0.05$).

DISCUSSION

The present study investigated that the AIx of former smokers did not differ significantly from that of non-smokers. However, the dose-response association between smoking cessation and AIx was revealed after considering the duration of smoking cessation. The shorter the duration after smoking cessation, the higher were the AIx values. In particular, participants who quit smoking less than 1-year prior showed similar AIx values to those of current smokers, while those who quit smoking more than a decade prior showed AIx values similar to those of non-smokers.

Arterial stiffness improved in former smokers according to cessation duration. In other words, former smokers who abstained from smoking for >10 years showed a reduction in arterial stiffness to the level of non-smokers, pointing toward improved vascular function. Although it is hard to conclude the definite duration of smoking cessation needed to reverse arterial stiffness, the longer duration of cessation seemed to benefit the arterial stiffness in our study. In our sensitivity analyses, various cutoffs of smoking cessation years were applied and ≥ 10 years of smoking cessation found to be the most similar level of AIx compared with that of non-smokers among various cutoff values. Our finding that former smokers with a longer duration of smoking cessation had a similar level of arterial stiffness compared to non-smokers is consistent with previous studies. According to the cross-sectional study of hypertensive patients, the duration of smoking cessation was significantly related to improvements in arterial stiffness including pulse wave velocity, AIx, and transit time.¹¹⁾ Another UK study of 20 patients who participated in the smoking clinic at a hospital found that successful smoking quitters had significant reductions in AIx after 4 weeks of tobacco smoking abstinence compared to relapsed smokers.⁵⁾

Arterial stiffness can be determined by 3 components of the arterial wall: first, quantitative structural components such as the proportion of elastin and collagen; second, qualitative structural components such as cross-linking of glycated collagen; and third, functional components such as vasoconstrictor tone exercised by smooth muscle cells.⁴⁾ Smoking can affect every physiological aspect of arterial stiffening through many potential mechanisms, such as endothelial dysfunction, increased oxidative stress, and inflammation.²⁾ A relevant mechanism related to the reversible smoking cessation effect is the autonomic nervous system (ANS) activated by smoking exposure.¹²⁾ The elevated ANS is correlated with increased blood pressure and peripheral resistance, a result of the concurrent interaction with the weakened baroreflex sensitivity that accompanies aging.¹³⁾ Through acute smoking exposure, nicotine leads to a predominance of sympathetic nerve activity¹²⁾ that can contribute to the risk of peripheral vasoconstriction and an accelerated heart rate.¹⁴⁾ Several studies in agreement with our result reported the possibility of a reversible effect of smoking cessation in relation to the sympathetic nervous system. Chronic habitual smokers¹⁵⁾ and heavy smokers¹⁶⁾ showed improved results in heart rate variability, a measure of the interplay between the sympathetic and parasympathetic nervous systems, within 7 days of smoking cessation. In addition, the sympathetic nerve activity patterns of former smokers were not distinguished from those of non-smokers, which had an implication that smoking effect may be reversible to the normalized sympathetic nerve activity.¹⁷⁾

The changeable effect of smoking cessation on arterial stiffness may also be explained by the role of nitric oxide synthesis related to the sympathetic nervous system.¹⁸⁾ Impaired nitric oxide synthesis is reportedly a detrimental factor in the regulation of vasomotor tone and

blood pressure¹⁹⁾ through a loss of vasodilator tone at the vascular smooth muscle cells and by progressing central neural vasoconstrictor outflow.²⁰⁾ The central action of nitric oxide, reducing sympathetic vasoconstrictor outflow to the sympathetic neurons, and peripheral action of nitric oxide, directly decreasing peripheral vascular smooth muscle tone, contribute to vasodilation or decreased arterial stiffness.²⁰⁾ One study supports the fact that the effect of smoking cessation is correlated with a normalized nitric oxide level. By achieving smoking cessation, the plasma nitric oxide concentration in former smokers was not significantly different from that of the non-smokers.²¹⁾ Another study reported that the elevated oxidative stress assessed by the level of nitric oxide gradually return to normal after a long period of smoking cessation.²²⁾ In addition, several studies reported the interactions between nitric oxide and cardiac autonomic control in animal data²³⁾ or human data.²⁴⁾ Nitric oxide appears to work as a sympatholytic agent,²³⁾ attenuating central sympathetic outflow and lowering cardiac responses to sympathetic stimulation.²⁵⁾ Considering the moderating role of nitric oxide on the sympathetic nervous system, it is reasonable to assume that there might be a reversible effect of smoking cessation on arterial stiffness through an ANS mechanism.

However, there are several limitations should be considered. First, as in all cross-sectional studies, we cannot conclude that there is a causal effect of smoking cessation on AIx. Unmeasured factors such as environmental smoking may be responsible for this relationship.²⁶⁾ Second, we utilized the radial AIx, a surrogate measurement of wave reflection. there are some data supporting that radial AIx reflects vascular stiffening and aging,¹⁰⁾ it may not be considered as a standard measurement of arterial stiffness.²⁷⁾ Third, smoking information including current status, duration of smoking cessation, and the number of cigarettes smoked were self-reported, which might lead to some misclassification bias. However, the validity of self-reported smoking status had been proven through many studies.²⁸⁾²⁹⁾ Fourth, the interpretation of results in this study is limited to male participants. However, in spite of the limited statistic power, we found a similar trend of results with primary analyses in sensitivity analyses with female participants. Fifth, the analyses were based on small numbers of former smokers who quit smoking more than 10 years prior, which limited the power of our statistical analysis. Finally, we found no study, in the literature, evaluating the simultaneous changes on nitric oxide synthesis and sympathetic nervous system, and this could limit our explanation of possible mechanisms.

We found a significant dose-response association between the duration of smoking cessation and AIx in male subjects recruited from the general population. Therefore, our findings may help inspire current smokers to modify their smoking habits to delay or reverse disease progression. Nonetheless, the effect of smoking cessation on arterial stiffness should be further evaluated in longitudinal studies in the future.

ACKNOWLEDGMENTS

The authors appreciate all cohort members for participating in the study and members of the research staff for their passion.

SUPPLEMENTARY MATERIALS

Supplementary Table 1

Association between smoking cessation and augmentation index in women

[Click here to view](#)

Supplementary Figure 1

Association between smoking cessation and augmentation index according to cumulative smoking exposure, different cutoffs of former smokers, and including subjects taking anti-hypertensive medication. Data are presented as β (95% confidence interval) of subjects. Fully adjusted model was adjusted for age, systolic blood pressure, diastolic blood pressure, height, weight, diabetes diagnosis, low-density lipoprotein cholesterol, alcohol intake, and sedentary time.

[Click here to view](#)

REFERENCES

1. Ng M, Freeman MK, Fleming TD, et al. Smoking prevalence and cigarette consumption in 187 countries, 1980–2012. *JAMA* 2014;311:183-92.
[PUBMED](#) | [CROSSREF](#)
2. Ambrose JA, Barua RS. The pathophysiology of cigarette smoking and cardiovascular disease: an update. *J Am Coll Cardiol* 2004;43:1731-7.
[PUBMED](#) | [CROSSREF](#)
3. McGill HC Jr. The cardiovascular pathology of smoking. *Am Heart J* 1988;115:250-7.
[PUBMED](#) | [CROSSREF](#)
4. Tanaka H, Safar ME. Influence of lifestyle modification on arterial stiffness and wave reflections. *Am J Hypertens* 2005;18:137-44.
[PUBMED](#) | [CROSSREF](#)
5. Rehill N, Beck CR, Yeo KR, Yeo WW. The effect of chronic tobacco smoking on arterial stiffness. *Br J Clin Pharmacol* 2006;61:767-73.
[PUBMED](#) | [CROSSREF](#)
6. Wei YJ, Liu HL, Liu B, Zhang L, Jin ZG. Impact of smoking and smoking cessation on arterial stiffness in healthy participants. *Angiology* 2013;64:273-80.
[PUBMED](#) | [CROSSREF](#)
7. van den Berkmortel FW, Wollersheim H, van Langen H, Smilde TJ, den Arend J, Thien T. Two years of smoking cessation does not reduce arterial wall thickness and stiffness. *Neth J Med* 2004;62:235-41.
[PUBMED](#)
8. Shim JS, Song BM, Lee JH, et al. Cardiovascular and Metabolic Diseases Etiology Research Center (CMERC) cohort: study protocol and results of the first 3 years of enrollment. *Epidemiol Health* 2017;39:e2017016.
[PUBMED](#) | [CROSSREF](#)
9. Chun MY. Validity and reliability of Korean version of international physical activity questionnaire short form in the elderly. *Korean J Fam Med* 2012;33:144-51.
[PUBMED](#) | [CROSSREF](#)
10. Kohara K, Tabara Y, Oshiumi A, Miyawaki Y, Kobayashi T, Miki T. Radial augmentation index: a useful and easily obtainable parameter for vascular aging. *Am J Hypertens* 2005;18:11S-14S.
[PUBMED](#) | [CROSSREF](#)
11. Jatoi NA, Jerrard-Dunne P, Feely J, Mahmud A. Impact of smoking and smoking cessation on arterial stiffness and aortic wave reflection in hypertension. *Hypertension* 2007;49:981-5.
[PUBMED](#) | [CROSSREF](#)
12. Middlekauff HR, Park J, Moheimani RS. Adverse effects of cigarette and noncigarette smoke exposure on the autonomic nervous system: mechanisms and implications for cardiovascular risk. *J Am Coll Cardiol* 2014;64:1740-50.
[PUBMED](#) | [CROSSREF](#)

13. Sun Z. Aging, arterial stiffness, and hypertension. *Hypertension* 2015;65:252-6.
[PUBMED](#) | [CROSSREF](#)
14. Oren S, Isakov I, Goltzman B, et al. The influence of smoking cessation on hemodynamics and arterial compliance. *Angiology* 2006;57:564-8.
[PUBMED](#) | [CROSSREF](#)
15. Minami J, Ishimitsu T, Matsuoka H. Effects of smoking cessation on blood pressure and heart rate variability in habitual smokers. *Hypertension* 1999;33:586-90.
[PUBMED](#) | [CROSSREF](#)
16. Yotsukura M, Koide Y, Fujii K, et al. Heart rate variability during the first month of smoking cessation. *Am Heart J* 1998;135:1004-9.
[PUBMED](#) | [CROSSREF](#)
17. Middlekauff HR, Park J, Agrawal H, Gornbein JA. Abnormal sympathetic nerve activity in women exposed to cigarette smoke: a potential mechanism to explain increased cardiac risk. *Am J Physiol Heart Circ Physiol* 2013;305:H1560-7.
[PUBMED](#) | [CROSSREF](#)
18. Scherrer U, Sartori C. Defective nitric oxide synthesis: a link between metabolic insulin resistance, sympathetic overactivity and cardiovascular morbidity. *Eur J Endocrinol* 2000;142:315-23.
[PUBMED](#) | [CROSSREF](#)
19. Haynes WG, Noon JP, Walker BR, Webb DJ. Inhibition of nitric oxide synthesis increases blood pressure in healthy humans. *J Hypertens* 1993;11:1375-80.
[PUBMED](#) | [CROSSREF](#)
20. Sartori C, Lepori M, Scherrer U. Interaction between nitric oxide and the cholinergic and sympathetic nervous system in cardiovascular control in humans. *Pharmacol Ther* 2005;106:209-20.
[PUBMED](#) | [CROSSREF](#)
21. Node K, Kitakaze M, Yoshikawa H, Kosaka H, Hori M. Reversible reduction in plasma concentration of nitric oxide induced by cigarette smoking in young adults. *Am J Cardiol* 1997;79:1538-41.
[PUBMED](#) | [CROSSREF](#)
22. Zhou JF, Yan XF, Guo FZ, Sun NY, Qian ZJ, Ding DY. Effects of cigarette smoking and smoking cessation on plasma constituents and enzyme activities related to oxidative stress. *Biomed Environ Sci* 2000;13:44-55.
[PUBMED](#)
23. Sakuma I, Togashi H, Yoshioka M, et al. NG-methyl-L-arginine, an inhibitor of L-arginine-derived nitric oxide synthesis, stimulates renal sympathetic nerve activity in vivo. A role for nitric oxide in the central regulation of sympathetic tone? *Circ Res* 1992;70:607-11.
[PUBMED](#) | [CROSSREF](#)
24. Chowdhary S, Vaile JC, Fletcher J, Ross HF, Coote JH, Townend JN. Nitric oxide and cardiac autonomic control in humans. *Hypertension* 2000;36:264-9.
[PUBMED](#) | [CROSSREF](#)
25. Balligand JL, Kelly RA, Marsden PA, Smith TW, Michel T. Control of cardiac muscle cell function by an endogenous nitric oxide signaling system. *Proc Natl Acad Sci U S A* 1993;90:347-51.
[PUBMED](#) | [CROSSREF](#)
26. Howard G, Wagenknecht LE, Burke GL, et al. Cigarette smoking and progression of atherosclerosis: The Atherosclerosis Risk in Communities (ARIC) Study. *JAMA* 1998;279:119-24.
[PUBMED](#) | [CROSSREF](#)
27. Cheng LT, Tang LJ, Cheng L, Huang HY, Wang T. Limitation of the augmentation index for evaluating arterial stiffness. *Hypertens Res* 2007;30:713-22.
[PUBMED](#) | [CROSSREF](#)
28. Wagenknecht LE, Burke GL, Perkins LL, Haley NJ, Friedman GD. Misclassification of smoking status in the CARDIA study: a comparison of self-report with serum cotinine levels. *Am J Public Health* 1992;82:33-6.
[PUBMED](#) | [CROSSREF](#)
29. Vartiainen E, Seppälä T, Lillsunde P, Puska P. Validation of self reported smoking by serum cotinine measurement in a community-based study. *J Epidemiol Community Health* 2002;56:167-70.
[PUBMED](#) | [CROSSREF](#)