



저작자표시-비영리-변경금지 2.0 대한민국

이용자는 아래의 조건을 따르는 경우에 한하여 자유롭게

- 이 저작물을 복제, 배포, 전송, 전시, 공연 및 방송할 수 있습니다.

다음과 같은 조건을 따라야 합니다:



저작자표시. 귀하는 원저작자를 표시하여야 합니다.



비영리. 귀하는 이 저작물을 영리 목적으로 이용할 수 없습니다.



변경금지. 귀하는 이 저작물을 개작, 변형 또는 가공할 수 없습니다.

- 귀하는, 이 저작물의 재이용이나 배포의 경우, 이 저작물에 적용된 이용허락조건을 명확하게 나타내어야 합니다.
- 저작권자로부터 별도의 허가를 받으면 이러한 조건들은 적용되지 않습니다.

저작권법에 따른 이용자의 권리는 위의 내용에 의하여 영향을 받지 않습니다.

이것은 [이용허락규약\(Legal Code\)](#)을 이해하기 쉽게 요약한 것입니다.

[Disclaimer](#)

Effect of intentional restriction of venous return on tissue oxygenation in a porcine model of acute limb ischemia

Wonho Kim

Department of Medicine

The Graduate School, Yonsei University

Effect of intentional restriction of venous return on tissue oxygenation in a porcine model of acute limb ischemia

Wonho Kim

Department of Medicine

The Graduate School, Yonsei University

Effect of intentional restriction of venous return on tissue oxygenation in a porcine model of acute limb ischemia

Directed by Professor Myeong-Ki Hong

The Doctoral Dissertation
submitted to the Department of Medicine,
the Graduate School of Yonsei University
in partial fulfillment of the requirements for the degree
of Doctor of Philosophy

Wonho Kim

June 2021

This certifies that the Doctoral
Dissertation of Wonho Kim is
approved.

Thesis Supervisor: Myeong-Ki Hong

Thesis Committee Member#1: Yangsoo Jang

Thesis Committee Member#2: Donghoon Choi

Thesis Committee Member#3: Chung Mo Nam

Thesis Committee Member#4: Seung-Ho Hur

The Graduate School
Yonsei University

June 2021

ACKNOWLEDGEMENTS

I owe my deepest gratitude to my supervisor, Professor Myeong-Ki Hong for inspiring and guiding me to the enlightening field of cardiology and cardiovascular intervention. His mentorship was paramount in providing key experiences in the process of cardiologic research. I am also grateful for his guidance which helped me pursue my academic career. I also would like to thank heartily Professor Yangsoo Jang who gave me an academic support throughout this study. His help was crucial in proceeding with this basic research study. I thank professors Donghoon Choi, Chung Mo Nam and Seung-Ho Hur for providing the important and constructive comments and advices. Without the on-site help and hands-on experience of Professor Se-Il Park in the laboratory procedures, conducting this research would have been impossible. I also appreciate his prompt and reliable help, and the experiences in problem-solving we have shared.

Most importantly, I am very grateful to my wife, Jiyoung Shin for her love and emotional support throughout this study and during my difficult times. I dedicate this work to my parents who have always stood by my side. Lastly, I would like to share this joy with my daughter, Yumi Kim.

<TABLE OF CONTENTS>

ABSTRACT	9
I. INTRODUCTION	10
II. MATERIALS AND METHODS	12
1. Lesion preparation	12
2. Procedures	12
3. Lower limb ischemia model	14
4. Blood sampling and measurement	15
5. Experimental study protocol	15
6. Statistical analysis	16
III. RESULTS	17
1. There are no significant differences between the values observed at T1-T4 and T5-T7 in the arterial oxygen saturation	17
2. After partial venous occlusion, the venous oxygen saturations (T5-T7) are significantly increased	18
3. The venous oxygen saturations evaluated in the experimental limb and in the control limb are significant over time	19
4. A significant decline after arterial embolization and re-elevation were observed in the mean venous pressure	20
5. Hypoxic damage by arterial embolization is positively associated with treatment effect with venous partial occlusion	21
IV. DISCUSSION	22
V. CONCLUSION	28
REFERENCES	29
ABSTRACT (IN KOREAN)	33

LIST OF FIGURES

Figure 1. Preparation of an animal model of ALI	13
Figure 2. Representative images of the porcine model of ALI	14
Figure 3. Experimental study protocol and blood sampling for evaluating oxygen saturation (right SFA and the right CIV)	16
Figure 4. Blood sampling of the right SFA (A) and right CIV (B) at different moments (From T0 to T7).....	17
Figure 5. Representative image of the venous blood pressure response in the ischemic limb	20
Figure 6. Spearman correlation was performed for bivariate correlations between hypoxic damage by arterial embolization	21

LIST OF TABLES

Table 1. Blood sampling of the SFA and CIV at different moments (From T0 to T7)	18
Table 2. Comparison of the venous oxygen saturations between the experimental limbs with arterial embolization and the control limbs	19

ABSTRACT

Effect of intentional restriction of venous return on tissue oxygenation in a porcine model of acute limb ischemia

Wonho Kim

*Department of Medicine
The Graduate School, Yonsei University*

(Directed by Professor Myeong-Ki Hong)

A sufficient oxygen supply to ischemic limb tissue is the most important requirement for wound healing and limb salvage. We investigated whether partial venous occlusion in the common iliac vein (CIV) causes a further increase of venous oxygenation in a porcine model of acute hindlimb ischemia. In 7 pigs, the model of acute hindlimb ischemia was created with embolization of the common iliac artery. Oxygen saturation was evaluated at baseline (T0), just after the arterial embolization (T1), at 10 minutes (T2), at 20 minutes (T3), and at 40 minutes (T4). Next, an intentional partial venous occlusion was achieved by inflating the vascular balloon at the level of the right CIV. Then, blood sampling was repeated at 5 minutes (T5), at 15 minutes (T6), and at 25 minutes (T7). The arterial oxygen saturation in the right SFA was similar during all phases. In contrast, after arterial embolization, an immediate reduction of venous oxygen saturation was observed (from 85.57 ± 1.72 at T0 to 71.86 ± 7.58 at T4). After the partial venous occlusion, the venous oxygen saturations (T5-T7) were significantly increased, again. The venous oxygen saturations evaluated in the hindlimb ischemia with partial venous occlusion and in the control limb were significantly over time. Venous oxygen saturations in the experimental limbs were higher than those in the control limbs (79.71 ± 4.78 vs 60.00 ± 4.24 at T7, p-value <0.001). Partial venous occlusion results in an increase of venous oxygen saturation in the ischemic limb, while significant changes in venous oxygen saturation are not observed in the control limb. An explanation for this may be that the oxygen consumption in the limb tissue is increased because it gets congested with the partial venous occlusion in the right CIV.

Keywords: oxygenation, acute limb ischemia, partial venous occlusion.

Effect of intentional restriction of venous return on tissue oxygenation in a porcine model of acute limb ischemia

Wonho Kim

*Department of Medicine
The Graduate School, Yonsei University*

(Directed by Professor Myeong-Ki Hong)

I. INTRODUCTION

Peripheral artery disease (PAD) is a cause of significant morbidity, mortality, and disability due to limb loss.¹ In particular, acute limb ischemia (ALI) occurs when there is a sudden decrease in limb perfusion that threatens the viability of the limb and requires urgent treatment to prevent loss of limb.² A lack of oxygen due to an interruption of the blood supply to an acutely occluded limb causes the accumulation of various metabolites, production of reactive oxygen species (ROS) and an inflammatory reaction in conjunction with tissue swelling, threatening the possibility of limb loss and even death.³ Subsequently, a sufficient oxygen supply to the ischemic limb tissue is the most important requirement for the promotion of wound healing and limb salvage. For limb salvage and restoring ambulatory function, therapies targeting ALI are directed primarily towards mechanical revascularization with an endovascular or surgical approach because essentially no effective medical therapies exist. Unfortunately, revascularization is often of limited benefit, and the clinical outcomes of the revascularization are not always satisfactory.^{2,4} Interestingly, previous investigators showed that an intentional narrowing of the coronary sinus lumen elevates the coronary venous pressure, leading to an increase in

capillary and arteriolar dilatation, a lower resistance to blood flow, and the restoration of the normal endocardial/epicardial blood flow ratio in patients with refractory angina.^{5,6} As a consequence of the enhanced blood flow to the ischemic myocardium, this process increases the oxygen supply and its metabolism in ischemic myocardium.

In this study, a porcine model of acute hindlimb ischemia was created by intra-vascular embolization of the common iliac artery (CIA). By using models, we investigated if partial venous occlusion in the common iliac vein (CIV) might cause a further increase of venous oxygenation.

II. MATERIALS AND METHODS

1. Lesion preparation

This study was approved by the Yonsei University Institutional Animal Care and Use Committee. Sexually mature female swine (N=8, 6 to 8 months of age) were obtained from a local vendor. The animals used in this study were miniature pigs originated from Korean Jeju island native (Cronex Co., Ltd., Osong, Republic of Korea). All the animals received humane care in compliance with the Animal Welfare Act and “The Guide for the Care and Use of Laboratory Animals” formulated by the Institute of Laboratory Animal Research.⁷ The animals were housed individually following standard laboratory conditions (temperature: 19-25 °C; humidity: 30-70%; ventilation: 10-15 per hour; light: 150-300 Lux; light cycle: twice per day (8 AM- 8PM); noise: 45 dB) and fed a standard laboratory pellet diet and water ad libitum. A veterinarian dedicated to our laboratory supervised and helped with the study. At the end of this study, animals were euthanized, according to the American Veterinary Medical Association guidelines for the euthanasia of animals. Briefly, under deep anesthesia, the animals received bolus injections of potassium chloride.

2. Procedures

Anesthesia was induced by using intra-muscular injections of ketamine (20 mg/kg) and xylazine (2 mg/kg). After adequate systemic anesthesia was attained, the animals were placed in the supine position under mechanical ventilation, and isoflurane (1-2%) was delivered with a precision vaporizer and a circle absorption breathing system with periodic arterial blood gas monitoring. Cardiovascular monitoring was used during the whole procedure to control the heart rate and blood pressure. After surgical exposure of both carotid arteries

and one carotid vein under sterile conditions, two 8 Fr arterial sheaths and one 9 Fr vein sheath were inserted (Fig 1). Intra-venous heparin (150 units/kg) was injected to maintain an activated clotting time of ≥ 250 seconds. First, a 0.014-inch hydrophilic guidewire was carefully inserted into the left carotid artery and navigated down to the right CIA. An 8 Fr multi-purpose guiding catheter was then advanced over the guidewire into the right CIA. A micro-catheter through the right carotid artery was placed in the right superficial femoral artery (SFA). The role of micro-catheter is to obtain arterial blood sampling in the right SFA when the right CIA is embolized to induce hindlimb ischemia. Second, a 0.014-inch hydrophilic guidewire through the left carotid vein was advanced in the inferior vena cava (IVC) and selectively, down into the right CIV. A 9 Fr multi-purpose guiding catheter was moved down over the guidewire and positioned in the CIV.




Fig. 1. Preparation of an animal model of ALI. After surgical exposure of both carotid arteries (white arrows) and one carotid vein (black arrow) under sterile conditions, two 8 Fr arterial sheaths (white arrows) and one 9 Fr vein sheath (black arrow) were inserted. Note that a micro-catheter (grey arrow) is inserted in the right carotid artery, an 8 Fr multi-purpose guiding catheter in the left carotid artery, and a 9 Fr multi-purpose guiding catheter in the left carotid vein. Abbreviations: ALI, acute limb ischemia.

3. Lower limb ischemia model

Baseline angiography was performed in the right leg (Fig 2). Embolization to induce hindlimb ischemia in the right CIA was performed using an Amplatzer Vascular Plug (AVP; St/ Jude Medical, St. Paul, MN), which acts as an embolic agent by promoting clot formation. To increase the thrombogenicity of the AVP, it was oversized by 30-50% of the estimated diameter of the right CIA. The occlusion time is highly variable depending on the caliber of the vessel, the coagulation status, the flow dynamics, and the size of the AVP. Therefore, repeated angiography was frequently performed to see if complete stasis of the injected contrast was noted.




Fig. 2. Representative images of the porcine model of ALI. The pigs underwent intravascular induction of ALI. Baseline angiography (A), Embolization of the CIA. ALI was generated by the deployment of an Amplatzer Vascular Plug II in the CIA, followed by an angiography to verify occlusion (B), representative angiography after CIA occlusion (C), Baseline venography (D), Venous blood flow restriction with partial balloon inflation (E). Abbreviations: ALI, acute limb ischemia; CIA, common iliac artery.

4. Blood sampling and measurement

Blood gas analysis was performed to measure oxygen saturation, carbon dioxide, bicarbonate, and acid-base balance (pH). A proper blood sample for blood gas analysis consists of a 2 to 3 ml blood specimen collected anaerobically in a 3 ml plastic syringe fitted with a small-bore needle. Heparin was added to the syringe as an anticoagulant. Any air bubbles inadvertently introduced during sampling were promptly evacuated. Mean venous pressure was measured during the study period. All the mean venous pressure values were calculated over a stable period.

5. Experimental study protocol

Blood sampling (Fig 3) for evaluating oxygen saturation was obtained from the right SFA and CIV at baseline (T0), just after the arterial occlusion with embolization (T1), at 10 minutes (T2), at 20 minutes (T3), and at 40 minutes (T4). Next, an intentional partial (90%) venous occlusion was achieved by inflating the vascular balloon at the level of the right CIV. Subsequently, blood sampling was repeated 5 minutes (T5) after the partial venous occlusion, at 15 minutes (T6), and at 25 minutes (T7). The left leg in two pigs served as a control limb; the left CIA was embolized while the left CIV was not occluded. In those legs, blood sampling was obtained 40 minutes after the left CIA occlusion in the left CIV with the same method as above (T5-1, T6-1, and T7-1). Hypoxic damage by arterial embolization was defined as T4 minus T0 in vein saturation, and treatment effect with partial venous occlusion was defined as T7 minus T4 in vein saturation, respectively.




Fig. 3. Experimental study protocol and blood sampling for evaluating oxygen saturation (right SFA and the right CIV). Blood sampling for evaluating oxygen saturation was obtained from the right SFA and the right CIV at baseline (T0), just after the arterial occlusion with embolization (T1), at 10 minutes (T2), at 20 minutes (T3), and at 40 minutes (T4). Next, an intentional partial (90%) venous occlusion was achieved by inflating the vascular balloon at the level of the right CIV. Subsequently, blood sampling was repeated at 5 minutes (T5) after the partial venous occlusion, at 15 minutes (T6), and at 25 minutes (T7). The left leg in two pigs served as a control limb; the left CIA was embolized while the left CIV was not occluded. In those legs, blood sampling was obtained 40 minutes after the left CIA occlusion in the left CIV with the same method as above (T5-1, T6-1, and T7-1). Abbreviations: CIA, common iliac artery; SFA, superficial femoral artery; CIV, common iliac vein.

6. Statistical analysis

Dichotomous data were reported as a number and percentage. Continuous data were reported as the mean and the standard deviation. Student t test was used to analyze continuous data and a chi-square test for dichotomous data. Arterial and venous oxygen saturations at different momentums were submitted to a repeated measures analysis of variance (ANOVA). All tests were two-sided, and statistical significance was defined as a p-value of less than 0.05. The analyses were performed using the SPSS version 20 software (SPSS Inc., Chicago, Illinois).

III. RESULTS

1. There are no significant differences between the values observed at T1-T4 and T5-T7 in the arterial oxygen saturation.

The model of acute hindlimb ischemia was successfully created in all 8 pigs. One pig died of sudden cardiac death due to iatrogenic catheter induced ventricular fibrillation. The seven surviving pigs developed acute hindlimb ischemia. The arterial oxygen saturation in the right SFA was similar during all phases. There were no significant differences between the values observed at T1-T4 and T5-T7 (Fig 4A and Table 1).




Fig. 4. Blood sampling of the right SFA (A) and right CIV (B) at different moments (From T0 to T7). Effect of partial venous occlusion on venous oxygen saturation over time (from T4 to T7) (C). Abbreviations: SFA, superficial femoral artery; CIV, common iliac vein.

Table 1. Blood sampling of the SFA and CIV at different moments (From T0 to T7).

SFA (percent)		CIV (Percent)	
T0	100	T0	85.57 ± 1.72
T1	98.43 ± 1.27	T1	75.43 ± 8.44
T2	98.29 ± 1.11	T2	71.71 ± 5.93
T3	97.57 ± 1.51	T3	72.57 ± 6.35
T4	97.71 ± 1.97	T4	71.86 ± 7.58
T5	98.29 ± 2.13	T5	74.14 ± 9.56
T6	98.29 ± 1.60	T6	79.29 ± 4.82
T7	98.57 ± 0.97	T7	79.71 ± 4.78

Values are n (%) or mean ± SD, as appropriate. Abbreviations: SFA, supercifical femoral artery; CIV, common iliac vein.FA and CIV at different moments (From T0 to T7).

2. After partial venous occlusion, the venous oxygen saturations (T5-T7) are significantly increased.

After arterial embolization, an immediate reduction of venous oxygen saturation was observed (from 85.57 ± 1.72 at T0 to 71.86 ± 7.58 at T4). After partial venous occlusion, interestingly, the venous oxygen saturations (T5-T7) were significantly increased, again. Repeated measures ANOVA show a significant difference after the partial venous occlusion concerning venous oxygen saturation (p-value < 0.001) (Fig 4B and Table 1).

3. The venous oxygen saturations evaluated in the experimental limb and in the control limb are significant over time.

The venous oxygen saturations evaluated in the hindlimb ischemia with partial venous occlusion and in the control limb (without partial venous occlusion) were significant over time. Venous oxygen saturations in the experimental limb were higher than those in the control limb (79.28 ± 4.82 vs 59.00 ± 2.82 at T6, p-value <0.001 , 79.71 ± 4.78 vs 60.00 ± 4.24 at T7, p-value <0.001) (Table 2). The largest changes in venous oxygen saturation during partial venous occlusion were observed at T6 (from 74.14 ± 9.56 at T5 to 79.28 ± 4.82 at T6). Overall, venous oxygenation was significantly increased after partial venous occlusion (from 74.14 ± 9.56 at T5 to 79.71 ± 4.78 at T7, p-value <0.001) (Fig 4C), while its largest change was observed at T6 (from 74.14 ± 9.56 at T5 to 79.28 ± 4.82 at T6).

Table. 2. Comparison of the venous oxygen saturations between the experimental limbs with arterial embolization and the control limbs.

	Experimental limb	Control limb	p-value
T5	74.14 ± 9.56	57.50 ± 7.77	0.061
T6	79.28 ± 4.82	59.00 ± 2.82	< 0.001
T7	79.71 ± 4.78	60.00 ± 4.24	< 0.001

Values are n (%) or mean \pm SD, as appropriate.

4. A significant decline after arterial embolization and re-elevation were observed in the mean venous pressure.

Fig 5 shows a representative image of the mean venous blood pressure response during the study period. A significant decline after arterial embolization (from 18.9 ± 1.3 at T0 to 3.0 ± 1.2 mmHg at T1, p-value <0.001) and re-elevation were observed (from 3.0 ± 1.2 at T1 to 7.1 ± 1.3 mmHg at T7, p-value <0.001) in the mean venous pressure.



Fig. 5. Representative image of the venous blood pressure response in the ischemic limb. White arrows show a recording of the mean venous pressure, at baseline (T0), just after the arterial occlusion (T1), and 25 minutes after partial vein occlusion (T7). A decline after arterial occlusion (B) and an elevation after partial vein occlusion were observed in the mean venous pressure (C).

5. Hypoxic damage by arterial embolization is positively associated with treatment effect with venous partial occlusion.

Fig 6 shows that hypoxic damage by arterial embolization is positively associated with treatment effect with venous partial occlusion ($r = 0.788$, $P\text{-value}=0.035$). This strong positive correlation might be interpreted that the more severe hypoxic damage, the greater would be the treatment effect with partial venous occlusion. On the other hand, we did not detect a significant change in the level of carbon dioxide, bicarbonate or pH during the study period.




Fig. 6. Spearman correlation was performed for bivariate correlations between hypoxic damage by arterial embolization (defined as T4 minus T0 in vein saturation) and treatment effect with partial venous occlusion (defined as T7 minus T4 in vein saturation) to establish a relationship.

IV. DISCUSSION

In this porcine model of acute hindlimb ischemia, arterial embolization decreases venous oxygen saturation over time. However, intentional partial venous occlusion results in an increase of venous oxygen saturation in the ischemic limb, while significant changes in venous oxygen saturation are not observed in the control limb. An explanation for this may be that the oxygen consumption in the limb tissue is increased because it gets congested with the partial venous occlusion in the right CIV. Long term effects of the partial venous occlusion are beyond the spectrum of this study. However, our result may lead to investigating an optional therapy targeted at increasing tissue oxygenation, which could be used alone or in combination with mechanical revascularization to improve clinical care.

Mechanical revascularization is a fundamental strategy for limb preservation. However, this strategy is occasionally unsuccessful at improving limb salvage. Investigational approaches derived from animal models or preclinical studies, like stem cell-based or gene therapies, have also been largely ineffective.^{8,9} Because there is still much debate on the best clinical and therapeutic strategies, it remains unclear just how good is good enough when it comes to increasing the blood supply to an ischemic limb. In clinical scenarios, there are patients for whom the revascularization method is not feasible for them. The intentional partial venous restriction in the CIV for oxygenation in the ischemic limb may assist in wound healing and decrease the risk of major amputation.

Our experimental idea got inspiration from Beck's study, which demonstrated in a dog model of acute coronary artery ligation that the partial narrowing of the coronary sinus is associated with a reduction in the infarct size and mortality by a doubled degree of retrograde backflow from the distal stump of the occluded coronary artery.¹⁰ Coronary sinus narrowing leads to the development of an upstream pressure gradient that results in the redistribution

of blood from the less ischemic epicardium to the ischemic endocardium.¹¹

An increase in oxygenated venous blood might be seen in the ischemic limb with both partial and complete venous occlusion. Enriched oxygen in tissue is considered beneficial to the metabolism in the muscle. The obstruction of arterial blood flow causes impaired oxygen delivery, tissue hypoxia and leads to the dysfunction of the electron transport chain in mitochondria. That induces anaerobic metabolism in cells, and the retention of lactic acid may lead to metabolic acidosis. Because tissue oxygen extraction/utilization is impaired in the ischemic limb, an increase in oxygen consumption in the ischemic tissue might not decrease the venous oxygen saturation.¹² Partial venous occlusion in the CIV induces local stasis of highly oxygenated blood in the ischemic area. It causes a slight venous pressure elevation in the vein (Fig 5), and the sustaining high oxygen level would be helpful for oxygen uptake by ischemic limb tissue, restoring cell metabolism and limb perfusion.

A total venous occlusion may develop a fatal deep vein thrombosis (DVT); there is a risk that a venous thrombosis starts as a partial venous occlusion and progresses over time. In animal experiments, the methods of creating a DVT model are mainly based on flow impairment induced by surgical ligation, endovascular balloon occlusion, or the assistance of specifically designed endovascular devices.¹³ However, venous stasis alone, unless prolonged, does not consistently induce venous thrombosis in an intact vein.¹⁴ Nevertheless, we often check if venous blood flow is patent during balloon occlusion to avoid a total venous thrombosis. On the other hand, there is concern that a venous ulcer or venous stasis related life-threatening condition may occur due to venous congestion by an intentional partial venous occlusion, causing pressure in the veins to increase. However, those catastrophic events are unlikely to develop because the intentional partial venous occlusion would be applied, temporarily.

The majority of studies evaluating susceptibility to tissue necrosis in hindlimb ischemia have focused on vascular contributions, such as angiogenesis

and collateral artery growth.^{15,16} In those studies, limb salvage is logically assumed to be dependent on perfusion and subsequent oxygen delivery. A critical component of limb tissue regeneration after ischemic injury is a sufficient oxygen concentration. On the other hand, oxygen has been used as a target of non-invasive measurements of tissue perfusion in limb ischemia. In clinical practice, the transcutaneous measurement of oxygen partial pressure is a good tool to assess the perfusion status of the foot in limb ischemia.¹⁷ One preliminary study showed that hyperbaric oxygen therapy has anti-microbial and neovascularization effects in patients with successful endovascular revascularization for the affected limb.¹⁸

The most common model of PAD is the hindlimb ischemia model, which is most frequently performed in small animals like rodents and rabbits.^{19,20} Generally, the model entails surgical ligation of the femoral artery and its intervening side-branches followed by excision of the vessel, resulting in occlusion of the blood flow and induction of ALI. One might argue that the most important weakness in this study is to use a model of ALI. A criticism of this model is that ALI does not represent the natural history of the most common PAD with gradual development of progressive atherosclerotic lesions. A chronic limb ischemic model with ischemic changes similar to those seen in patients must be ideal, but this work needs several weeks after induction of hindlimb ischemia. They are logistically more complex to handle and not cost effective.²¹ Compared to small animals, pigs have a size and physiology that more closely mimic humans.

Meanwhile, previous studies with the porcine hindlimb ischemic model have been limited by rapid collateralization.^{22,23} Surgical CIA ligation would completely exclude the possibility of collateralization to the ischemic hindlimb and make a more durable and reproducible ischemic state in the porcine model.²¹ However, the human peripheral limb vascular system has numerous collaterals and inter-individual disparities. Generally, irreversible tissue damage

in ALI usually occurs within 6 hours after complete arterial occlusion. Although endovascular CIA embolization leads to a stronger reduction in the total amount of blood flow into the ipsilateral limb, it would seem barely feasible to occlude all collateral branches at the initial stage of ALI. Tissue blood flow and perfusion to the skin or limb muscles are significantly decreased, whereas intraluminal arterial saturation in the occluded artery may not be decreased and still has highly saturated blood. In addition, arterial anatomy of the porcine hind limb is different from that in human peripheral limb. In the pig, the distal abdominal aorta generally trifurcates into the right and left iliac arteries, and the common internal iliac trunk which then bifurcates into the right and left internal iliac arteries.²⁴ Development of collateral vasculature via those internal iliac arteries might be one of the important mechanisms in compensating for arterial occlusion in the porcine limb ischemic model. Our model was induced with arterial embolization in the CIA, not at the level of distal abdominal aorta. Thus, the patent common internal iliac trunk might facilitate the development of collateral blood flow into the ischemic limb. However, we did not actually show the limb perfusion through collaterals, which could be assessed with peripheral angiography, non-invasive imaging modalities, or histological staining of the ischemic tissues. Swine models fulfilling all ALI characteristics, with evaluating the adequacy of limb perfusion through collaterals intraprocedurally, seem to be lacking in the scientific literature. When it comes to further researches following this study, a major challenge may be having technologies available that can show numerous collaterals in the ischemic limb.

Abrupt total occlusion of the CIA, like surgical CIA ligation, is a rare medical scenario and is hardly seen in PAD patients clinically. Considering the mentioned points, our model is likely to show ischemic changes similar to those seen in patients with ALI. We focused on a straight-forward reliable and reproducible endovascular method. Moreover, our method is well equipped for performing our study which is designed to investigate the effect of partial

venous occlusion on the oxygenation in porcine hindlimb ischemia.

It is important to recognize several other limitations of this study. First, our model is based on the non-atherosclerotic and normo-lipidemic hindlimb ischemia model. Because the model is done in a swine without any cardiovascular risk factors or comorbidities, the induction of hindlimb ischemia will never perfectly replicate the pathophysiology of clinical PAD. Although the diseased porcine models provide more similar biological responses, the normo-lipemic porcine artery is still recommended as the choice of the porcine model to evaluate pathophysiologic responses in consensus documents.^{21,25} This study aimed to investigate the effect of partial venous occlusion on tissue oxygenation in porcine hindlimb ischemia. The main target is not atherosclerotic arteries, but the vein. The presence of cardiovascular risk factors or the onset of ischemia may not influence our expected study outcomes. Second, there are concerns about the small number of experimental animals and a relatively short period of observation time. With the current amount of observations, however, there was an obvious trend towards improvement of the venous oxygen saturation over time, following intentional partial venous occlusion in the CIV. Possibly, the inclusion of an additional time point, e.g. 6 hours after the partial venous occlusion, may have revealed more information about the effect of the intentional restriction of venous return on tissue oxygenation in the ischemic limb. Third, partial venous occlusion was created with percutaneous balloon inflation, causing a 90% stenosis artificially in the CIV by visual estimation. The designated percutaneous device, like a coronary sinus reducer, may provide a more accurate way in which to achieve a stable and durable venous blood flow. However, those devices are not yet available, and it is not likely to be cost effective. Future investigators should clarify the question concerning the designated vein narrowing device. In our opinion, our porcine model with percutaneous balloon inflation could prove suitable for evaluating the effect of partial venous occlusion on venous oxygenation. Fourth,

there is a limitation to the use of non-invasive measurements of tissue perfusion such as transcutaneous oxygen pressure, indocyanine green fluorescence angiography, laser Doppler flow-metry, or positron emission tomography (PET). It might be argued that conventional peripheral angiography (macrocirculation) with intraluminal oxygen saturation does not allow reliable assessment of the tissue perfusion and microcirculation, and could, therefore, underestimate the effect of intentional restriction of venous return on tissue oxygenation in our model. At present, the object of this study is to prove the similarity between the mechanism of coronary reducer and that of partial venous occlusion in the CIV. Tissue perfusion imaging for evaluating the effect of partial venous occlusion on tissue oxygenation in the limb ischemia would be an important part at next research. Finally, our study was insufficient to demonstrate an effect of partial venous occlusion on the inflammatory marker, oxidative stress, and acid-base balance in the ischemic limb. However, this study was not designed or powered to look at the impact on those parameters, although those are likely of great interest. The study period may not have been long to detect significant changes.

V. CONCLUSION

Given the limited evidence-based research to date, the clinical efficacy of using a non-revascularization strategy in limb ischemia is still questionable. Our results suggest a potential positive therapeutic effect of partial venous occlusion to induce the enhancement of oxygen uptake in the ischemic limb tissue. The effect of a partial venous occlusion on tissue oxygenation in an ischemic limb may deserve to be investigated in patients for whom revascularization therapy is unsuccessful.

REFERENCES

1. Fowkes FG, Rudan D, Rudan I, Aboyans V, Denenberg JO, McDermott MM, et al. Comparison of global estimates of prevalence and risk factors for peripheral artery disease in 2000 and 2010: a systematic review and analysis. *Lancet* 2013;382:1329-40.
2. Aboyans V, Ricco JB, Bartelink MEL, Björck M, Brodmann M, Cohnert T, et al; ESC Scientific Document Group. 2017 ESC Guidelines on the Diagnosis and Treatment of Peripheral Arterial Diseases, in collaboration with the European Society for Vascular Surgery (ESVS): Document covering atherosclerotic disease of extracranial carotid and vertebral, mesenteric, renal, upper and lower extremity arteries Endorsed by: the European Stroke Organization (ESO) The Task Force for the Diagnosis and Treatment of Peripheral Arterial Diseases of the European Society of Cardiology (ESC) and of the European Society for Vascular Surgery (ESVS). *Eur Heart J* 2018;39:763-816.
3. Tang GL, Chang DS, Sarkar R, Wang R, Messina LM. The effect of gradual or acute arterial occlusion on skeletal muscle blood flow, arteriogenesis, and inflammation in rat hindlimb ischemia. *J Vasc Surg* 2005;41:312-20.
4. Romiti M, Albers M, Brochado-Neto FC, Durazzo AE, Pereira CA, De Luccia N. Meta-analysis of infrapopliteal angioplasty for chronic critical limb ischemia. *J Vasc Surg* 2008;47:975-81.
5. Verheye S, Jolicœur EM, Behan MW, Pettersson T, Sainsbury P, Hill J, et al. Efficacy of a device to narrow the coronary sinus in refractory angina. *N Engl J Med* 2015;372:519-27.
6. Ielasi A, Todaro MC, Grigis G, Tespili M. Coronary Sinus Reducer system™: A new therapeutic option in refractory angina patients unsuitable for revascularization. *Int J Cardiol* 2016;209:122-30.

7. Institute for Laboratory Animal Resources (US), Guide for the Care and Use of Laboratory Animals, Institute for Laboratory Animal Resources (US), Washington, DC, USA, 1996.
8. Rajagopalan S, Mohler ER 3rd, Lederman RJ, Mendelsohn FO, Saucedo JF, Goldman CK, et al. Regional angiogenesis with vascular endothelial growth factor in peripheral arterial disease: a phase II randomized, double-blind, controlled study of adenoviral delivery of vascular endothelial growth factor 121 in patients with disabling intermittent claudication. *Circulation* 2003;108:1933-8.
9. Fadini GP, Agostini C, Avogaro A. Autologous stem cell therapy for peripheral arterial disease meta-analysis and systematic review of the literature. *Atherosclerosis* 2010;209:10-7.
10. Beck CS, Leighninger DS. Scientific basis for the surgical treatment of coronary artery disease. *JAMA* 1955;159:1264-71.
11. Verheye S, Jolicœur EM, Behan MW, Pettersson T, Sainsbury P, Hill J, et al. Efficacy of a device to narrow the coronary sinus in refractory angina. *N Engl J Med* 2015;372:519-27.
12. Kapanadze T, Bankstahl JP, Wittneben A, Koestner W, Ballmaier M, Gamrekelashvili J. Multimodal and Multiscale Analysis Reveals Distinct Vascular, Metabolic and Inflammatory Components of the Tissue Response to Limb Ischemia. *Theranostics* 2019;9:152-66.
13. Diaz JA, Obi AT, Myers Jr DD, Wroblewski SK, Henke PK, Mackman N, et al. Critical Review of Mouse Models of Venous Thrombosis. *Arterioscler Thromb Vasc Biol* 2012;32:556-62.
14. Shi WY, Hu LY, Wu S, Liu CJ, Gu JP. Two swine models of iliac vein occlusion: Which form most contributes to venous thrombosis? *Thromb Res* 2015;135:1172-8.
15. Dokun AO, Keum S, Hazarika S, Li Y, Lamonte GM, Wheeler F, et al. A

- quantitative trait locus (LSq-1) on mouse chromosome 7 is linked to the absence of tissue loss after surgical hindlimb ischemia. *Circulation* 2008;117:1207-15.
16. Sealock R, Zhang H, Lucitti JL, Moore SM, Faber JE. Congenic fine-mapping identifies a major causal locus for variation in the native collateral circulation and ischemic injury in brain and lower extremity. *Circ Res* 2014;114:660-71.
 17. Beckert S et al. The impact of the MicroLightguide O2C for the quantification of tissue ischemia in diabetic foot ulcers. *Diabetes Care* 2004;27:2863-86.
 18. Nakamura H, Makiguchi T, Atomura D, Yamatsu Y, Shirabe K, Yokoo S. Changes in Skin Perfusion Pressure After Hyperbaric Oxygen Therapy Following Revascularization in Patients With Critical Limb Ischemia: A Preliminary Study. *Int J Low Extrem Wounds* 2020;19:57-62.
 19. McClung JM, McCord TJ, Southerland K, Schmidt CA, Padgett ME, Ryan TE, Kontos CD. Subacute limb ischemia induces skeletal muscle injury in genetically susceptible mice independent of vascular density. *J Vasc Surg* 2016;64:1101-11.
 20. Chen M, Fan H, Ledford BT, Farah Z, Barron C, Liu Z, He JQ. Impacts of femoral artery and vein excision versus femoral artery excision on the hindlimb ischemic model in CD-1 mice. *Microvasc Res* 2017;110:48-55.
 21. Long CA, Timmins LH, Koutakis P, Goodchild TT, Lefer DJ, Pipinos II, et al. An endovascular model of ischemic myopathy from peripheral arterial disease. *J Vasc Surg* 2017;66:891-901.
 22. Naehle CP, Steinberg VA, Schild H, Mommertz G. Assessment of peripheral skeletal muscle microperfusion in a porcine model of peripheral arterial stenosis by steady-state contrast-enhanced ultrasound and Doppler flow measurement. *J Vasc Surg* 2015;61:1312-20.

23. Ziegler MA, Distasi MR, Bills RG, Miller SJ, Alloosh M, Murphy MP, et al. Marvels, mysteries, and misconceptions of vascular compensation to peripheral artery occlusion. *Microcirculation* 2010;17:3-20.
24. Gao Y, Aravind S, Patel NS, Fuglestad MA, Ungar JS, Mietus CJ, et al. Collateral Development and Arteriogenesis in Hindlimbs of Swine After Ligation of Arterial Inflow. *J Surg Res* 2020;249:168-79.
25. Rushing AM, Donnarumma E, Polhemus DJ, Au KR, Victoria SE, Schumacher JD, et al. Effects of a novel hydrogen sulfide prodrug in a porcine model of acute limb ischemia. *J Vasc Surg* 2019;69:1924-35.

ABSTRACT(IN KOREAN)

돼지 급성 하지허혈 모형에서 인위적 장골정맥 폐쇄를 이용한
정맥 환류 제한이 조직 산소포화도에 미치는 효과

<지도교수 홍 명 기>

연세대학교 대학원 의학과

김 원 호

하지 허혈 조직에 충분한 산소의 공급은 상처 회복 (wound healing)과 사지 구제 (limb salvage)에 중요하다. 본 연구는 돼지 급성 하지허혈 모형 (acute limb ischemia)에서 인위적 총장골정맥 (common iliac vein)의 부분 폐쇄 (partial occlusion)를 통한 정맥 환류 (venous return) 제한이 하지 허혈 조직 산소포화도 (tissue oxygenation)에 미치는 영향에 대하여 조사하였다. 실험은 총 7마리 돼지를 사용하였다. 급성 하지허혈 모형을 만들기 위하여 혈관 폐색장치 (vascular plug)를 이용하여 색전술을 통한 우측 총장골동맥 (common iliac artery)의 통한 완전 폐쇄를 유도하였다. 조직 산소 포화도 측정은 우측 표피대퇴동맥 (superficial femoral artery)과 동측 총장골정맥에서 채취한 혈액으로 혈액가스검사 (blood gas analysis)를 시행하였다. 순서는 처음 (T0)과, 동맥 색전술 직후 (T1), 동맥 색전술 10분 후 (T2), 20분 후 (T3), 40분 후 (T4), 이어서 인위적으로 혈관내 풍선 확장술 (intravascular balloon inflation)로 총장골정맥을 부분 폐쇄를 시켜 정맥 환류 제한을 유도한 다음 5분 후 (T5), 15분 후 (T6), 마지막으로 25분 후 (T7)으로, 총 8번을 평가하였다. 결과 상, 표피대퇴동맥에서 측정된 동맥혈의 조직 산소포화도는 모든 단계에서 큰 변화가 없었다. 반면에, 총장골동맥 색전술 후 총장골정맥의 조직 산소포화도는 의미 있게 감소하였다 (T0 단계 85.57 ± 1.72 %에서 T4 단계 71.86 ± 7.58 % 로 감소). 흥미롭게도 총장골정맥을 부분 폐쇄한 후에는, 정맥혈 조직 산소포화도가 다시 증가하였다 (T5 단계 $74.14 \pm$

9.56 %에서 T7 단계 $79.71 \pm 4.78\%$ 로 증가). 한편, 좌측 하지는 총장골정맥을 폐쇄하지 않은 급성 하지허혈 대조군 (control limb)으로 설정하여, 우측 실험군 (experimental limb)과 조직 산소포화도의 차이를 분석하였는데, 정맥혈 조직 산소포화도는 시간이 지남에 따라 실험군에서 대조군보다 의미있게 높았다 (T7 단계에서 실험군은 $79.71 \pm 4.78\%$, 대조군은 $60.00 \pm 4.24\%$). 위 결과들을 종합할 때, 급성 하지허혈 모형에서 총장골정맥의 부분 폐쇄는 정맥혈의 일시적인 울혈 및 압력 상승을 유도하여, 조직산소포화도를 증가시키고, 허혈 조직이 산소를 이용할 수 있는 시간을 늘려, 상처 회복에 도움이 될 수 있는 가능성을 보여주었다. 향후, 위와 같은 기전의 유용성을 검증하는 후속 연구가 필요하다.

핵심되는 말: 급성 하지허혈, 산소포화도, 총장골정맥 부분 폐쇄