

Original Article

Current Prevalence of the *crpP* Gene in Carbapenemase-Producing *Pseudomonas aeruginosa* Blood Isolates in Korea

Jinho Heo¹, Yu Jeong Choi¹, Young Ah Kim², Seok Hoon Jeong^{1,3}, Jong Hee Shin⁴, Kyeong Seob Shin⁵, Jeong Hwan Shin⁶, Young Ree Kim⁷, Hyun Soo Kim⁸, Young Uh⁹, Nam Hee Ryo¹⁰

¹Department of Laboratory Medicine, Yonsei University College of Medicine, Seoul, ²Department of Laboratory Medicine, National Health Insurance Service Ilsan Hospital, Goyang, ³Research Institute of Bacterial Resistance, Yonsei University College of Medicine, Seoul, ⁴Department of Laboratory Medicine, Chonnam National University Medical School, Gwangju, ⁵Department of Laboratory Medicine, Chungbuk National University College of Medicine, Cheongju, ⁶Department of Laboratory Medicine and Paik Institute for Clinical Research, Inje University College of Medicine, Busan, ⁷Department of Laboratory Medicine, Jeju National University, College of Medicine, Jeju, ⁸Department of Laboratory Medicine, Hallym University Dongtan Sacred Heart Hospital, Hallym University College of Medicine, Hwaseong, ⁹Department of Laboratory Medicine, Yonsei University Wonju College of Medicine, Wonju, ¹⁰Department of Laboratory Medicine, Keimyung University School of Medicine, Daegu, Korea

국내 혈액분리 카바페넴 분해효소 생성 녹농균에서의 *crpP* 유전자 확산 현황

허진호¹, 최유정¹, 김영아², 정석훈^{1,3}, 신종희⁴, 신경섭⁵, 신정환⁶, 김영리⁷, 김현수⁸, 어영⁹, 류남희¹⁰

¹연세대학교 의과대학 진단검사의학교실, ²국민건강보험 일산병원 진단검사의학과, ³연세대학교 의과대학 세균내성연구소, ⁴전남대학교병원 진단검사의학과, ⁵충북대학교병원 진단검사의학과, ⁶인제대학교 부산백병원 진단검사의학과, ⁷제주대학교병원 진단검사의학과, ⁸한림대학교병원 진단검사의학과, ⁹연세대학교 원주의과대학 진단검사의학과, ¹⁰계명대학교병원 진단검사의학과



OPEN ACCESS

pISSN : 2288-0585
eISSN : 2288-6850

Ann Clin Microbiol 2022 June, 25(2): 59-65
<https://doi.org/10.5145/ACM.2022.25.2.4>

Corresponding author

Young Ah Kim

E-mail: yakim@nhimc.or.kr

Tel: +82-31-900-0908

Fax: +82-31-900-9012

Received: January 25, 2022

Revised: May 01, 2022

Accepted: May 17, 2022

© 2022 Korean Society of Clinical Microbiology.



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Recently, CrpP enzymes have been described as a novel cause of ciprofloxacin resistance. The *crpP* gene encodes a novel protein that specifically confers resistance to ciprofloxacin through an adenosine triphosphate-dependent mechanism that phosphorylates the antimicrobial. In this study, the current prevalence of the *crpP* gene in carbapenemase-producing *Pseudomonas aeruginosa* blood isolates was evaluated.

Methods: During the study of the Antimicrobial Resistance Surveillance System in Korea, 22 blood isolates of carbapenemase-producing *P. aeruginosa* were collected from nine general hospitals and two nursing homes in the year 2020. Resistance genes and phylogenetic trees were analyzed with the whole genome sequencing data.

Results: A total of 11 *P. aeruginosa* blood isolates harbored the *crpP* and carbapenemase genes (nine IMP-6 producers and two GES-5-producers). Nine NDM-1-producers harbored *aac(6')-Ib-cr* and *qnrVC1*. One GES-9-producer also carried *aac(6')-Ib-cr*, and one NDM-1-producer also carried *qnrVC1*. The phylogenetic tree showed no epidemiologic link among the 22 carbapenemase-producing *P. aeruginosa* isolates.

Conclusion: This is the first report on the current prevalence of the *crpP* gene in carbapenemase-producing *P. aeruginosa* blood isolates in Korea.

Keywords: Fluoroquinolone, Resistance, *crpP* gene, *Pseudomonas aeruginosa*

INTRODUCTION

Pseudomonas aeruginosa, one of the antimicrobial-resistant ESKAPE (*Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *P. aeruginosa*, and *Enterobacter* species) pathogens, represents a global threat to human health [1]. *P. aeruginosa* usually shows multidrug resistance. Its co-resistance to carbapenems, aminoglycosides, polymyxins, and tigecycline has also increased [2]. There are limited treatment options for serious infections caused by multidrug-resistant *P. aeruginosa*. Many combination therapies have been tried [3]. Recently, it has been reported that combination therapy including ciprofloxacin is correlated with a lower mortality. A combination of a beta-lactam and ciprofloxacin has been proposed for ciprofloxacin-susceptible *P. aeruginosa* bacteremia [4].

According to data from the Korean global antimicrobial resistance surveillance system (Kor-GLASS) [5], of *P. aeruginosa* blood isolates in 2017-2019, 12.3% showed resistance to piperacillin, 13.1% to ceftazidime, 12.9% to cefepime, 21.3% to imipenem, 22.1% to meropenem, 9.6% to amikacin, and 18.9% to ciprofloxacin. Most the carbapenemase-producers exhibited co-resistance to amikacin. However, data about its co-resistance to ciprofloxacin are unavailable. This study provides the newest data about the resistance mechanism to fluoroquinolone of carbapenemase-producing *P. aeruginosa*.

Resistance mechanisms of *P. aeruginosa* to fluoroquinolone are known mostly through the acquisition of mutations in genes encoding target proteins of fluoroquinolone and regulators of efflux pumps, leading to overexpression of these pumps [6]. Quinolone resistance may also be attributable to mutations of target enzymes of topoisomerases II and IV encoded by *gyrA* and *parC*, respectively [7]. These mechanisms are known to be chromosomally mediated. However, plasmid-mediated quinolone resistance (PMQR) in *P. aeruginosa* is also important. It is associated with *qnrA-E*, *qnrS*, *qnrVC*, *qepA*, *oqxAB* (efflux pump), and *acc(6)-Ib-cr* (quinolone-modifying enzyme) [7]. Recently, CrpP enzymes have been described as a novel ciprofloxacin-resistance mechanism. The *crpP* gene encodes a novel protein, capable of specifically conferring resistance to ciprofloxacin in *Escherichia coli* through an adenosine triphosphate-dependent mechanism that involves phosphorylation of the antibiotic [8].

The purpose of this study was to report on the current situation of *crpP* gene spread in carbapenemase-producing *P. aeruginosa* blood isolates in Korea.

MATERIALS AND METHODS

In this study, non-duplicated *P. aeruginosa* blood isolates were isolated from nine nationwide general hospitals and two nursing homes according to the Kor-GLASS manual [5] in 2020. Briefly, pure colonies of *P. aeruginosa* were collected in 10% skim milk and stored at -70°C before all collected isolates were transferred to a single analysis center of the Korea Disease Control and Prevention Agency for analyses using approved methods [5]. Bacterial species were verified using Matrix-Assisted Laser Desorption/

Ionization Time-of-Flight mass spectrometry (Bruker Biotyper, Bruker Daltonics GmbH, Bremen, Germany). Antimicrobial susceptibility was mainly determined by the disk diffusion test according to the Clinical and Laboratory Standards Institute (CLSI) guidelines [9].

To detect carbapenemase-producers, *P. aeruginosa* isolates showing nonsusceptibility to imipenem or meropenem were PCR-sequenced to detect *bla*_{KPC}, *bla*_{NDM}, *bla*_{OXA-48}, *bla*_{VIM}, *bla*_{IMP}, and *bla*_{GES}. For whole-genome sequencing, DNAs of freshly subcultured isolates were extracted using a GenElute™ Bacterial Genomic DNA Kit (Sigma-Aldrich, St. Louis, MO, USA). Then 8 µg of input genomic DNA was used to sequence the entire genome using NextSeq 550 instrument (Illumina, San Diego, CA, USA). Sequences were assembled with Spades (version 3.11.1) and annotated with Prokka (version 1.13.7). Resistance genes were obtained with ResFinder 4.1 from the website of the center for genomic epidemiology [10]. A phylogenetic tree was generated based on whole-genome multilocus sequence typing using a BioNumerics software, version 7.6.3 (Applied Maths, St Martens Latem, Belgium) [11] (Fig. 1).




Fig. 1. Phylogenetic tree of 22 carbapenemase-producing *Pseudomonas aeruginosa* strains based on whole genome multilocus sequence typing. Dendrogram was generated with BioNumerics software.

RESULTS

Among 212 non-duplicated *P. aeruginosa* blood isolates, 22 carbapenemase-producing *P. aeruginosa* isolates showed resistance to both ciprofloxacin and imipenem. A total of 11 carbapenemase-producing *P. aeruginosa* blood isolates (nine IMP-6 producers and two GES-5-producers) co-harbored *aac(6)-Ib-cr* and *crpP* (Table 1). A total of nine NDM-1-producers co-harbored *aac(6)-Ib-cr* and *qnrVC1*. One GES-9-producer also carried *aac(6)-Ib-cr* and one NDM-1-producer also carried *qnrVC1*. The phylogenetic tree showed no epidemiologic link among 22 carbapenemase-producing *P. aeruginosa* isolates.

Table 1. Resistance to carbapenem and ciprofloxacin with corresponding resistance genes

Isolates	Phenotype (ZD)			Genotype			
	IPM	MEM	CIP	CRP	PMQR	<i>gyrA</i>	<i>parC</i>
F0020PA0023	R (11)	R (10)	R (6)	<i>bla</i> _{GES-9}	<i>aac(6')-Ib-cr</i>	Thr83Ile	Ser87Leu
A0020PA0022	R (7)	R (6)	R (6)	<i>bla</i> _{GES-5}	<i>aac(6')-Ib-cr, crpP</i>	Thr83Ile	Ser87Leu
A0020PA0005	R (7)	R (6)	R (6)	<i>bla</i> _{GES-5}	<i>aac(6')-Ib-cr, crpP</i>	Thr83Ile	WT
A0020PA0023	R (6)	R (6)	R (6)	<i>bla</i> _{IMP-6}	<i>aac(6')-Ib-cr, crpP</i>	Thr83Ile	Ser87Leu
A0020PA0007	R (6)	R (6)	R (6)	<i>bla</i> _{IMP-6}	<i>aac(6')-Ib-cr, crpP</i>	Thr83Ile	Ser87Leu
C0020PA0004	R (6)	R (6)	R (6)	<i>bla</i> _{IMP-6}	<i>aac(6')-Ib-cr, crpP</i>	Thr83Ile	Ser87Leu
E0020PA0006	R (6)	R (6)	R (6)	<i>bla</i> _{IMP-6}	<i>aac(6')-Ib-cr, crpP</i>	Thr83Ile	Ser87Leu
E0020PA0012	R (6)	R (6)	R (6)	<i>bla</i> _{IMP-6}	<i>aac(6')-Ib-cr, crpP</i>	Thr83Ile	Ser87Leu
E0020PA0025	R (6)	R (6)	R (6)	<i>bla</i> _{IMP-6}	<i>aac(6')-Ib-cr, crpP</i>	Thr83Ile	Ser87Leu
E0020PA0035	R (6)	R (6)	R (6)	<i>bla</i> _{IMP-6}	<i>aac(6')-Ib-cr, crpP</i>	Thr83Ile	Ser87Leu
E0020PA0039	R (6)	R (6)	R (6)	<i>bla</i> _{IMP-6}	<i>aac(6')-Ib-cr, crpP</i>	Thr83Ile	Ser87Leu
I0020PA0016	R (6)	R (6)	R (6)	<i>bla</i> _{IMP-6}	<i>aac(6')-Ib-cr, crpP</i>	Thr83Ile	Ser87Leu
B0020PA0004	R (6)	R (6)	R (6)	<i>bla</i> _{NDM-1}	<i>aac(6')-Ib-cr, qnrVC1</i>	Thr83Ile	Ser87Leu
B0020PA0010	R (6)	R (6)	R (6)	<i>bla</i> _{NDM-1}	<i>aac(6')-Ib-cr, qnrVC1</i>	Thr83Ile	Ser87Leu
F0020PA0002	R (6)	R (6)	R (6)	<i>bla</i> _{NDM-1}	<i>aac(6')-Ib-cr, qnrVC1</i>	Thr83Ile	Ser87Leu
F0020PA0003	R (6)	R (6)	R (6)	<i>bla</i> _{NDM-1}	<i>aac(6')-Ib-cr, qnrVC1</i>	Thr83Ile	Ser87Leu
I0020PA0007	R (6)	R (6)	R (6)	<i>bla</i> _{NDM-1}	<i>aac(6')-Ib-cr, qnrVC1</i>	Thr83Ile	Ser87Leu
I0020PA0009	R (6)	R (6)	R (6)	<i>bla</i> _{NDM-1}	<i>aac(6')-Ib-cr, qnrVC1</i>	Thr83Ile	Ser87Leu
I0020PA0021	R (6)	R (6)	R (6)	<i>bla</i> _{NDM-1}	<i>aac(6')-Ib-cr, qnrVC1</i>	Thr83Ile	Ser87Leu
I0020PA0024	R (6)	R (6)	R (6)	<i>bla</i> _{NDM-1}	<i>aac(6')-Ib-cr, qnrVC1</i>	Thr83Ile	Ser87Leu
I0020PA0026	R (6)	R (6)	R (6)	<i>bla</i> _{NDM-1}	<i>aac(6')-Ib-cr, qnrVC1</i>	Thr83Ile	Ser87Leu
I0020PA0004	R (6)	R (6)	R (6)	<i>bla</i> _{NDM-1}	<i>qnrVC1</i>	Thr83Ile	Ser87Leu

Abbreviations: IMP, imipenem; MEM, meropenem; CIP, ciprofloxacin; CRP, carbapenemase; PMQR, plasmid-mediated quinolone resistance; R, resistant; ZD, zone diameter (mm); WT, wild type.

DISCUSSION

Quinolones have been commonly used as a treatment option for a large number of bacterial infections. Ciprofloxacin is the most active antibiotic in this group. The increase in quinolone resistance has limited the effect of quinolones. There was no report about the prevalence of quinolone resistance genes in *P. aeruginosa* in Korea to the best of our knowledge. Recently, novel mutations in *gyrA* and *parC* genes were first found in *P. aeruginosa* isolated from companion dogs in South Korea [12]. In that study, a total of 84 nonduplicated *P. aeruginosa* strains were obtained from healthy dogs and infected dogs. The resistance rate was 14.3% for levofloxacin and 13.1% for ciprofloxacin. The resistance was commonly associated with *gyrA* mutations [12].

The information about the prevalence of quinolone resistance genes of clinically important pathogens in Korea is limited. For *K. pneumoniae*, mutations in *gyrA* and *parC* were found in 78.9% and 65.5% of 142 extended-spectrum β -lactamases-producers, respectively [13]. The common PMQR gene was *qnrB-aac(6')-Ib-cr-oqxAB* (58/142, 40.8%) [13]. Lee et al. [14] have reported the presence of PMQR genes in *Salmonella enterica* isolated from human salmonellosis patients in South Korea from 2016 to 2019 [14]. Among 34 *Salmonella* strains with reduced susceptibility to quinolones, 25 strains harbored one or two of *qnrA*, *qnrB*, *qnrS* (most common), and *aac(6')-Ib-cr* genes.

Out of 22 carbapenemase-producers from nationwide collections, we detected 11 *P. aeruginosa* blood isolates that co-harbored *crpP* and carbapenemase genes. This is the first report about the current situation of *crpP* gene spread in carbapenemase-producing *P. aeruginosa* blood isolates in Korea. Because all strains also have *aac(6)-Ib-cr* and chromosomal mutations in *gyrA* or *parC*, the effect of the presence of *crpP* gene on quinolone resistance is hardly speculated in this study.

The *crpP* gene obtained from the pUM505 plasmid isolated from a *P. aeruginosa* clinical isolate was identified in 1986 [8]. Regarding the expression of CrpP in the transconjugants test, CrpP proteins increased minimal inhibitory concentration values of ciprofloxacin concerning *E. coli* J53-3. However, these changes were not enough to be considered as resistance according to the breakpoint value (≥ 4 mg/L) reported for this interpretation by the CLSI [8]. Nevertheless, the *crpP* gene conferring low-level resistance could facilitate the selection of mutants with a higher level of quinolone resistance. In addition, plasmids contain other genes in addition to the *crpP* gene, which encodes additional quinolone resistance mechanisms [8].

In this study, the type of carbapenemase of *crpP* gene-carrying *P. aeruginosa* was mostly IMP-6, known to be a common type in Korea [15]. NDM-1-producing *P. aeruginosa* isolates did not carry *crpP* genes, which seemed to become one of the majority types of carbapenemase in *P. aeruginosa*. The shift in the molecular epidemiology of carbapenemase genes might change the molecular epidemiology of PMQR genes.

Mobile genetic elements play a key role in the spread of resistance genes, and high-risk clones frequently integrate such determinants into their genomes [16]. In this study, the linkage of *crpP* and *aac(6)-Ib-cr* gene with *bla*_{IMP-6} is suspected. And there is high possibility that *aac(6)-Ib-cr* and *qnrVC1* are associated with *bla*_{NDM-1} in *P. aeruginosa*.

If only PMQR is present, it is expected to be susceptible or low-grade resistance phenotypes [7,8]. Unfortunately, the only effect of *crpP* gene is hard to know in this study, because all isolates had chromosomal mutations in *gyrA* or *parC*, in addition to PMQRs. These strains are usually associated with a large reduction in biological fitness with the accumulation of more resistance-associated mutations [17]. One more thing to consider is that NDM-1 carrying *P. aeruginosa* is mostly separated from hospitals in Daegu or Busan and the regional differences could have an effect. Therefore, we suggest continuous monitoring of the change of PMQR in *P. aeruginosa*.

In conclusion, the spread of *P. aeruginosa* isolates, coharboring *crpP* and carbapenemase genes were found in this study. This is the first report about the current situation of *crpP* gene spread in carbapenemase-producing *P. aeruginosa* blood isolates in Korea.

요약

배경: 새로운 퀴놀론 내성기전으로 알려진 *crpP* 내성유전자는 CrpP 효소를 생성하고 이 단백질은 adenosine triphosphate에 의존하여 항균제의 phosphorylation을 유발하여 내성을 유발하는 것으로 알려져 있다. 본 연구에서는 혈액에서 분리된 카바페넴 분해효소를 생성하는 *Pseudomonas aeruginosa*에서 *crpP* 내성유전자 확산의 최근 현황을 살펴보고자 한다.

방법: 국내 내성현황 조사를 통해 9개의 종합병원과 2개의 요양병원에서 총 22주의 카바페넴 분해효소를 생성하는 *P. aeruginosa*를 분리하였고, 전장유전체 분석을 통해 내성유전자와 계통도를 분석하였다.

결과: 총 11주의 *P. aeruginosa* 혈액분리주가 CrpP와 카바페넴 분해효소를 코딩하는 내성 유전자를 동시에 가지고 있었다(IMP-6 생성 9주와 GES-5 생성 2주). 총 9주의 NDM-1-생성주는 *aac(6)-Ib-cr*와 *qnrVCI*를 가지고 있었다. 또 GES-9 생성주 한 주는 *aac(6)-Ib-cr*를 가지고 있었고, 나머지 NDM-1 생성주 한주는 *qnrVCI*를 가지고 있었다. 계통도 분석에서 역학적 연관성은 확인되지 않았다.

결론: 본 연구는 국내 카바페넴 분해효소를 생성하는 혈액분리 *P. aeruginosa*에서의 *crpP* 내성유전자의 확산에 관한 최근 현황을 최초로 보고하고 있다.

CONFLICTS OF INTEREST

No potential conflicts of interest relevant to this article were reported.

FUNDING

This work was supported by a fund (2020E540600) by the Research of Korea Disease Control and Prevention Agency.

REFERENCES

1. De Oliveira DMP, Forde BM, Kidd TJ, Harris PNA, Schembri MA, Beatson SA, et al. Antimicrobial resistance in ESKAPE pathogens. *Clin Microbiol Rev* 2020;33:e00181-19.
2. Karakonstantis S, Kritsotakis EI, Gikas A. Treatment options for *K. pneumoniae*, *P. aeruginosa* and *A. baumannii* co-resistant to carbapenems, aminoglycosides, polymyxins and tigecycline: an approach based on the mechanisms of resistance to carbapenems. *Infection* 2020;48:835-51.
3. Doi Y. Treatment options for carbapenem-resistant gram-negative bacterial infections. *Clin Infect Dis* 2019;69:S565-75.
4. Paulsson M, Granrot A, Ahl J, Tham J, Resman F, Riesbeck K, et al. Antimicrobial combination treatment including ciprofloxacin decreased the mortality rate of *Pseudomonas aeruginosa* bacteraemia: a retrospective cohort study. *Eur J Clin Microbiol Infect Dis* 2017;36:1187-96.
5. Kim D, Yoon EJ, Hong JS, Choi MH, Kim HS, Kim YR, et al. Major bloodstream infection-causing bacterial pathogens and their antimicrobial resistance in South Korea, 2017-2019: phase I report from Kor-GLASS. *Front Microbiol* 2022;12:799084.
6. Rehman A, Patrick WM, Lamont IL. Mechanisms of ciprofloxacin resistance in *Pseudomonas aeruginosa*: new approaches to an old problem. *J Med Microbiol* 2019;68:1-10.
7. Ming DS, Chen QQ, Chen XT. Detection of 5 kinds of genes related to plasmid-mediated quinolone resistance in four species of nonfermenting bacteria with 2 drug resistant phenotypes. *Can J Infect Dis Med Microbiol* 2020;2020:3948719.
8. Chávez-Jacobo VM, Hernández-Ramírez KC, Romo-Rodríguez P, Pérez-Gallardo RV, Campos-García J, Gutiérrez-Corona JF, et al. CrpP is a novel ciprofloxacin-modifying enzyme encoded by the *Pseudomonas aeruginosa* pUM505 plasmid. *Antimicrob Agents Chemother* 2018;62:e02629-17.

9. Clinical and Laboratory Standards Institute (CLSI). Performance standards for antimicrobial susceptibility testing; CLSI Supplement M100. Wayne, PA: 2020.
10. Center for Genomic Epidemiology. Center for Genomic Epidemiology Web site. www.genomicepidemiology.org [Online] (last visited on 13 June 2022).
11. Blanc DS, Magalhães B, Koenig I, Senn L, Grandbastien B. Comparison of whole Genome (wg-) and core genome (cg-) MLST (BioNumerics™) versus SNP variant calling for epidemiological investigation of *Pseudomonas aeruginosa*. *Front Microbiol* 2020;11:1729.
12. Park Y, Oh J, Park S, Sum S, Song W, Chae J, et al. Antimicrobial resistance and novel mutations detected in the *gyrA* and *parC* genes of *Pseudomonas aeruginosa* strains isolated from companion dogs. *BMC Vet Res* 2020;16:111.
13. Kim JO, Yoo IY, Yu JK, Kwon JA, Kim SY, Park YJ. Predominance and clonal spread of CTX-M-15 in cefotaxime-resistant *Klebsiella pneumoniae* in Korea and their association with plasmid-mediated quinolone resistance determinants. *J Infect Chemother* 2021;27:1186-92.
14. Lee S, Park N, Yun S, Hur E, Song J, Lee H, et al. Presence of plasmid-mediated quinolone resistance (PMQR) genes in non-typhoidal *Salmonella* strains with reduced susceptibility to fluoroquinolones isolated from human salmonellosis in Gyeonggi-do, South Korea from 2016 to 2019. *Gut Pathog* 2021;13:35.
15. Yoon EJ, Jeong SH. Mobile carbapenemase genes in *Pseudomonas aeruginosa*. *Front Microbiol* 2021;12:614058.
16. Kocsis B, Toth A, Gulyas D, Ligeti B, Katona K, Rokusz L, et al. Acquired *qnrVC1* and *bla_{NDM-1}* resistance markers in an international high-risk *Pseudomonas aeruginosa* ST773 clone. *J Med Microbiol* 2019;68:336-8.
17. Lindgren PK, Marcusson LL, Sandvang D, Frimodt-Møller N, Hughes D. Biological cost of single and multiple norfloxacin resistance mutations in *Escherichia coli* implicated in urinary tract infections. *Antimicrob Agents Chemother* 2005;49:2343-51.