



The Association Between Objectively Measured Vision Impairment and Self-Reported Physical Activity Among 34,129 Adults Aged ≥ 50 Years in Six Low- and Middle-Income Countries

Lee Smith, Shahina Pardhan, Trish J. Gorely, Yvonne Annette Barnett, Louis Jacob, Guillermo Felipe López-Sánchez, Mark A. Tully, Nicola Veronese, Jae-Il Shin, Ai Koyanagi

► To cite this version:

Lee Smith, Shahina Pardhan, Trish J. Gorely, Yvonne Annette Barnett, Louis Jacob, et al.. The Association Between Objectively Measured Vision Impairment and Self-Reported Physical Activity Among 34,129 Adults Aged ≥ 50 Years in Six Low- and Middle-Income Countries. *Journal of Aging and Physical Activity*, 2022, 30 (2), pp.316-322. 10.1123/japa.2021-0103 . hal-03673640

HAL Id: hal-03673640

<https://hal.science/hal-03673640>

Submitted on 17 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The association between objectively measured vision impairment and self-reported physical activity among 34,129 adults aged ≥ 50 years in six low- and middle-income countries

Lee Smith^{1*}, Shahina Pardhan², Trish Gorely³, Yvonne Barnett⁴, Louis Jacob^{5,6}, Guillermo F.

López-Sánchez^{2*}, Mark A. Tully⁷, Nicola Veronese⁸, Jae Il Shin⁹, Ai Koyanagi^{6,10}

1. The Cambridge Centre for Sport and Exercise Sciences, Anglia Ruskin University,
Cambridge CB1 1PT, UK

2. Vision and Eye Research Institute, School of Medicine, Faculty of Health, Education,
Medicine and Social Care, Anglia Ruskin University-Cambridge Campus, Cambridge,
United Kingdom.

3. Department of Nursing and Midwifery, University of the Highlands and Islands, Inverness,
UK

4. School of Life Sciences, Anglia Ruskin University, Cambridge, CB1 1PT, UK

5. Faculty of Medicine, University of Versailles Saint-Quentin-en-Yvelines, Montigny-le-
Bretonneux, 78180 Versailles, France

6. Research and Development Unit, Parc Sanitari Sant Joan de Déu, CIBERSAM, Dr. Antoni
Pujadas, 42, Sant Boi de Llobregat, 08830 Barcelona, Spain

7. Institute of Mental Health Sciences, School of Health Sciences, Ulster University,
Newtownabbey, UK

8. Department of Geriatric Medicine, University of Palermo, Italy

9. Department of Pediatrics, Yonsei University College of Medicine, Seoul, Republic of
Korea

10. ICREA, Pg. Lluís Companys 23, 08010, Barcelona, Spain

*** Corresponding authors:** Lee Smith. Lee.Smith@aru.ac.uk. Guillermo F. López-Sánchez.
guillermo.lopez-sanchez@aru.ac.uk

ABSTRACT

We investigated the association between vision impairment and physical activity among older adults from low- and middle-income countries (LMICs). Visual acuity was measured using the tumbling ElogMAR chart, and vision impairment was defined as visual acuity worse than 6/18 (0.48 logMAR) in the better seeing eye. Physical activity was assessed by the Global Physical Activity Questionnaire. Multivariable logistic regression and meta-analysis were conducted to assess associations. The sample included 34,129 individuals aged 50-114 years [mean (SD) age 62.4 (16.0) years; 47.9% males]. After adjustment for confounders, near vision impairment was not significantly associated with low physical activity, but far vision impairment showed a significant association (OR=1.32; 95%CI=[1.17-1.49], $I^2=0.0\%$). Far vision impairment was dose-dependently associated with low physical activity [e.g., severe (<6/10) vs. no ($\geq 6/12$) far vision impairment (OR=1.80; 95%CI=[1.03-3.15]). Interventions to address low levels of physical activity in the visually impaired in LMICs should target those with far vision impairment.

Key Words: Visual Impairment, Physical Activity, Low- and Middle-Income Countries, Epidemiology

INTRODUCTION

Physical activity (bodily movement caused by contraction of skeletal muscle that results in energy expenditure) can be categorized into multiple domains including, structured exercise and sport, active travel (walking and cycling), occupational activity and household chores/gardening (Caspersen, Powell, & Christenson, 1985). Participation in physical activity is beneficial for both the physical and mental health of adults. For example, a recent systematic review of review articles found that physically active older adults are at a reduced risk of cardiovascular mortality, breast and prostate cancer, fractures, recurrent falls, functional limitation, and depression. Moreover, the review found that physically active older adults experience healthier ageing trajectories, better quality of life and improved cognitive functioning (Cunningham, O'Sullivan, Caserotti, & Tully, 2020). Moreover, low levels of physical activity have an important economic burden. For example, in the UK alone, physical inactivity is expected to cost the National Health Service approximately £1.3 billion by 2030 (Sport England, 2019). In light of this evidence, the World Health Organization (WHO) produced guidance in relation to physical activity levels. The key message from this guidance is that adults including older adults should achieve at least 150 minutes of moderate physical activity and/or 75 minutes of vigorous physical activity per week (WHO, 2010). It is therefore important to ensure that all populations maintain adequate levels of physical activity for good health. However, literature shows that as adults age, levels of physical activity decline. For example, one study in a sample of 5022 participants (mean age 61 years; 2114 male) from the UK observed that there was an overall trend for increasing levels of inactivity and a reduction in vigorous activity over a period of 10 years (Smith, Gardner, Fisher, & Hamer, 2015).

Despite the known benefits, some groups of people engage in low levels of physical activity, jeopardizing health status. One such group are those with visual impairment. Low levels of physical activity in this group may be due to factors such as fear of falling, lack of access to adapted recreational and athletic programmes for those with vision impairment, and help or encouragement in developing suitable and safe physical recreation skills and habits specifically tailored for those with visual impairment (Capella-McDonnall, 2007). A previous study found in a sample of 6,634 UK older adults (mean (SD) age 65 (9.2) years) that those with poor vision were twice as likely to be physically inactive than those with good eyesight (Smith et al., 2017). Similar findings have been found in adults residing in the US and Spain (López-Sánchez, Grabovac, Pizzol, Yang, & Smith, 2019; Smith et al., 2019; Willis, Jefferys, Vitale, & Ramulu, 2012) .

However, the current literature has several limitations. First, only a few studies have used objective measures to record vision status with the majority of studies using self-report (Smith et al, 2019). Self-reported measures of vision are often crude in nature, for example “is your eyesight (using glasses or corrective lenses; if you use them) excellent/very good/good/fair/ or poor” (Smith et al., 2017). Thus, these measures are not able to determine acuity and consequently unable to diagnose or confirm visual impairment. Next, while there are a few studies on self-reported measures of visual acuity and physical activity from LMICs (Smith et al., 2021), there are currently no studies on objectively measured visual acuity and physical activity from low- and middle-income countries (LMICs). This is an important omission as visual difficulties have been reported to be more common in LMICs than in high-income countries (HICs) (Freeman et al., 2013), while it is possible that people with vision impairment may have particular difficulties in engaging in physical activity in LMICs due to factors such as lack of visually impaired accessible facilities. Furthermore, there are only a

few studies that have specifically focused on the older population despite the fact that the prevalence of visual impairment and low physical activity increase with age (Klaver, Wolfs, Vingerling, Hofman, & de Jong, 1998; Smith et al., 2015).

Therefore, the aim of the present study was to investigate the association between objectively measured visual impairment and self-reported physical activity among adults aged ≥ 50 years from six LMICs (China, Ghana, India, Mexico, Russia, South Africa), which broadly represent different geographical locations and levels of socio-economic and demographic transition. We hypothesized that those with visual impairment will report lower levels of physical activity.

METHODS

Publically available data from the SAGE (<http://www.who.int/healthinfo/sage/en/>) were analyzed. This survey was undertaken in China, Ghana, India, Mexico, Russia, and South Africa between 2007 and 2010. All countries were LMICs based on the World Bank classification at the time of the survey.

Details of the SAGE survey methodology have been published previously (Kowal et al., 2012). In brief, in order to obtain nationally representative samples, a multistage clustered sampling design method was used. The sample consisted of adults aged ≥ 18 years with oversampling of those aged ≥ 50 years. Trained interviewers conducted face-to-face interviews using a standard questionnaire. Standard translation procedures were undertaken to ensure comparability between countries. The survey response rates were: China 93%; Ghana 81%; India 68%; Mexico 53%; Russia 83%; and South Africa 75%. Sampling weights

were constructed to adjust for the population structure as reported by the United Nations Statistical Division. Ethical approval was obtained from the WHO Ethical Review Committee and local ethics research review boards. Written informed consent was obtained from all participants.

Physical activity

Levels of physical activity was assessed with the validated Global Physical Activity Questionnaire (Bull, Maslin, & Armstrong, 2009). The total amount of moderate-to-vigorous physical activity in a typical week was calculated based on self-report. Those scoring ≥ 150 minutes of moderate-to-vigorous intensity physical activity were classified as meeting the recommended guidelines (coded=0), and those scoring < 150 minutes (low physical activity) were classified as not meeting the recommended WHO guidelines (coded=1) (WHO, 2010).

Visual impairment

Visual acuity was measured using the tumbling ElogMAR chart for distance and near acuity separately for each eye. A string was used to measure 40 cm as the test distance for near visual acuity. The interviewer was instructed to check that the vision charts are well lit and to make sure that the surface does not reflect glare. Furthermore, the respondent was instructed to use glasses or contact lenses if they usually wear them. We defined vision impairment (at distance and near) according to the World Health Organization definition for moderate vision impairment, which refers to visual acuity worse than 6/18 (0.48 logMAR) in the better seeing eye (Ehrlich, Stagg, Andrews, Kumagai, & Musch, 2019). We also categorized far vision into the following levels of severity: no vision impairment (6/12 or better); mild vision

impairment = 6/18 or better but worse than 6/12; moderate vision impairment = 6/60 or better but worse than 6/18; severe vision impairment = worse than 6/60 (World Health Organization., 2019).

Control variables

The control variables, selected based on past literature (Smith et al., 2019), were age, sex, wealth quintiles based on country-specific income, highest level of education achieved (primary, secondary, tertiary), smoking (never, current, former), obesity, and chronic physical conditions (angina, arthritis, diabetes, stroke). A stadiometer and a routinely calibrated electronic weighting scale were used to measure height and weight respectively. Obesity was defined as body mass index $\geq 30 \text{ kg/m}^2$. Arthritis, diabetes, and stroke were based on self-reported lifetime diagnosis. For angina, in addition to a self-reported diagnosis, a symptom-based diagnosis based on the Rose questionnaire was also used (Rose, 1962). Chronic physical conditions referred to having at least one of angina, arthritis, diabetes, or stroke.

Statistical analysis

The statistical analysis was performed with Stata 14.1 (Stata Corp LP, College station, Texas). The analysis was restricted to those aged ≥ 50 years. The difference in sample characteristics between those with and without near or far vision impairment was tested by Chi-squared tests and Student's *t*-tests for categorical and continuous variables, respectively. Country-wise multivariable logistic regression analysis was conducted to assess the association between near or far vision impairment (exposures) and low physical activity (outcome). Interaction analysis was also conducted to assess whether the strength of the association between near or far vision impairment and low physical activity differs by age

group (50-64 and ≥ 65 years) by including the product term of age group X (near or far) visual impairment in the model. In order to assess the between-country heterogeneity that may exist in the association between near or far vision impairment and low physical activity, we calculated the Higgins' I^2 based on estimates for each country. The Higgins' I^2 represents the degree of heterogeneity that is not explained by sampling error with a value of $<40\%$ often considered as negligible and 40-60% as moderate heterogeneity (Higgins & Thompson, 2002). A pooled estimate was obtained by fixed-effect meta-analysis as the level of between-country heterogeneity was low. Finally, we also assessed whether there is a dose-dependent association between severity of far vision impairment and low physical activity with multivariable logistic regression using the overall sample.

All regression analyses were adjusted for age, sex, wealth, education, smoking, obesity, and chronic physical condition. The analysis with near vision impairment as the exposure was additionally adjusted for far vision impairment, while that of far vision impairment was adjusted for near vision impairment. Furthermore, the analysis on severity of far vision impairment and low physical activity was adjusted for country by including dummy variables for each country in the model as in previous SAGE publications (Field, 2013; Koyanagi et al., 2018; Koyanagi et al., 2019). All variables were included in the models as categorical variables with the exception of age (continuous variable). The sample weighting and the complex study design (i.e., strata and primary sampling units) were taken into account in all analyses with the use of the `svy` command in Stata, which relies on the Taylor linearization method. Results from the regression analyses are presented as odds ratios (ORs) with 95% confidence intervals (CIs). The level of statistical significance was set at $P < 0.05$.

RESULTS

The final sample included 34,129 individuals aged ≥ 50 years (China 13,175; Ghana 4,305; India 6,560; Mexico 2,313; Russia 3,938; South Africa 3,838). The sample characteristics are provided in **Table 1**. Overall, the mean (SD) age of the sample was 62.4 (16.0) and 47.9% were males. The overall prevalence of low physical activity (i.e., not meeting the recommended WHO guidelines), near vision impairment, and far vision impairment were 23.5%, 39.5%, and 15.8%, respectively. Furthermore, 17.1%, 15.4%, and 0.4% had mild, moderate, and severe far vision impairment, respectively. Individuals with near or far vision impairment were more likely to be older, females, poorer, have lower levels of education, and have chronic physical conditions. Overall, the prevalence of low physical activity among those with and without near vision impairment was 25.7% and 21.6%, respectively, while the corresponding figures for far vision impairment were 31.4% and 21.8%, respectively (**Figure 1**). The country-wise association between near vision impairment and low physical activity estimated by multivariable logistic regression is shown in **Figure 2**. Near vision impairment was not significantly associated with low physical activity, with the overall estimate based on a meta-analysis being $OR=1.07$ ($95\%CI=[0.97-1.10]$, $I^2=0.0\%$). On the other hand, far vision impairment was significantly associated with low physical activity with the pooled estimate being $OR=1.32$ ($95\%CI=[1.17-1.49]$, $I^2=0.0\%$) (**Figure 3**). For both near and far vision, there was no significant interaction by age group. Finally, there was a dose-dependent increase in the odds for low physical activity with severity of far vision impairment. Specifically, compared to no far vision impairment, mild, moderate, and severe far vision impairment were associated with 1.09 ($95\%CI = [0.94-1.26]$), 1.31 ($95\%CI = [1.06-1.61]$), and 1.80 ($95\%CI = [1.03-3.15]$) times higher odds for low physical activity, respectively (**Figure 4**).

DISCUSSION

In this large representative sample of older adults from six LMICs across multiple continents, far vision impairment was significantly associated with low physical activity, in a dose dependent manner. Those with severe far vision impairment (vs. no vision impairment) were 1.80 times more likely to report low physical activity and not meet current physical activity recommendations. The finding that far vision impairment is associated with low physical activity is in line with previous studies using objective (Smith et al., 2019) and subjective measures of visual acuity (López-Sánchez et al., 2019; Smith et al., 2017; Smith et al., 2019; Willis et al., 2012) conducted in HICs. Our study adds to the previous literature by identifying for the first time that far vision impairment as confirmed by objective measures is associated with low physical activity in older adults in LMICs. Furthermore, we show for the first time that near vision impairment is not associated with low physical activity.

There are several plausible pathways that may explain the association between far vision impairment and low physical activity levels. First, to participate in many sporting activities, optimal vision is required and having far vision impairment will likely hinder one's ability to perform at optimal levels. For example, in relation to ball sports, not being able to see a ball until it is up close will significantly impair one's ability to respond. Moreover, if correctives (spectacles) are worn, it is possible that one would be put off playing particular sports in fear of the corrective being damaged. Although this could be overcome through contact lenses wear, in LMICs, it is likely that a small proportion of the population wear contact lenses. For example, in a study of adults from Ghana, just 34.8% of the sample were aware of contact lens wear for vision correction (Abokyi, Manuh, Otchere, & Ilechie, 2017). In relation to this, it is also possible that people with visual difficulties may lack access to recreational and athletic programmes, especially in LMICs where these types of programmes may be scarce.

Next, some areas of LMICs have a high crime rate and can be hostile environments (Wolf, Gray, & Fazel, 2014). Thus, those who have far vision impairment may be concerned about their personal safety when carrying out free-living physical activity, such as walking to a destination, as they may not be able to clearly see signs of danger ahead, and may therefore choose to stay at home more or use alternative modes of motorized transport that may be perceived to be safer. Also, it is possible that those with far visual impairment are more likely to be unemployed, while they may also have difficulty in engaging in social activities (Royal National Institute of Blind People, 2020). Thereby, they may lose the opportunity to engage in occupational physical activity and incidental physical activity acquired during social activities.

Given these plausible pathways in terms of the link between far vision impairment and low physical activity, it may not be of surprise that near vision impairment was not associated with low physical activity. It is possible that, although not investigated in the present study, near vision impairment is less likely to interfere with the above-mentioned factors that may lead to low physical activity. Future research of a qualitative nature is now required to identify barriers and facilitators to physical activity participation in older adults with vision impairments residing in LMICs.

It has previously been suggested that to increase levels of physical activity for those who have a disability in LMICs, awareness of the benefits of physical activity needs to be increased among health care providers in this setting. This could be achieved through continued medical education in relation to the importance of assessing levels of and promoting participation in physical activity. Moreover, it is stated that physical activity promotion in this setting should utilize cognitive behavior principles (e.g., goal setting and

problem solving). Finally, it would be prudent to increase ophthalmic infrastructure in relation to visual impairment and implement health policies in terms of glasses and contact lens awareness and distribution.

The use of a large nationally representative dataset across multiple LMICs is a clear strength of the present study. However, findings must be interpreted in light of the study limitations. First, physical activity was assessed using self-report, and this may have introduced some level of bias (e.g., recall bias). Future studies using device-based data (e.g. accelerometers) on physical activity from LMICs are warranted. Second, our study included six LMICs with large populations but our study cannot be considered to be representative of all LMICs. Finally, the study is of a cross-sectional nature and it is not known whether lower levels of physical activity precede far vision impairment or whether far vision impairment precedes low levels of physical activity. For example, some studies have shown that those who engage in high levels of physical activity may be less likely to develop myopia (Suhr Thykjær, Lundberg, & Grauslund, 2017).

In conclusion, in this large representative sample of older adults from multiple LMICs, those with objectively measured far vision impairment reported lower levels of physical activity. Future studies should aim to identify the factors that lead to low physical activity in people with far vision impairment in this setting. Interventions to address low levels of physical activity in the visually impaired in LMICs should target those with far vision impairment and tailor interventions to this population's specific needs. Such interventions may include medical education on the benefits of participation in physical activity, goal setting and problem solving. Moreover, a recent systematic review and meta-analysis on interventions to promote physical activity among those with vision impairment concluded that physical

activity interventions in individuals with visual impairment incorporating activities such as tai chi, yoga and dance can have positive results, particularly in physical measures such as mobility and balance (Sweeting et al., 2020).

References

- Abokyi, S., Manuh, G., Otchere, H., & Ilechie, A. (2017). Knowledge, usage and barriers associated with contact lens wear in ghana. *Contact Lens and Anterior Eye*, 40(5), 329-334.
- Bull, F. C., Maslin, T. S., & Armstrong, T. (2009). Global physical activity questionnaire (GPAQ): Nine country reliability and validity study. *Journal of Physical Activity and Health*, 6(6), 790-804.
- Capella-McDonnall, M. (2007). The need for health promotion for adults who are visually impaired. *Journal of Visual Impairment & Blindness*, 101(3), 133-145.
- Caspersen, C. J., Powell, K. E., & Christenson, G. M. (1985). Physical activity, exercise, and physical fitness: Definitions and distinctions for health-related research. *Public Health Reports*, 100(2), 126.
- Cunningham, C., O'Sullivan, R., Caserotti, P., & Tully, M. A. (2020). Consequences of physical inactivity in older adults: A systematic review of reviews and meta-analyses. *Scandinavian Journal of Medicine & Science in Sports*, 30(5), 816-827.
- Ehrlich, J. R., Stagg, B. C., Andrews, C., Kumagai, A., & Musch, D. C. (2019). Vision impairment and receipt of eye care among older adults in low-and middle-income countries. *JAMA Ophthalmology*, 137(2), 146-158.

- 316 Field, A. (2013). *Discovering statistics using IBM SPSS statistics*. Los Angeles: SAGE.
- 317 Freeman, E. E., Roy-Gagnon, M., Samson, E., Haddad, S., Aubin, M., Vela, C., &
318 Zunzunegui, M. V. (2013). The global burden of visual difficulty in low, middle, and
319 high income countries. *PloS One*, 8(5), e63315.
- 320 Higgins, J. P., & Thompson, S. G. (2002). Quantifying heterogeneity in a meta-analysis.
321 *Statistics in Medicine*, 21(11), 1539-1558.
- 322 Klaver, C. C., Wolfs, R. C., Vingerling, J. R., Hofman, A., & de Jong, P. T. (1998). Age-
323 specific prevalence and causes of blindness and visual impairment in an older
324 population: The rotterdam study. *Archives of Ophthalmology*, 116(5), 653-658.
- 325 Kowal, P., Chatterji, S., Naidoo, N., Biritwum, R., Fan, W., Lopez Ridaura, R., . . . Williams,
326 S. (2012). Data resource profile: The world health organization study on global AGEing
327 and adult health (SAGE). *International Journal of Epidemiology*, 41(6), 1639-1649.
- 328 Koyanagi, A., Lara, E., Stubbs, B., Carvalho, A. F., Oh, H., Stickley, A., . . . Vancampfort, D.
329 (2018). Chronic physical conditions, multimorbidity, and mild cognitive impairment in
330 low-and middle-income countries. *Journal of the American Geriatrics Society*, 66(4),
331 721-727.
- 332 Koyanagi, A., Oh, H., Vancampfort, D., Carvalho, A. F., Veronese, N., Stubbs, B., & Lara,
333 E. (2019). Perceived stress and mild cognitive impairment among 32,715 community-
334 dwelling older adults across six low-and middle-income countries. *Gerontology*, 65(2),
335 155-163.
- 336 López-Sánchez, G. F., Grabovac, I., Pizzol, D., Yang, L., & Smith, L. (2019). The
337 association between difficulty seeing and physical activity among 17,777 adults residing

- 338 in Spain. *International Journal of Environmental Research and Public Health*, 16(21),
339 4267.
- 340 Rose, G. A. (1962). The diagnosis of ischaemic heart pain and intermittent claudication in
341 field surveys. *Bulletin of the World Health Organization*, 27(6), 645.
- 342 Royal National Institute of Blind People. (2020). Employment status and sight loss. Retrieved
343 from
344 [https://www.rnib.org.uk/sites/default/files/Employment%20status%20and%20sight%20l](https://www.rnib.org.uk/sites/default/files/Employment%20status%20and%20sight%20loss%202017.pdf)
345 [oss%202017.pdf](https://www.rnib.org.uk/sites/default/files/Employment%20status%20and%20sight%20loss%202017.pdf)
- 346 Smith, L., Gardner, B., Fisher, A., & Hamer, M. (2015). Patterns and correlates of physical
347 activity behaviour over 10 years in older adults: Prospective analyses from the English
348 longitudinal study of ageing. *BMJ Open*, 5(4), e007423.
- 349 Smith, L., Jackson, S. E., Pardhan, S., López-Sánchez, G. F., Hu, L., Cao, C., . . . Firth, J.
350 (2019). Visual impairment and objectively measured physical activity and sedentary
351 behaviour in US adolescents and adults: A cross-sectional study. *BMJ Open*, 9(4),
352 e027267.
- 353 Smith, L., Pardhan, S., Gorely, T., Barnett, Y., Jacob, L., López-Sánchez, G. F., . . .
354 Koyanagi, A. (2021). Physical activity and visual difficulties in 36 low-and middle-
355 income countries. *Eye*, Online ahead of print.
- 356 Smith, L., Timmis, M. A., Pardhan, S., Latham, K., Johnstone, J., & Hamer, M. (2017).
357 Physical inactivity in relation to self-rated eyesight: Cross-sectional analysis from the
358 English longitudinal study of ageing. *BMJ Open Ophthalmology*, 1(1), e000046.

- 359 Sport England. (2019). Physical inactivity in later life to cost the NHS more than £1.3bn by
 360 2030. Retrieved from [https://sportengland-production-files.s3.eu-west-](https://sportengland-production-files.s3.eu-west-2.amazonaws.com/s3fs-public/20190509-10-today-launched-by-sport-england.pdf)
 361 [2.amazonaws.com/s3fs-public/20190509-10-today-launched-by-sport-england.pdf](https://sportengland-production-files.s3.eu-west-2.amazonaws.com/s3fs-public/20190509-10-today-launched-by-sport-england.pdf)
- 362 Suhr Thykjær, A., Lundberg, K., & Grauslund, J. (2017). Physical activity in relation to
 363 development and progression of myopia—a systematic review. *Acta Ophthalmologica*,
 364 95(7), 651-659.
- 365 Sweeting, J., Merom, D., Astuti, P. A. S., Antoun, M., Edwards, K., & Ding, D. (2020).
 366 Physical activity interventions for adults who are visually impaired: A systematic review
 367 and meta-analysis. *BMJ Open*, 10(2), e034036.
- 368 WHO. (2010). Global recommendations on physical activity for health. *Geneva: World*
 369 *Health Organization*. <https://www.who.int/publications/i/item/9789241599979>
- 370 Willis, J. R., Jefferys, J. L., Vitale, S., & Ramulu, P. Y. (2012). Visual impairment,
 371 uncorrected refractive error, and accelerometer-defined physical activity in the united
 372 states. *Archives of Ophthalmology*, 130(3), 329-335.
- 373 Wolf, A., Gray, R., & Fazel, S. (2014). Violence as a public health problem: An ecological
 374 study of 169 countries. *Social Science & Medicine*, 104, 220-227.
- 375 World Health Organization. (2019). Blindness and vision impairment. Retrieved from
 376 <https://www.who.int/news-room/fact-sheets/detail/blindness-and-visual-impairment>

377 **Table 1**

Table 1 Sample characteristics (overall and by near and far vision impairment)

		Near vision impairment				Far vision impairment		
Characteristic		Overall	No	Yes	P-value ^a	No	Yes	P-value ^a
Age (years)	Mean (SD)	62.4 (16.0)	61.1 (15.5)	63.8 (16.6)	<0.001	61.2 (15.4)	67.1 (17.0)	<0.001
Sex	Male	47.9	51.2	44.1	<0.001	49.8	40.6	<0.001
	Female	52.1	48.8	55.9		50.2	59.4	
Wealth	Poorest	17.1	16.3	18.9	<0.001	15.9	24.8	<0.001
	Poorer	19.0	18.3	20.7		18.9	21.2	
	Middle	19.5	19.6	19.8		19.2	22.4	
	Richer	21.3	22.1	20.3		22.3	16.2	
	Richest	23.1	23.7	20.3		23.7	15.3	
Education	Primary	57.4	55.9	66.5	<0.001	58.8	67.6	0.033
	Secondary	35.2	36.4	28.5		34.1	28.2	
	Tertiary	7.4	7.8	5.0		7.1	4.2	
Smoking	Never	58.6	57.5	58.4	0.621	57.8	58.9	0.811
	Current	34.9	36.1	35.0		35.7	34.8	
	Former	6.6	6.4	6.6		6.5	6.2	

Vision impairment and physical activity in adults

Obesity	No	88.5	90.0	89.9	0.910	90.0	90.3	0.776
	Yes	11.5	10.0	10.1		10.0	9.7	
Chronic physical condition	No	62.0	65.0	59.8	<0.001	64.7	53.6	<0.001
	Yes	38.0	35.0	40.2		35.3	46.4	

Abbreviation: SD Standard deviation

Data are % unless otherwise stated.

P-value was calculated by Chi-squared tests and Student's t-tests for categorical and continuous variables, respectively.

FIGURE LEGENDS

Figure 1 Prevalence of low physical activity by presence of absence of (A) near or (B) far vision impairment (overall and by country)

Abbreviation: VI Vision impairment

Bars denote 95% confidence interval.

Figure 2 Country-wise association between near vision impairment and low physical activity (outcome) estimated by multivariable logistic regression

Abbreviation: OR Odds ratio; CI Confidence interval

Models are adjusted for age, sex, wealth, education, smoking, obesity, chronic physical conditions, and far vision impairment.

Overall estimate was obtained by meta-analysis with fixed effects.

Figure 3 Country-wise association between far vision impairment and low physical activity (outcome) estimated by multivariable logistic regression

Abbreviation: OR Odds ratio; CI Confidence interval

Models are adjusted for age, sex, wealth, education, smoking, obesity, chronic physical conditions, and near vision impairment.


Overall estimate was obtained by meta-analysis with fixed effects.

Figure 4 Association between severity of far vision impairment and low physical activity (outcome) estimated by multivariable logistic regression

Reference category is no vision impairment (6/12 or better). Mild vision impairment = 6/18 or better but worse than 6/12; Moderate vision impairment = 6/60 or better but worse than 6/18; Severe vision impairment = worse than 6/60.

Models are adjusted for age, sex, wealth, education, smoking, obesity, chronic physical conditions, near vision impairment, and country.

(A) Near vision impairment



(B) Far vision impairment

