



The Long-Term Outcome and Rehabilitative Approach of Intraventricular Hemorrhage at Preterm Birth

Juntaek Hong, Dong-wook Rha

Department and Research Institute of Rehabilitation Medicine, Yonsei University College of Medicine, Seoul, Korea

Technological advances in neonatology led to the improvement of the survival rate in preterm babies with very low birth weights. However, intraventricular hemorrhage (IVH) has been one of the major complications of prematurity. IVH is relevant to neurodevelopmental disorders, such as cerebral palsy, language and cognitive impairments, and neurosensory and psychiatric problems, especially when combined with brain parenchymal injuries. Additionally, severe IVH requiring shunt insertion is associated with a higher risk of adverse neurodevelopmental outcomes. Multidisciplinary and longitudinal rehabilitation should be provided for these children based on the patients' life cycles. During the infantile period, it is essential to detect high-risk infants based on neuromotor examinations and provide early intervention as soon as possible. As babies grow up, close monitoring of language and cognitive development is needed. Moreover, providing continuous rehabilitation with task-specific and intensive repetitive training could improve functional outcomes in children with mild-to-moderate disabilities. After school age, maintaining the level of physical activity and managing complications are also needed.

Key Words : Cerebral intraventricular hemorrhage · Premature birth · Neurodevelopmental disorders · Rehabilitation.

INTRODUCTION

Technological advances in neonatology have improved the survival rate of preterm infants with low birth weights. Intraventricular hemorrhage (IVH) is a major complication of premature birth. Approximately 12000 infants are born annually with IVH only in the United States⁷⁾. The frequency of IVH is 10–20% in preterm infants born before 30 weeks of gestation, and the rate of severe IVH increases to 35–45% in these infants with less than 750 g of birth weight^{1,11)}. Germinal matrix hemorrhage-IVH is the most common variety of neonatal intracranial hemorrhage⁴⁸⁾. The germinal matrix is a highly vas-

cular structure that is vulnerable to hemorrhage, as the thickness of the germinal matrix decreases after 24 gestational weeks and almost disappears by 36–37 gestational weeks¹⁰⁾. Disturbance in cerebral blood flow and fragility of germinal matrix vasculature primarily contributes to the development of germinal matrix hemorrhage-IVH⁹⁾. IVH causes complications such as post-hemorrhagic ventricular dilatation (PHVD) and brain parenchymal lesions, whose incidence is proportional to the severity of IVH⁷⁶⁾. Even though there was no apparent parenchymal lesion, ventricular blood clots and associated inflammatory reaction caused by blood breakdown products in the ventricles may result in brain injury⁴²⁾. More-

• Received : November 29, 2022 • Revised : January 20, 2023 • Accepted : January 31, 2023

• Address for reprints : **Dong-wook Rha**

Department and Research Institute of Rehabilitation Medicine, Yonsei University College of Medicine, 50 Yonsei-ro, Seodaemun-gu, Seoul 03722, Korea
Tel : +82-2-2228-3717, Fax : +82-2-363-2795, E-mail : medicus@yonsei.ac.kr, rehabkr@gmail.com, ORCID : <https://orcid.org/0000-0002-7153-4937>

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

over, some cases of PHVD require a permanent ventriculo-peritoneal shunt¹. These structural brain lesions as well as the presence of a shunt are related to unfavorable neurodevelopmental outcomes^{71,91,100}. When these long-term sequelae remain, multimodal approaches including family and social support should be integrated²². Especially, the need for rehabilitation throughout the patients' life is becoming more important²⁶. In this review, we discuss the long-term effects of IVH in premature babies and rehabilitative approaches throughout their life.

LONG-TERM OUTCOME OF IVH AT PRETERM BIRTH BASED ON THE CLINICAL DISEASE ENTITY

The survival rate of preterm infants has improved over the past three decades^{5,31,93}. Ironically, the advances in neonatal intensive care improving the survival of extremely preterm infants have, by extension, led to a relative increase in the number of infants with IVH development⁸⁹. Many survivors diagnosed with IVH may live with a long-term neurodevelopmental disability, and the severity of the disability is proportional to the IVH grade^{32,78}. Moreover, severe IVH that requires shunt insertion poses a great risk for adverse neurodevelopmental outcomes when compared to IVH without a shunt¹. And prolonged and frequent treatments for shunt-related complications including revision could be one of the factors associated with increased disability and reduced quality of life^{51,60}. Meanwhile, approximately 20–30% of extreme preterm survivors (gestational age <28 weeks) with IVH were diagnosed with cerebral palsy (CP), 30–35% with minor motor dysfunction which was not enough to be diagnosed with CP, 20% with a major neurosensory disability, and 15% with impaired academic skills^{44,88}. A quarter of preterm survivors (gestational age <36 weeks) without neuromotor symptoms develop psychiatric disorders, such as major depressive disorder and obsessive-compulsive disorder⁸⁸. In this section, we describe the long-term outcomes of preterm babies diagnosed with IVH according to disease entities.

Cerebral palsy

CP is a group of permanent, but changing, disorders that affect the ability to move and maintain posture and balance

caused by non-progressive interference, lesions, or abnormality of the immature brain^{12,85}. There are many risk factors associated with the prevalence of CP, such as low birth weight and premature birth⁹⁶. In South Korea, the overall prevalence of CP was 6.2–6.6% in extremely low birth weight infants, and it had an inverse relationship with gestational age⁵⁴. IVH at preterm birth is one of the major causes of CP⁹². Overall, approximately 10–20% of preterm babies with IVH develop CP^{6,88}. The incidence of CP is related directly to the grade of IVH, and almost 60–100% of grade 4 IVH results in CP⁷². IVH may induce brain parenchymal lesions such as periventricular leukomalacia (PVL) or PHVD, or both. The association between IVH without parenchymal injury and CP incidence is still controversial⁷¹. In contrast, IVH with parenchymal injury is closely related to the prevalence and incidence of CP. PVL is a predominant form of brain injury and an independent risk factor for CP²⁸. In 50% of children with CP, CP is associated with PVL²⁹. Meanwhile, PHVD is also one of the main complication of IVH. When ventricular dilation is more severe and clinical symptoms develop (full fontanelle, splaying sutures, irritability, apneas, sunseting), the term post-hemorrhagic hydrocephalus is used⁵⁷. 60% of infants with PHVD become stable without surgical treatment or need transient treatment such as external ventricular drainage. About 25% of PHVD require permanent surgical intervention¹. Overall, CP is diagnosed in about 40% of infants with PHVD, and the risk factors for this poor developmental outcome are high-grade IVH and increased head circumference³².

CP usually involves a variety of neuromuscular problems that depend on the variable location and severity of brain lesions^{34,50}. These problems include not only muscle weakness but also spasticity, dystonia, poor balance, and loss of selective motor control. All of these are correlated in a complex manner, which disturbs children's gross motor function and performance of daily activities. Non-ambulatory children with CP reach their maximum gross motor function earlier than ambulatory children do. Age₉₀, the average age at which children achieve 90% of their expected limit in gross motor ability, significantly differs according to the Gross Motor Function Classification System (GMFCS) level. The GMFCS is a five-level scale that rates a child's gross motor function with an emphasis on movement initiation, sitting control, and walking. Level I represents the highest gross motor function, whereas level V represents the lowest. The average of Age₉₀ for

GMFCS I and II is between 4 and 5 years of age, less than 4 years for GMFCS III to IV, and 2.7 years for GMFCS V. Moreover, the children with GMFCS III-V reach their peak of gross motor function before adolescence (7 years 11 months, 6 years 11 months, and 6 years 11 months, respectively) which decline as they grow up^{41,84}.

Meanwhile, children with CP could have progressive musculoskeletal complications caused by various factors, mainly due to failure of longitudinal growth of skeletal muscle³⁶. Clinically, deficiency in muscle relaxation as well as reduced activity could induce contracture of the joints, especially at the ankle⁵⁰. The imbalance between the growth of long bones and muscle-tendon units itself could induce relative muscle shortening and limited range of motion¹⁰⁶. Moreover, bone and joint changes are usually progressive, especially for the spine and lower extremities, such as scoliosis, hip subluxation, and subluxation of the talonavicular joint⁶⁶, which could result in functional regression and chronic pain³⁷. As a result, children with CP have a considerably higher burden of medical, neurological, and mental/behavioral disorders than the non-CP population⁴⁵. Regular monitoring and management of musculoskeletal pain are also needed across the entire range of motor impairment⁷⁹.

Cognitive abnormality and language development without neuromotor symptoms

Children with a history of IVH at preterm birth have a higher risk of cognitive impairment than children born full-term. It has been reported that 35% of children with a history of IVH at preterm birth without brain parenchymal injury and up to 76% with injury, such as grade III-IV IVH, PVL, or ventriculomegaly, need special education whereas only 10% of children born full-term need it⁵⁸. IVH was reported to be an independent risk factor for cognitive development, with the severity of impairments being proportional to IVH grade⁸⁸. The total IQ was significantly lower in children with post-hemorrhagic ventricular dilatation, especially in those who had undergone surgical intervention such as ventriculoperitoneal shunt and external ventricular drainage, than in children with a history of preterm birth but no ventricular dilatation⁴⁶. In contrast, the presence of cognitive dysfunction was not associated with low-grade IVH^{55,88,98}. However, when the follow-up period is extended up to adolescence, low-grade IVH is also a risk factor for cognitive dysfunction^{7,73}.

Language disability is also common in children with a history of extremely preterm birth (32 weeks) and very low birth weight (<1500 g)⁸⁰, and functional communication skills in preterm toddlers with IVH could be impaired⁵².

The pathophysiological mechanism of these symptoms has not yet been established in children with IVH in the absence of brain parenchymal lesions. The preterm brain itself showed reduced white matter connectivity⁹⁴. However, a previous study analyzing brain images demonstrated that very premature infants with low-grade IVH may present with regional abnormalities, such as a decrease in the total volume of cortical gray matter associated with the visual association, functional anisotropy, and microstructural immaturity of major white matter tracts⁸. Further studies are needed to clarify the mechanism.

Other complications

Several other complications are associated with IVH. All grades of IVH correlate with a higher incidence of optic atrophy⁷⁰. Mild IVH (grade I or II) is associated with deleterious effects on cortical vision development and function based on visually evoked potential in infants with 5–7 months of corrective age⁹⁸. Infants with severe IVH (grade III or IV) showed a higher incidence of visual-motor coordination dysfunction⁹⁸. Moreover, in children who had no evident white matter damage, a history of IVH at preterm birth was associated with double the risk of microcephaly and impaired visual fixation. Children with IVH and white matter damage show a higher risk of visual field impairment⁷¹.

Hearing impairment is also one of the severe consequences of prematurity, and its prevalence is inversely proportional to gestational age¹⁰³. However, there was no evidence of increased risk of hearing impairment in preterm infants with any degree of IVH compared to preterm infants without IVH. However, in children with PVL, the risk of hearing impairment increases^{35,40}.

Epilepsy is one of the most common complications in extremely preterm infants. Seizure disorders were present in about 4–5% of preterm infants, and the incidence of epilepsy was inversely proportional to gestational age at birth^{99,102}. IVH with white matter injury at preterm birth and a history of neonatal convulsions were strongly associated with an increased risk of epilepsy. However, there have been no reports of epilepsy incidence in children with IVH without white

matter injury⁴³). Further studies are needed to accurately analyze the correlation between IVH and epilepsy.

Considering psychiatric disorders, children with IVH were three times more likely to develop autism spectrum disease, especially post-hemorrhagic ventricular dilatation in a low birth weight population⁶⁷. Moreover, IVH without brain parenchymal injury increased the risk of adolescent major depressive disorder and obsessive-compulsive disorder, whereas IVH with brain injury increased the risk of tic disorder, obsessive-compulsive disorder, and attention deficit hyperactivity disorder-inattentive type only¹⁰¹. In contrast, prematurity and low birth weight were significantly correlated with all types of attention deficit hyperactivity disorders at the school age²⁵.

REHABILITATIVE APPROACH : FROM EARLY DETECTION TO LONG-TERM CARE BASED ON LIFE CYCLE

For patients diagnosed with brain injury, including IVH, several variables could affect long-term outcomes. We should consider the severity of brain lesions, current level of function, degree of spasticity, and anticipatory growth of bone and muscle and accordingly provide multidisciplinary and lifelong rehabilitation plans. In this section, we review the rehabilitative approach according to age.

Early detection and intervention

For children with brain injuries, including IVH, a comprehensive and individualized rehabilitation plan based on the patient's life cycle is needed. Around birth, it is essential to detect high-risk babies and provide early interventions. Brain magnetic resonance imaging (MRI) is the most suitable modality for early evaluation of the site and severity of brain lesions¹⁵. Meanwhile, observing general movements (GMs) in preterm and term-born infants during their first few months also has similar sensitivity and specificity to detect neurological abnormalities as MRI⁶³. GMs are characterized as spontaneous gross movements of the entire body with no distinctive pattern or sequence in movement of each body part. Prechtl et al.⁷⁷ developed a highly predictive tool for later neurological impairments by observing these GMs, particularly for spastic CP³⁰ as well as the dyskinetic type. Moreover, the prediction was significantly enhanced when the GM assessment was

combined with MRI findings⁹⁰. Some studies claimed that the quality of GMs can also predict cognitive and behavioral development other than motor abnormalities^{16,19,39}. Other neurological examinations, such as the Hammersmith Infant Neurological Examination, could also play an important role in the diagnosis and prognosis of infants at risk of poor neurodevelopmental outcomes^{38,83}. Even after the early infant period, it is necessary to closely observe the delay of development in motor, language, cognitive, and socio-emotional functions.

After neurological impairments are detected, early intervention is important in terms of neural plasticity. In South Korea, rehabilitative support after discharge was provided to approximately 35–38% of very low birth weight infants⁵⁴. For infants who exhibit difficulty following commands for motor training, providing an enriched environment is important for maximizing neurological outcomes. A key feature of an enriched environment is the provision of environmental complexity with the enrichment of objects that provide a range of opportunities for visual, somatosensory, and olfactory stimulation. Providing enriched environments could promote structural and chemical brain plasticity in animal studies^{68,97}. Although the term “enriched environment” has been widely used, there is no agreed definition for an enriched environment for human infants. One previous study proposed the definition of “enriched environment” for infants, which aims to enrich the multiple aspects of the infant's environment to promote learning : e.g., interventions aiming to enhance parent-infant interaction and stimulate self-generated motor activity by adapting the physical and play environment⁶⁵. Recently, early intervention incorporating self-initiated movement was reported to have a positive effect on motor development as well as intellectual development in high-risk infants with brain injury^{56,64}.

Rehabilitative approach in toddlers and preschoolers

Language and cognitive development progress rapidly in toddlers and preschoolers²⁰. Moreover, children who have not been diagnosed with CP often show language and cognitive abnormalities at this age². Therefore, attention should be paid to language and cognitive development, especially in children without neuromotor symptoms.

Continuous rehabilitation should be provided to children with neuromotor impairment. Task-oriented and intensive repetitive motor training has been shown to improve motor function, especially in children with mild-to-moderate dis-

abilities^{49,87}. Even for children with severe disability, rehabilitation could have a potentially positive impact on pediatric CP survival rate through the complication management⁴⁷. In accordance with the principles of motor learning, various rehabilitation treatment methods, including conventional physical therapy, occupational therapy, hydrotherapy, and electrical stimulation therapy, have been developed to improve motor function and activities of daily living. Recently, advanced technologies have been applied to achieve more effective motor learning. For example, robot-assisted training has been introduced, and its effectiveness has been reported in many studies^{33,104}. Moreover, the application of virtual reality technologies improved not only motor function²⁴ but also cognitive function⁵³ by boosting motivation and participation⁵⁹.

Spasticity is one of the most common neuromotor symptoms that can lead to long-term complications. There are many modalities to reduce spasticity: from non-surgical management, such as conventional stretching, serial casting, and anti-spastic medication, to surgical management, such as insertion of an intrathecal baclofen pump, selective dorsal rhizotomy, and orthopedic surgery⁹⁵. One of the most effective non-surgical treatments for spasticity is intramuscular botulinum toxin injection. Botulinum toxin injection not only promotes selective motor control by reducing spasticity¹³ but also avoids the development of bony deformities secondary to abnormal muscle pull and contracted tendons. Botulinum toxin injection could delay or decrease the need for orthopedic or neurosurgical operations⁷⁴. Even after these complications are managed by surgical interventions, postoperative intensive rehabilitation would be very helpful not only for recovering pre-functional levels but also for improving functional outcomes^{4,61}.

Rehabilitative approach for adolescents and adults

After school age, long-term negative health consequences of inactivity could increase the risk of musculoskeletal complications such as scoliosis⁸⁶, osteoporosis, fracture, and arthritis⁶⁹, as well as systemic metabolic diseases such as sarcopenia, obesity⁷⁵, and cardiovascular disease⁶². Considering the influence of adolescent lifestyle on adult health status, the importance of interventions for adolescents cannot be overemphasized¹⁷. Nevertheless, across all ages and levels of motor function, young people with neuromotor impairments participated in 13–53% less habitual physical activity than able-bodied per-

sons²¹. Therefore, it is important to encourage physical activity through a multidisciplinary and integrated approach involving medical rehabilitation as well as social and family support^{18,82}.

Complication management is also needed. Patients with neuromotor disorders are very vulnerable to musculoskeletal chronic pain, which could adversely influence life satisfaction²⁷. Therefore, alleviating chronic pain is critical, and treatment for chronic pain includes pharmacological interventions such as conventional pain-modulating medication as well as new medications such as nerve growth factor and calcitonin gene-related peptide¹⁴. However, the most important factor for alleviating pain is the maintenance of a light level of physical activity, which is an efficient pain coping strategy, and perseverance in participation⁸¹.

NECESSITY OF A REGISTRY SYSTEM FOR HIGH-RISK BABIES DIAGNOSED WITH IVH

In South Korea, there have been some retrospective cross-sectional studies on the incidence, prevalence, and various characteristics of preterm children with IVH^{3,105}. Only a few studies have analyzed the outcomes of high-risk infants. However, they retrospectively analyzed the payment data of the Korean National Health Insurance, which was not collected for academic study²³. The Korean Neonatal Network⁵⁴, which is a web-based registry of very low birth weight infants formed through voluntary participation of hospitals nationwide from 2013 to 2016, has reported useful data to help improve the outcomes of premature infants and their health. However, we do not have a prospective registry system for children with IVH or CP which is a major neuromotor complication in preterm children with IVH. A few Western countries only have their own CP registry systems: The Cerebral Palsy Research Registry in the USA, the Surveillance of Cerebral Palsy in Europe, and the Australian Cerebral Palsy Register. These registry systems have provided informative data to build clinical practice guidelines and public healthcare policies to improve the health status of children with CP. However, there is no registry system for children with IVH. If we have our own registry system encompassing IVH and CP, it could be a cornerstone not only for the care of Korean children but also for East Asian children with IVH and/or CP.

CONCLUSION

IVH is a major complication of prematurity and can induce poor long-term developmental outcomes, including gross motor dysfunction, language and cognitive impairments, and psychiatric problems, especially in children with brain parenchymal injuries. Lifelong management is needed, which ranges from early detection and intervention to long-term care.

AUTHORS' DECLARATION

Conflicts of interest

No potential conflict of interest relevant to this article was reported.

Informed consent

This type of study does not require informed consent.

Author contributions

Conceptualization : JH, DR; Data curation : JH; Formal analysis : JH, DR; Methodology : JH, DR; Project administration : JH; Visualization : JH; Writing - original draft : JH; Writing - review & editing : DR

Data sharing

None

Preprint

None

ORCID

Juntaek Hong <https://orcid.org/0000-0002-1597-4137>
Dong-wook Rha <https://orcid.org/0000-0002-7153-4937>

References

1. Adams-Chapman I, Hansen NI, Stoll BJ, Higgins R; NICHD Research Network : Neurodevelopmental outcome of extremely low birth weight infants with posthemorrhagic hydrocephalus requiring shunt insertion. **Pediatrics** **121** : e1167-e1177, 2008
2. Adams-Chapman I, Heyne RJ, DeMauro SB, Duncan AF, Hintz SR, Pappas A, et al. : Neurodevelopmental impairment among extremely preterm infants in the neonatal research network. **Pediatrics** **141** : e20173091, 2018
3. Ahn SY, Shim SY, Sung IK : Intraventricular hemorrhage and post hemorrhagic hydrocephalus among very-low-birth-weight infants in Korea. **J Korean Med Sci** **30 Suppl 1(Suppl 1)** : S52-S58, 2015
4. Amen JF, El-Mikkawy DM, Yosry AH, El-Gebely MA, El-Sherbini MH : Outcome of intensive rehabilitation following single-event multilevel surgery for crouch gait in children with cerebral palsy. **ERAR** **46** : 78-84, 2019
5. Ancel PY, Goffinet F; EPIPAGE-2 Writing Group; Kuhn P, Langer B, Matis J, et al. : Survival and morbidity of preterm children born at 22 through 34 weeks' gestation in france in 2011: results of the epipage-2 cohort study. **JAMA Pediatr** **169** : 230-238, 2015
6. Ancel PY, Livinec F, Larroque B, Marret S, Arnaud C, Pierrat V, et al. : Cerebral palsy among very preterm children in relation to gestational age and neonatal ultrasound abnormalities: the epipage cohort study. **Pediatrics** **117** : 828-835, 2006
7. Ann Wy P, Rettiganti M, Li J, Yap V, Barrett K, Whiteside-Mansell L, et al. : Impact of intraventricular hemorrhage on cognitive and behavioral outcomes at 18 years of age in low birth weight preterm infants. **J Perinatol** **35** : 511-515, 2015
8. Argyropoulou MI, Astrakas LG, Xydis VG, Drougia A, Mouka V, Goel I, et al. : Is low-grade intraventricular hemorrhage in very preterm infants an innocent condition? Structural and functional evaluation of the brain reveals regional neurodevelopmental abnormalities. **AJNR Am J Neuroradiol** **41** : 542-547, 2020
9. Ballabh P : Intraventricular hemorrhage in premature infants: mechanism of disease. **Pediatr Res** **67** : 1-8, 2010
10. Ballabh P : Pathogenesis and prevention of intraventricular hemorrhage. **Clin Perinatol** **41** : 47-67, 2014
11. Bassan H : Intracranial hemorrhage in the preterm infant: understanding it, preventing it. **Clin Perinatol** **36** : 737-762, v, 2009
12. Bax M, Goldstein M, Rosenbaum P, Leviton A, Paneth N, Dan B, et al. : Proposed definition and classification of cerebral palsy, april 2005. **Dev Med Child Neurol** **47** : 571-576, 2005
13. Bjornson K, Hays R, Graubert C, Price R, Won F, McLaughlin JF, et al. : Botulinum toxin for spasticity in children with cerebral palsy: a comprehensive evaluation. **Pediatrics** **120** : 49-58, 2007
14. Blackman JA, Svensson CI, Marchand S : Pathophysiology of chronic pain in cerebral palsy: implications for pharmacological treatment and research. **Dev Med Child Neurol** **60** : 861-865, 2018
15. Bosanquet M, Copeland L, Ware R, Boyd R : A systematic review of tests to predict cerebral palsy in young children. **Dev Med Child Neurol** **55** : 418-426, 2013
16. Bruggink JL, Van Braeckel KN, Bos AF : The early motor repertoire of children born preterm is associated with intelligence at school age. **Pediatrics** **125** : e1356-e1363, 2010
17. Buffart LM, van den Berg-Emons RJ, van Wijlen-Hempel MS, Stam HJ, Roebroek ME : Health-related physical fitness of adolescents and young adults with myelomeningocele. **Eur J Appl Physiol** **103** : 181-

- 188, 2008
18. Bulekbayeva S, Daribayev Z, Ospanova S, Vento S : Cerebral palsy: a multidisciplinary, integrated approach is essential. **Lancet Glob Health** **5** : e401, 2017
 19. Butcher PR, van Braeckel K, Bouma A, Einspieler C, Stremmelaar EF, Bos AF : The quality of preterm infants' spontaneous movements: an early indicator of intelligence and behaviour at school age. **J Child Psychol Psychiatry** **50** : 920-930, 2009
 20. Capute AJ, Accardo PJ : The infant neurodevelopmental assessment: a clinical interpretive manual for cat-clams in the first two years of life, part 2. **Curr Probl Pediatr** **26** : 279-306, 1996
 21. Carlon SL, Taylor NF, Dodd KJ, Shields N : Differences in habitual physical activity levels of young people with cerebral palsy and their typically developing peers: a systematic review. **Disabil Rehabil** **35** : 647-655, 2013
 22. Cavonius-Rintahaka D, Aho AL, Voutilainen A, Billstedt E, Gillberg C : Health, functionality, and social support in families with a child with a neurodevelopmental disorder - a pilot study. **Neuropsychiatr Dis Treat** **15** : 1151-1161, 2019
 23. Cha JH, Ahn JH, Kim YJ, Lee BG, Kim JI, Park HK, et al. : Impact of preterm birth on neurodevelopmental disorders in South Korea: a nationwide population-based study. **J Clin Med** **11** : 2476, 2022
 24. Choi JY, Yi SH, Ao L, Tang X, Xu X, Shim D, et al. : Virtual reality rehabilitation in children with brain injury: a randomized controlled trial. **Dev Med Child Neurol** **63** : 480-487, 2021
 25. Chu SM, Tsai MH, Hwang FM, Hsu JF, Huang HR, Huang YS : The relationship between attention deficit hyperactivity disorder and premature infants in Taiwanese: a case control study. **BMC Psychiatry** **12** : 85, 2012
 26. Cieza A, Causey K, Kamenov K, Hanson SW, Chatterji S, Vos T : Global estimates of the need for rehabilitation based on the global burden of disease study 2019: a systematic analysis for the global burden of disease study 2019. **Lancet** **396** : 2006-2017, 2021
 27. Colver A, Rapp M, Eisemann N, Ehlinger V, Thyen U, Dickinson HO, et al. : Self-reported quality of life of adolescents with cerebral palsy: a cross-sectional and longitudinal analysis. **Lancet** **385** : 705-716, 2015
 28. Deng W, Pleasure J, Pleasure D : Progress in periventricular leukomalacia. **Arch Neurol** **65** : 1291-1295, 2008
 29. Drougia A, Giapros V, Krallis N, Theocharis P, Nikaki A, Tzoufi M, et al. : Incidence and risk factors for cerebral palsy in infants with perinatal problems: a 15-year review. **Early Hum Dev** **83** : 541-547, 2007
 30. Einspieler C, Cioni G, Paolicelli PB, Bos AF, Dressler A, Ferrari F, et al. : The early markers for later dyskinetic cerebral palsy are different from those for spastic cerebral palsy. **Neuropediatrics** **33** : 73-78, 2002
 31. EXPRESS Group, Fellman V, Hellström-Westas L, Norman M, Westgren M, Källén K, et al. : One-year survival of extremely preterm infants after active perinatal care in Sweden. **JAMA** **301** : 2225-2233, 2009
 32. Gilard V, Chadie A, Ferracci FX, Brasseur-Daudruy M, Proust F, Marret S, et al. : Post hemorrhagic hydrocephalus and neurodevelopmental outcomes in a context of neonatal intraventricular hemorrhage: an institutional experience in 122 preterm children. **BMC Pediatrics** **18** : 288, 2018
 33. Gilliaux M, Renders A, Dispa D, Holvoet D, Sapin J, Dehez B, et al. : Upper limb robot-assisted therapy in cerebral palsy: a single-blind randomized controlled trial. **Neurorehabil Neural Repair** **29** : 183-192, 2015
 34. Gormley ME Jr : Treatment of neuromuscular and musculoskeletal problems in cerebral palsy. **Pediatr Rehabil** **4** : 5-16, 2001
 35. Gotardo JW, Volkmer NFV, Stangler GP, Dornelles AD, Bohrer BBA, Carvalho CG : Impact of peri-intraventricular haemorrhage and periventricular leukomalacia in the neurodevelopment of preterms: a systematic review and meta-analysis. **PLoS One** **14** : e0223427, 2019
 36. Graham HK : Painful hip dislocation in cerebral palsy. **Lancet** **359** : 907-908, 2002
 37. Graham HK, Thomason P, Willoughby K, Hastings-Ison T, Stralen RV, Dala-Ali B, et al. : Musculoskeletal pathology in cerebral palsy: a classification system and reliability study. **Children (Basel)** **8** : 252, 2021
 38. Haataja L, Mercuri E, Guzzetta A, Rutherford M, Counsell S, Flavia Frisone M, et al. : Neurologic examination in infants with hypoxic-ischemic encephalopathy at age 9 to 14 months: use of optimality scores and correlation with magnetic resonance imaging findings. **J Pediatr** **138** : 332-337, 2001
 39. Hadders-Algra M, Groothuis AM : Quality of general movements in infancy is related to neurological dysfunction, adhd, and aggressive behaviour. **Dev Med Child Neurol** **41** : 381-391, 1999
 40. Han JH, Shin JE, Lee SM, Eun HS, Park MS, Park KI : Hearing impairments in preterm infants: factors associated with discrepancies between screening and confirmatory test results. **Neonatal Med** **27** : 126-132, 2020
 41. Hanna SE, Rosenbaum PL, Bartlett DJ, Palisano RJ, Walter SD, Avery L, et al. : Stability and decline in gross motor function among children and youth with cerebral palsy aged 2 to 21 years. **Dev Med Child Neurol** **51** : 295-302, 2009
 42. Hinson HE, Hanley DF, Ziai WC : Management of intraventricular hemorrhage. **Curr Neurol Neurosci Rep** **10** : 73-82, 2010
 43. Hirvonen M, Ojala R, Korhonen P, Haataja P, Eriksson K, Gissler M, et al. : The incidence and risk factors of epilepsy in children born preterm: a nationwide register study. **Epilepsy Res** **138** : 32-38, 2017
 44. Hollebrandse NL, Spittle AJ, Burnett AC, Anderson PJ, Roberts G, Doyle LW, et al. : School-age outcomes following intraventricular haemorrhage in infants born extremely preterm. **Arch Dis Child Fetal Neonatal Ed** **106** : 4-8, 2021
 45. Hollung SJ, Bakken IJ, Vik T, Lydersen S, Wiik R, Aaberg KM, et al. : Comorbidities in cerebral palsy: a patient registry study. **Dev Med Child Neurol** **62** : 97-103, 2020
 46. Holwerda JC, Van Braeckel KNJA, Roze E, Hoving EW, Maathuis CGB, Brouwer OF, et al. : Functional outcome at school age of neonatal post-hemorrhagic ventricular dilatation. **Early Hum Dev** **96** : 15-20, 2016
 47. Hsu CL, Hung CL, Huang SJ, Yin CH, Chu CH, Kuo TJ, et al. : Impact of rehabilitation intensity on 3-year mortality among children with moderate to severe cerebral palsy: a population-based cohort study. **Int J Environ Res Public Health** **18** : 9932, 2021
 48. Inder TE, Perlman JM, Volpe JJ : Chapter 24 - preterm intraventricular

- hemorrhage/posthemorrhagic hydrocephalus in Volpe JJ, Inder TE, Darvas BT, de Vries LS, du Plessis AJ, Neil JJ, et al. (eds) : **Volpe's neurology of the newborn (sixth edition)**. Amsterdam : Elsevier, 2018, pp637-698.e621
49. Katz-Leurer M, Rotem H, Keren O, Meyer S : The effects of a 'home-based' task-oriented exercise programme on motor and balance performance in children with spastic cerebral palsy and severe traumatic brain injury. **Clin Rehabil** 23 : 714-724, 2009
 50. Kerr Graham H, Selber P : Musculoskeletal aspects of cerebral palsy. **J Bone Joint Surg Br** 85 : 157-166, 2003
 51. Kulkarni AV, Shams I : Quality of life in children with hydrocephalus: results from the hospital for sick children, Toronto. **J Neurosurg** 107(5 Suppl) : 358-364, 2007
 52. Landry SH, Schmidt M, Richardson MA : The effects of intraventricular hemorrhage on functional communication skills in preterm toddlers. **J Dev Behav Pediatr** 10 : 299-306, 1989
 53. Larson EB, Feigon M, Gagliardo P, Dvorkin AY : Virtual reality and cognitive rehabilitation: a review of current outcome research. **NeuroRehabilitation** 34 : 759-772, 2014
 54. Lee JH, Youn Y, Chang YS : Short- and long-term outcomes of very low birth weight infants in Korea: Korean neonatal network update in 2019. **Clin Exp Pediatr** 63 : 284-290, 2020
 55. Legge N, Lutz T, Wocadlo C, Rieger I : Long-term neurodevelopmental outcome in preterm infants with intraventricular haemorrhage. **J Paediatr Child Health** 58 : 1797-1802, 2022
 56. Li X, Kang Y, Tang H : The effects of early rehabilitation in high-risk infants with brain injury: a systematic review and meta-analysis. **Transl Pediatr** 10 : 2467-2478, 2021
 57. Limbrick DD Jr, de Vries LS : New insights into the management of post-hemorrhagic hydrocephalus. **Semin Perinatol** 46 : 1515-1527, 2022
 58. Luu TM, Ment LR, Schneider KC, Katz KH, Allan WC, Vohr BR : Lasting effects of preterm birth and neonatal brain hemorrhage at 12 years of age. **Pediatrics** 123 : 1037-1044, 2009
 59. Maggio MG, Maresca G, De Luca R, Stagnitti MC, Porcari B, Ferrera MC, et al. : The growing use of virtual reality in cognitive rehabilitation: fact, fake or vision? A scoping review. **J Natl Med Assoc** 111 : 457-463, 2019
 60. Mansoor N, Solheim O, Fredrikli OA, Gulati S : Shunt complications and revisions in children: a retrospective single institution study. **Brain Behav** 11 : e2390, 2021
 61. McLaughlin J, Bjornson K, Temkin N, Steinbok P, Wright V, Reiner A, et al. : Selective dorsal rhizotomy: meta-analysis of three randomized controlled trials. **Dev Med Child Neurol** 44 : 17-25, 2002
 62. McPhee PG, Claridge EA, Noorduyn SG, Gorter JW : Cardiovascular disease and related risk factors in adults with cerebral palsy: a systematic review. **Dev Med Child Neurol** 61 : 915-923, 2019
 63. Morgan C, Crowle C, Goyen TA, Hardman C, Jackman M, Novak I, et al. : Sensitivity and specificity of general movements assessment for diagnostic accuracy of detecting cerebral palsy early in an Australian context. **J Paediatr Child Health** 52 : 54-59, 2016
 64. Morgan C, Darrah J, Gordon AM, Harbourne R, Spittle A, Johnson R, et al. : Effectiveness of motor interventions in infants with cerebral palsy: a systematic review. **Dev Med Child Neurol** 58 : 900-909, 2016
 65. Morgan C, Novak I, Badawi N : Enriched environments and motor outcomes in cerebral palsy: systematic review and meta-analysis. **Pediatrics** 132 : e735-e746, 2013
 66. Morrell DS, Pearson JM, Sauser DD : Progressive bone and joint abnormalities of the spine and lower extremities in cerebral palsy. **Radiographics** 22 : 257-268, 2002
 67. Movsas TZ, Pinto-Martin JA, Whitaker AH, Feldman JF, Lorenz JM, Korzeniewski SJ, et al. : Autism spectrum disorder is associated with ventricular enlargement in a low birth weight population. **J Pediatr** 163 : 73-78, 2013
 68. Nithianantharajah J, Hannan AJ : Enriched environments, experience-dependent plasticity and disorders of the nervous system. **Nat Rev Neurosci** 7 : 697-709, 2006
 69. O'Connell NE, Smith KJ, Peterson MD, Ryan N, Liverani S, Anokye N, et al. : Incidence of osteoarthritis, osteoporosis and inflammatory musculoskeletal diseases in adults with cerebral palsy: a population-based cohort study. **Bone** 125 : 30-35, 2019
 70. O'Keefe M, Kafil-Hussain N, Flitcroft I, Lanigan B : Ocular significance of intraventricular haemorrhage in premature infants. **Br J Ophthalmol** 85 : 357-359, 2001
 71. O'Shea TM, Allred EN, Kuban KC, Hirtz D, Specter B, Durfee S, et al. : Intraventricular hemorrhage and developmental outcomes at 24 months of age in extremely preterm infants. **J Child Neurol** 27 : 22-29, 2012
 72. Özek E, Kersin SG : Intraventricular hemorrhage in preterm babies. **Turk Pediatr Ars** 55 : 215-221, 2020
 73. Patra K, Wilson-Costello D, Taylor HG, Mercuri-Minich N, Hack M : Grades I-II intraventricular hemorrhage in extremely low birth weight infants: effects on neurodevelopment. **J Pediatr** 149 : 169-173, 2006
 74. Pavone V, Testa G, Restivo DA, Cannavò L, Condorelli G, Portinaro NM, et al. : Botulinum toxin treatment for limb spasticity in childhood cerebral palsy. **Front Pharmacol** 7 : 29, 2016
 75. Peterson MD, Gordon PM, Hurvitz EA : Chronic disease risk among adults with cerebral palsy: the role of premature sarcopenia, obesity and sedentary behaviour. **Obes Rev** 14 : 171-182, 2013
 76. Pinto C, Malik P, Desai R, Shelar V, Bekina-Sreenivasan D, Satnarine TA, et al. : Post-hemorrhagic hydrocephalus and outcomes amongst neonates with intraventricular hemorrhage: a systematic review and pooled analysis. **Cureus** 13 : e18877, 2021
 77. Prechtl HF, Einspieler C, Cioni G, Bos AF, Ferrari F, Sontheimer D : An early marker for neurological deficits after perinatal brain lesions. **Lancet** 349 : 1361-1363, 1997
 78. Radic JA, Vincer M, McNeely PD : Outcomes of intraventricular hemorrhage and posthemorrhagic hydrocephalus in a population-based cohort of very preterm infants born to residents of Nova Scotia from 1993 to 2010. **J Neurosurg Pediatr** 15 : 580-588, 2015
 79. Ramstad K, Jahnsen R, Skjeldal OH, Diseth TH : Characteristics of recurrent musculoskeletal pain in children with cerebral palsy aged 8 to 18 years. **Dev Med Child Neurol** 53 : 1013-1018, 2011
 80. Reidy N, Morgan A, Thompson DK, Inder TE, Doyle LW, Anderson PJ :

- Impaired language abilities and white matter abnormalities in children born very preterm and/or very low birth weight. **J Pediatr** **162** : 719-724, 2013
81. Riquelme I, do Rosário RS, Vehmaskoski K, Natunen P, Montoya P : Influence of chronic pain in physical activity of children with cerebral palsy. **NeuroRehabilitation** **43** : 113-123, 2018
 82. Rohrer-Baumgartner N, Holthe IL, Svendsen EJ, Røe C, Egeland J, Borgen IMH, et al. : Rehabilitation for children with chronic acquired brain injury in the child in context intervention (CICI) study: study protocol for a randomized controlled trial. **Trials** **23** : 169, 2022
 83. Romeo DM, Ricci D, Brogna C, Mercuri E : Use of the hammersmith infant neurological examination in infants with cerebral palsy: a critical review of the literature. **Dev Med Child Neurol** **58** : 240-245, 2016
 84. Rosenbaum PL, Walter SD, Hanna SE, Palisano RJ, Russell DJ, Raina P, et al. : Prognosis for gross motor function in cerebral palsy: creation of motor development curves. **JAMA** **288** : 1357-1363, 2002
 85. Sadowska M, Sarecka-Hujar B, Kopyta I : Cerebral palsy: current opinions on definition, epidemiology, risk factors, classification and treatment options. **Neuropsychiatr Dis Treat** **16** : 1505-1518, 2020
 86. Saito N, Ebara S, Ohotsuka K, Kumeta H, Takaoka K : Natural history of scoliosis in spastic cerebral palsy. **Lancet** **351** : 1687-1692, 1998
 87. Salem Y, Godwin EM : Effects of task-oriented training on mobility function in children with cerebral palsy. **NeuroRehabilitation** **24** : 307-313, 2009
 88. Sherlock RL, Anderson PJ, Doyle LW; Victorian Infant Collaborative Study Group : Neurodevelopmental sequelae of intraventricular haemorrhage at 8 years of age in a regional cohort of elbw/very preterm infants. **Early Hum Dev** **81** : 909-916, 2005
 89. Siffel C, Kistler KD, Sarda SP : Global incidence of intraventricular hemorrhage among extremely preterm infants: a systematic literature review. **J Perinat Med** **49** : 1017-1026, 2021
 90. Skiöld B, Eriksson C, Eliasson AC, Adén U, Vollmer B : General movements and magnetic resonance imaging in the prediction of neuromotor outcome in children born extremely preterm. **Early Hum Dev** **89** : 467-472, 2013
 91. Sobana M, Halim D, Aviani JK, Gamayani U, Achmad TH : Neurodevelopmental outcomes after ventriculoperitoneal shunt placement in children with non-infectious hydrocephalus: a meta-analysis. **Childs Nerv Syst** **37** : 1055-1065, 2021
 92. Stavsky M, Mor O, Mastrolia SA, Greenbaum S, Than NG, Erez O : Cerebral palsy-trends in epidemiology and recent development in prenatal mechanisms of disease, treatment, and prevention. **Front Pediatr** **5** : 21, 2017
 93. Stoll BJ, Hansen NI, Bell EF, Shankaran S, Laptook AR, Walsh MC, et al. : Neonatal outcomes of extremely preterm infants from the nichd neonatal research network. **Pediatrics** **126** : 443-456, 2010
 94. Thompson DK, Chen J, Beare R, Adamson CL, Ellis R, Ahmadzai ZM, et al. : Structural connectivity relates to perinatal factors and functional impairment at 7years in children born very preterm. **Neuroimage** **134** : 328-337, 2016
 95. Tilton AH : Management of spasticity in children with cerebral palsy. **Semin Pediatr Neurol** **11** : 58-65, 2004
 96. Van Naarden Braun K, Doernberg N, Schieve L, Christensen D, Goodman A, Yeargin-Allsopp M : Birth prevalence of cerebral palsy: a population-based study. **Pediatrics** **137** : 1-9, 2016
 97. van Praag H, Kempermann G, Gage FH : Neural consequences of environmental enrichment. **Nat Rev Neurosci** **1** : 191-198, 2000
 98. Vohr BR, Garcia-Coll C, Mayfield S, Brann B, Shaul P, Oh W : Neurologic and developmental status related to the evolution of visual-motor abnormalities from birth to 2 years of age in preterm infants with intraventricular hemorrhage. **J Pediatr** **115** : 296-302, 1989
 99. Vohr BR, Wright LL, Dusick AM, Mele L, Verter J, Steichen JJ, et al. : Neurodevelopmental and functional outcomes of extremely low birth weight infants in the national institute of child health and human development neonatal research network, 1993-1994. **Pediatrics** **105** : 1216-1226, 2000
 100. Vollmer B, Roth S, Riley K, Sellwood MW, Baudin J, Neville BG, et al. : Neurodevelopmental outcome of preterm infants with ventricular dilatation with and without associated haemorrhage. **Dev Med Child Neurol** **48** : 348-352, 2006
 101. Whitaker AH, Feldman JF, Lorenz JM, McNicholas F, Fisher PW, Shen S, et al. : Neonatal head ultrasound abnormalities in preterm infants and adolescent psychiatric disorders. **Arch Gen Psychiatry** **68** : 742-752, 2011
 102. Whitehead E, Dodds L, Joseph KS, Gordon KE, Wood E, Allen AC, et al. : Relation of pregnancy and neonatal factors to subsequent development of childhood epilepsy: a population-based cohort study. **Pediatrics** **117** : 1298-1306, 2006
 103. Wroblewska-Seniuk K, Greczka G, Dabrowski P, Szyfter-Harris J, Mazela J : Hearing impairment in premature newborns-analysis based on the national hearing screening database in poland. **PLoS One** **12** : e0184359, 2017
 104. Yazıcı M, Livanelioğlu A, Gücüyener K, Tekin L, Sümer E, Yakut Y : Effects of robotic rehabilitation on walking and balance in pediatric patients with hemiparetic cerebral palsy. **Gait Posture** **70** : 397-402, 2019
 105. Yim SY, Yang CY, Park JH, Kim MY, Shin YB, Kang EY, et al. : Korean database of cerebral palsy: a report on characteristics of cerebral palsy in South Korea. **Ann Rehabil Med** **41** : 638-649, 2017
 106. Ziv I, Blackburn N, Rang M, Koreska J : Muscle growth in normal and spastic mice. **Dev Med Child Neurol** **26** : 94-99, 1984