



## 저작자표시-비영리-변경금지 2.0 대한민국

이용자는 아래의 조건을 따르는 경우에 한하여 자유롭게

- 이 저작물을 복제, 배포, 전송, 전시, 공연 및 방송할 수 있습니다.

다음과 같은 조건을 따라야 합니다:



저작자표시. 귀하는 원저작자를 표시하여야 합니다.



비영리. 귀하는 이 저작물을 영리 목적으로 이용할 수 없습니다.



변경금지. 귀하는 이 저작물을 개작, 변형 또는 가공할 수 없습니다.

- 귀하는, 이 저작물의 재이용이나 배포의 경우, 이 저작물에 적용된 이용허락조건을 명확하게 나타내어야 합니다.
- 저작권자로부터 별도의 허가를 받으면 이러한 조건들은 적용되지 않습니다.

저작권법에 따른 이용자의 권리는 위의 내용에 의하여 영향을 받지 않습니다.

이것은 [이용허락규약\(Legal Code\)](#)을 이해하기 쉽게 요약한 것입니다.

[Disclaimer](#)

**The consecutive 200 cases of endoscopic-  
combined intrarenal surgery: comparison  
between standard and miniature  
surgeries**

Young Joon Moon

Department of Medicine

The Graduate School, Yonsei University

# **The consecutive 200 cases of endoscopic- combined intrarenal surgery: comparison between standard and miniature surgeries**


Directed by Professor Joo Yong Lee

The Master's Thesis  
submitted to the Department of Medicine,  
the Graduate School of Yonsei University  
in partial fulfillment of the requirements for the degree of  
Master of Medical Science

Young Joon Moon

December 2023

This certifies that the Master's Thesis of  
Young Joon Moon is approved.



-----  
Thesis Supervisor : Joo Yong Lee



-----  
Thesis Committee Member#1 : Kang Su Cho



-----  
Thesis Committee Member#2 : Dae Chul Jung

The Graduate School  
Yonsei University

December 2023

## ACKNOWLEDGEMENTS

Thanks for everyone who contributed to writing this paper.

## <TABLE OF CONTENTS>

|  |     |
|--|-----|
| ABSTRACT .....   | iii |
| I. INTRODUCTION .....  | 1   |
| II. MATERIALS AND METHODS .....  | 2   |
| 1. ECIRS Patient Population and Technique .....                                  | 2   |
| 2. Baseline Patient and Stone Characteristics .....                              | 3   |
| 3. Statistical Analyses .....  | 4   |
| 4. Ethics Statement .....  | 4   |
| III. RESULTS .....   | 5   |
| 1. Baseline Characteristics and Treatment Outcomes .....                         | 5   |
| 2. Comparison of Propensity Score-Matched Standard ECIRS and Mini-ECIRS<br>..... | 6   |
| 3. Predictive Factors of Successful ECIRS .....                                  | 8   |
| IV. DISCUSSION .....   | 8   |
| V. CONCLUSION .....  | 15  |
| REFERENCES .....   | 16  |
| APPENDICES .....   | 21  |
| ABSTRACT(IN KOREAN) .....  | 22  |
| PUBLICATION LIST .....   | 24  |

## LIST OF TABLES

|  |   |
|--|---|
| Table 1. Baseline Characteristics and Treatment Outcomes of standard ECIRS vs. mini-ECIRS .....  | 5 |
| Table 2. Baseline Characteristics and Treatment Outcomes of standard ECIRS vs. mini-ECIRS Groups after Propensity Score Matching using MSL, MSD, SHI, VCSD and LCD ..... | 7 |
| Table 3. Logistic Regression Analyses of Factors Associated with Successful ECIRS .....  | 8 |

## ABSTRACT

### **The consecutive 200 cases of endoscopic-combined intrarenal surgery: comparison between standard and miniature surgeries**

Young Joon Moon

*Department of Medicine  
The Graduate School, Yonsei University*

(Directed by Professor Joo Yong Lee)

**Introduction:** Percutaneous nephrolithotomy (PCNL) is still the gold-standard treatment for large and/or complex renal stones. Endoscopic combined intrarenal surgery (ECIRS) was developed with the goal of minimizing the number of access tracts of PCNL while simultaneously improving the one-step stone free rate (SFR). The aim of this study was to share the experience of the consecutive 200 cases of ECIRS in one institute and analyze surgical outcomes of mini-ECIRS and standard ECIRS.

**Methods:** We performed ECIRS for 200 adult patients between July 2017 and January 2020. An ECIRS was performed with the patient under general anesthesia in the intermediate-supine position. Surgeries were finished using tubeless technique with a simple ureteral stent insertion.

**Results:** There were significant differences in mean maximal stone length (MSL), variation coefficient of stone density (VCSD), linear calculus density (LCD), Seoul National University Renal Stone Complexity (S-ReSC), and modified S-ReSC scores in stone characteristics and estimated blood loss (EBL) and operation time in peri-operative outcomes between conventional and mini-ECIRS. After propensity-score matching, there was only difference in EBL two groups. In logistic regression models, MSL [odds ration (OR) 0.953; 95% confidence interval (CI) 0.926-0.979;  $P < 0.001$ ], LCD (OR 4.702; 95%



CI 1.613-18.655;  $P=0.013$ ) were significant factors for success rate after ECIRS.

**Conclusion:** In patients underwent mini-ECIRS, stones were relatively smaller and less complex and operation time was shorter. However, if the size of stones was similar, there was no difference in success rate, but EBL can be lower in mini-ECIRS than standard surgery.

---

Key words : kidney; nephrolithiasis; lithotripsy; urinary calculi

**The consecutive 200 cases of endoscopic-combined intrarenal surgery:  
comparison between standard and miniature surgeries**

Young Joon Moom

*Department of Medicine  
The Graduate School, Yonsei University*

(Directed by Professor Joo Yong Lee)

## I. INTRODUCTION

Renal stone is a common disease in urology, and to date, the prevalence is increasing worldwide[1]. Especially, complex renal stone is defined as having multiple stones or having anatomical or functional abnormalities. If complex renal stone is not treated, it can cause serious complications such as renal failure, sepsis, and septic shock, thus appropriate treatment is essential. Complex renal stone is a challenging case to manage in the urology field. In guidelines, percutaneous nephrolithotomy (PCNL) is recommended as the primary option in the treatment of large and/or complex renal stones[2,3]. However, El-Nahas AR et al. reported that the stone free rate (SFR) of PCNL monotherapy in the treatment of staghorn stones was only 56%[4]. In the treatment of patients with staghorn stones, performing PCNL using multiple tracts or sessions can achieve a high SFR, but there is a problem of increased related complications[5]. For this reason, after PCNL was first introduced by Fernstrom et al. in 1976[6], various developments, such as minimally invasive PCNL, have been made to reduce complications and increase SFR.

Endoscopic combined intrarenal surgery (ECIRS) was started with the goal of reducing complications and improving safety by reducing the number of access tracts while simultaneously increasing SFR[7]. ECIRS was first reported by Scoffone et al. in 2008[8,9].

It simultaneously combines PCNL and retrograde intrarenal surgery (RIRS) to explore whole urinary tract including renal cavities and aims to resolve urolithiasis with one-step surgical treatment[10]. The major advantages of using simultaneous antegrade-retrograde approach for complex renal calculi are better SFRs and lower bleeding complications by avoiding multiple tract PCNL. Until recently, many studies have been conducted and reported on clinical results and safety of ECIRS[11]. In addition, not only is radiation exposure less than standard PCNL, but it can also be a new alternative for treating stones in patients with renal congenital anomalies[12]. In 2014, mini-ECIRS was first reported by Hamamoto et al.[13]. Although the implementation of mini-ECIRS has increased since then, there is still a lack of studies on the indications, clinical outcomes, and complications of mini-ECIRS and comparative studies with standard ECIRS. The purpose of this study was to report the experience of the consecutive 200 cases of ECIRS in one institute and analyze clinical outcomes of mini-ECIRS and standard ECIRS.

## II. MATERIALS AND METHODS

### 1. ECIRS Patient Population and Technique

We retrospectively analyzed 200 consecutive ECIRS cases between July 2017 to January 2020 at one institution (Severance Hospital, Seoul, Korea). Renal calyceal puncture was performed by interventional radiologist using ultrasonic guidance before surgery or performed by urologic surgeon using ultrasonic and fluoroscopic guidance during the surgery. ECIRS was performed with the intermediate-supine [Galdakao modified supine Valdivia (GMSV)] position under general anesthesia. First, an Amplatz Super Stiff guidewire was indwelled under cystoscopic guidance, and then an 11/13-Fr ureteral access sheath (Uropass; Olympus Corp., Tokyo, Japan or Navigator, Boston Scientific) was inserted through the guidewire. Next, the collecting system was identified using a flexible ureteroscope [LithoVue™ single-use digital flexible ureteroscope (Boston Scientific)] and the puncture point was visualized. If renal calyceal puncture was not

performed before surgery, ultrasonic and fluoroscopic guidance puncture was performed by a urologist at this point. Afterwards, the formed tract was dilated with balloon dilator or one-step dilator. Fifty-five patients underwent standard ECIRS using 20-Fr nephroscope and 145 underwent mini-ECIRS with 12-Fr mini-nephroscope. Lithotripsy was performed with a holmium:YAG laser lithotripter (VersaPulse™ PowerSuite™ 100W; Lumenis, Tel Aviv, Israel). All cases were performed simultaneously using a flexible ureteroscope and tubeless technique with a simple ureteral stent indwelling.

## 2. Baseline Patient and Stone Characteristics

Patient characteristics, detailed history of urolithiasis, and stone characteristics including maximal stone length (MSL), mean stone density (MSD), stone heterogeneity index (SHI), variation coefficient of stone density (VCSD), and linear calculus density (LCD) were collected for analysis. Non-contrast computed tomography (NCCT) images of all patients were acquired and analyzed to collect stone characteristic data. MSD was defined as the mean Hounsfield units (HU) over the region of interest, and was measured using magnified axial NCCT images at level of the largest stone diameter. SHI was defined as the standard deviation of HU for the same region of interest. Stone complexity was assessed and measured using the Seoul National University Renal Stone Complexity (S-ReSC) and modified S-ReSC scores[14,15]. Stone analysis method consisted of quantitative analysis of stone composition through Fourier transform infrared (FTIR) spectroscopy, and stones were sent to the GC laboratory in Yongin, Korea for analysis. The stone composition data was stratified following the Mayo Clinic classification[16]. The stones were categorized into six groups: (1) calcium oxalate (CaOx) monohydrate group (stones containing >50% CaOx with or without any hydroxyapatite); (2) CaOx dihydrate group (stones containing >50% CaOx with or without any hydroxyapatite); (3) struvite group (stones containing any amount of struvite); (4) cystine group (stones containing any amount of cystine); (5) uric acid group (stones containing any amount of uric acid); and (6) calcium phosphate (CaP) group (stones containing >50% carbonate apatite and stones

containing any amount of brushite). To evaluate treatment results, complication rate and post-treatment complication data were collected including changes of hemoglobin, creatinine, and estimated glomerular filtration rate. Successful treatment was assessed 3 months after treatment and was defined as the absence of residual stones and the presence of residual fragments sized 3 mm or less on NCCT images.

### 3. Statistical Analyses

In comparing the basic characteristics of patients, Student's two-sample t-test and Pearson chi-square test with Yates' continuity correction were used. In addition, univariate and multivariate logistic regression analyzes were performed to identify factors associated with successful treatment. Welch's two sample t-test was performed to confirm homoscedasticity between the two groups, and the Shapiro-Wilk test was performed to analyze the normality of each group. Additionally, the Mann-Whitney U test and Wilcoxon rank-sum test were performed to confirm the independence of both groups. After comparing the variances of two samples from normal populations, the presence or absence of variation equality was confirmed in a t-test.

To further specify the characteristics of the subjects, we performed propensity score matching. Each mini-ECIRS patient was matched 1:1 with standard-ECIRS patient. Mini-ECIRS patient and standard-ECIRS patient each conducted 1:1 matching, and logistic regression with multivariate analysis using MSL, MSD, SHI, VCSD and LCD was used to calculate propensity scores. In order to present a better comparison group, propensity score matching is used to improve matching according to stone characteristics[17]. Regarding the propensity scores between the two groups, pretreatment characteristics distribution did not differ[18].

All statistical analyzes in this study were performed using R software (version 4.1.3).

### 4. Ethics Statement

The Institutional Review Board of our medical institution reviewed and approved the protocol for this study (Approval No. 2022-3041-002).

### III. RESULTS

#### 1. Baseline Characteristics and Treatment Outcomes

Of the 200 ECIRS patients, 120 patients were male and 80 were female. The mean age was  $58.49 \pm 13.46$  years. MSL was  $29.76 \pm 14.35$  mm. The average MSD was  $995.12 \pm 366.80$  HU and mean SHI was  $209.28 \pm 104.45$  HU. VCSD was  $21.61 \pm 20.98$  and LCD was  $0.98 \pm 1.37$ . The S-ReSC score was  $4.49 \pm 2.63$  and the average modified S-ReSC score was  $6.68 \pm 3.34$ . Mean operative time was  $79.69 \pm 32.50$  minutes, and the patient was discharged on postoperative day 2 (IQR 1.75-3). Twenty-two patients underwent simultaneous ipsilateral ECIRS and contralateral RIRS, and 11 patients underwent ipsilateral ureteroscopic ureterolithotomy. There were significant differences in MSL, VCSD, LCD, S-ReSC, and modified S-ReSC scores in stone characteristics and estimated blood loss (EBL) and operation time in peri-operative outcomes between conventional and mini-ECIRS (Table 1). The results of stone composition analysis were as follows: struvite group 40%, uric acid group 22%, CaOx monohydrate group 20.5%, CaOx dihydrate group 8.5%, CaP group 8.5%, and cystine group 0.5%.

**Table 1.** Baseline Characteristics and Treatment Outcomes of standard ECIRS vs. mini-ECIRS

|                          | Total             | Standard ECIRS    | Mini-ECIRS        | P-value |
|--------------------------|-------------------|-------------------|-------------------|---------|
| No.                      | 200               | 55                | 145               |         |
| Age (yr)                 | $58.49 \pm 13.46$ | $57.75 \pm 16.17$ | $58.77 \pm 12.32$ | 0.671   |
| Sex                      |                   |                   |                   | 1.000   |
| Female                   | 80 (40.00%)       | 22 (40.00%)       | 58 (40.00%)       |         |
| Male                     | 120 (60.00%)      | 33 (60.00%)       | 87 (60.00%)       |         |
| BMI (kg/m <sup>2</sup> ) | $24.93 \pm 5.26$  | $24.77 \pm 3.06$  | $24.99 \pm 5.89$  | 0.732   |

|                |              |                |               |        |
|----------------|--------------|----------------|---------------|--------|
| MSL (mm)       | 29.76±14.35  | 36.95±14.93    | 27.04±13.18   | <0.001 |
| MSD (HU)       | 995.12±366.8 | 1047.05±404.92 | 975.43±350.75 | 0.218  |
| SHI (HU)       | 204.28±104.4 | 188.80±103.91  | 210.15±104.41 | 0.198  |
| VCSD           | 21.61±20.98  | 17.34±6.62     | 23.23±24.13   | 0.008  |
| LCD            | 0.98±1.37    | 0.57±0.37      | 1.13±1.57     | <0.001 |
| S-ReSC         | 4.49±2.63    | 5.62±2.33      | 4.06±2.61     | <0.001 |
| MoS-ReSC       | 6.68±3.34    | 7.82±3.27      | 6.25±3.28     | 0.003  |
| Multiplicity   |              |                |               | 1.000  |
| No             | 38 (19.00%)  | 10 (18.18%)    | 28 (19.31%)   |        |
| Yes            | 162 (81.00%) | 45 (81.82%)    | 117 (80.69%)  |        |
| Bilateral      |              |                |               | 0.276  |
| No             | 128 (64.00%) | 39 (70.91%)    | 89 (61.38%)   |        |
| Yes            | 72 (36.00%)  | 16 (29.09%)    | 56 (38.62%)   |        |
| EBL            | 27.88±71.53  | 56.27±110.78   | 17.10±45.21   | 0.014  |
| Operation time | 79.69±32.50  | 91.67±36.62    | 75.15±29.68   | 0.001  |
| Success        |              |                |               | 0.158  |
| No             | 24 (12.00%)  | 10 (18.18%)    | 14 (9.66%)    |        |
| Yes            | 176 (88.00%) | 45 (81.82%)    | 131 (90.34%)  |        |

ECIRS, endoscopic combined intrarenal surgery; MSL, maximal stone length; MSD, mean stone density; HU, Hounsfield units; SHI, stone heterogeneity index; VCSD, Variation coefficient of stone density; LCD, linear calculus density; S-ReSC, Seoul National University Renal Stone Complexity; MoS-ReSC, modified S-ReSC; EBL, estimated blood loss.

Data are presented as means±standard deviations, numbers (percentage)

## 2. Comparison of Propensity Score-Matched Standard ECIRS and Mini-ECIRS

As a result of propensity score matching, the number of patients in each group was 37, and there were no significant differences in age ( $p=0.499$ ), sex ( $p=0.811$ ), MSL ( $p=0.856$ ), MSD ( $p=0.880$ ), or SHI ( $p=0.927$ ) between the two groups. Also, there were no significant differences in S-ReSC and modified S-ReSC scores between two groups ( $p=0.737$ ;  $p=0.744$ ). After propensity score matching, only EBL showed a significant difference

between the two groups, being significantly higher in standard ECIRS than mini ECIRS ( $p=0.010$ ) (Table 2).

**Table 2.** Baseline Characteristics and Treatment Outcomes of standard ECIRS vs. mini-ECIRS Groups after Propensity Score Matching using MSL, MSD, SHI, VCSD and LCD

|                          | Total         | Standard ECIRS | Mini-ECIRS     | P-value |
|--------------------------|---------------|----------------|----------------|---------|
| No.                      | 74            | 37             | 37             |         |
| Age (yr)                 | 57.99±13.82   | 59.08±15.62    | 56.89±11.86    | 0.499   |
| Sex                      |               |                |                | 0.811   |
| Female                   | 28 (37.84%)   | 15 (40.54%)    | 13 (35.14%)    |         |
| Male                     | 46 (62.16%)   | 22 (59.46%)    | 24 (64.86%)    |         |
| BMI (kg/m <sup>2</sup> ) | 25.40±3.56    | 24.84±3.05     | 25.95±3.97     | 0.182   |
| MSL (mm)                 | 30.48±11.87   | 30.74±10.22    | 30.23±13.46    | 0.856   |
| MSD (HU)                 | 1010.95±355.8 | 1017.24±343.57 | 1004.66±372.31 | 0.880   |
| SHI (HU)                 | 191.09±100.39 | 192.18±99.85   | 190.01±102.29  | 0.927   |
| VCSD                     | 18.23±6.38    | 18.32±6.76     | 18.13±6.06     | 0.895   |
| LCD                      | 0.70±0.39     | 0.68±0.39      | 0.71±0.39      | 0.733   |
| S-ReSC                   | 4.69±2.40     | 4.78±2.03      | 4.59±2.74      | 0.737   |
| MoS-ReSC                 | 6.85±3.17     | 6.73±3.03      | 6.97±3.35      | 0.744   |
| Multiplicity             |               |                |                | 0.247   |
| No                       | 15 (20.27%)   | 10 (27.03%)    | 5 (13.51%)     |         |
| Yes                      | 59 (79.73%)   | 27 (72.97%)    | 32 (86.49%)    |         |
| EBL                      | 39.32±94.77   | 67.84±125.50   | 10.81±28.12    | 0.010   |
| Operation time           | 79.95±28.34   | 82.59±30.32    | 77.30±26.37    | 0.425   |
| Success                  |               |                |                | 1.000   |
| No                       | 12 (16.22%)   | 6 (16.22%)     | 6 (16.22%)     |         |
| Yes                      | 62 (83.78%)   | 31 (83.78%)    | 31 (83.78%)    |         |

ECIRS, endoscopic combined intrarenal surgery; MSL, maximal stone length; MSD, mean stone density; HU, Hounsfield units; SHI, stone heterogeneity index; VCSD, Variation coefficient of stone density; LCD, linear calculus density; S-ReSC, Seoul National University Renal Stone Complexity; MoS-ReSC, modified



S-ReSC; EBL, estimated blood loss.

Data are presented as means±standard deviations, numbers (percentage)

### 3. Predictive Factors of Successful ECIRS

In logistic regression models, MSL [odds ratio (OR) 0.953; 95% confidence interval (CI) 0.926-0.979;  $p<0.001$ ], LCD (OR 4.702; 95% CI 1.613-18.655;  $p=0.013$ ) were significant factors for success rate after ECIRS (Table 3).

**Table 3.** Logistic Regression Analyses of Factors Associated with Successful ECIRS

| Variables  | Odds ratio | 95% CI       | P-value |
|------------|------------|--------------|---------|
| Age        | 1.016      | 0.984-1.048  | 0.318   |
| BMI        | 0.982      | 0.919-1.070  | 0.602   |
| MSL        | 0.953      | 0.926-0.979  | <0.001  |
| MSD        | 0.999      | 0.997-1.000  | 0.171   |
| SHI        | 0.999      | 0.995-1.004  | 0.841   |
| VCSD       | 1.014      | 0.988-1.071  | 0.573   |
| LCD        | 4.702      | 1.613-18.655 | 0.013   |
| S-ReSC     | 0.865      | 0.738-1.012  | 0.069   |
| MoS-ReSC   | 0.890      | 0.781-1.011  | 0.075   |
| Mini-ECIRS | 2.080      | 0.843-4.983  | 0.103   |

ECIRS, endoscopic combined intrarenal surgery; MSL, maximal stone length; MSD, mean stone density; HU, Hounsfield units; SHI, stone heterogeneity index; VCSD, Variation coefficient of stone density; LCD, linear calculus density; S-ReSC, Seoul National University Renal Stone Complexity; MoS-ReSC, modified S-ReSC; EBL, estimated blood loss.

## IV. DISCUSSION

Renal stone is a common disorder, and the prevalence has been increasing worldwide[1]. In the United States, it is reported that 5% of the population is affected by nephrolithiasis[19]. Renal staghorn stones account for 10-20% of all renal stones and are a

serious disease that, if left untreated, can lead to renal failure, severe infection, septic shock, and even death. Therefore, treatment of these complex renal stones is essential to prevent complications, and as it is a challenging treatment in the field of urology, many studies are being conducted[20]. The European Association of Urology (EAU) and American Urological Association (AUA) guidelines for the management of renal calculi > 20 mm recommend PCNL as the first-line therapy[2, 3, 21].

PCNL was first reported by Fernstrom et al. in 1976[6, 22], and it showed relatively high SFRs for partial and complete staghorn stones[23]. As PCNL has been widely performed, various patient positions have been suggested and Ibarluzea et al. first reported PCNL using a intermediate-supine (GMSV) position in 1992[24]. Additionally, much effort has been made to increase the stone free rate in the treatment of complex renal stones, and as part of that effort, multitract PCNL using multiple PCN tracts has been performed. Multitract PCNL has the advantage of relatively efficiently increasing the stone free rate by accessing stones through various routes, but its major disadvantages are increased blood loss and renal parenchymal injury caused by the increased number of punctures. Chen et al.[20] reported a single-center experience comparing multitract mini-PCNL and ECIRS in the treatment of renal staghorn stones. From January 2018 to September 2021, a retrospective analysis was conducted on a total of 34 patients, including 17 ECIRS patients and 17 multitract mini-PCNL patients. There was no significant difference in stone-free rate and mean operative time between the two groups ( $p=0.94$  and  $p=0.63$ , respectively). There were no significant differences between the two groups in complications including hemoglobin loss and postoperative blood transfusion rate. However, postoperative pain was significantly lower in the ECIRS group ( $p<0.001$ ). In conclusion, they described that both ECIRS and multitract mini-PCNL were effective and safe in the treatment of renal staghorn stones, and that ECIRS showed less postoperative pain.

However, PCNL is not the only option for the treatment of patients with a greater stone burden. In 1992, Ibarluzea et al.[24] proposed a method of performing PCNL for the treatment of large renal stones while simultaneously removing stone fragments through an

Amplatz catheter using a ureteroscope with clear vision. Improvements in surgical techniques and the development of endourologic devices have led to significant advances in the treatment of complex renal stones. RIRS plays a major role in reducing complications and improving clinical outcomes through combination with PCNL[25]. Later in 2008, Scoffone et al.[9] coined the term ECIRS and operated under the GMSV position. SFR of ECIRS was higher than PCNL monotherapy, and due to these advantages, it has been in the spotlight as a new treatment method for complex renal stones.

Recently, mini-ECIRS using mini-PCNL for safer and more effective ECIRS has been reported. Kwon et al.[26] conducted prospectively study to analyze the feasibility of carrying out ECIRS with supine mini-PCNL and RIRS for patients with bilateral stones in a single session. The study was conducted on a total of 26 patients. The SFR was 76.9% and 92.3% on the ECIRS side and the contralateral RIRS side, respectively. Complications occurred in two patients, but they completely improved with appropriate medical treatment after surgery. Accumulation of surgical experience (OR 117.3,  $p=0.046$ ) was a significant predictor of stone-free status. They concluded that carrying out ECIRS and contralateral RIRS in a single session for patients with bilateral stones is feasible and safe.

In our prior study, we retrospectively analyzed the first 100 cases of ECIRS in Korea and compared outcomes with shock-wave lithotripsy (SWL)[27]. ECIRS had a higher SFR and success rate compared to SWL. As in this study, multivariate logistic regression analysis was performed to identify predictors of successful ECIRS, and the results showed that smaller stone size (OR: 0.947,  $p=0.002$ ) and lower S-ReSC score (OR: 0.759,  $p=0.011$ ) were independent predictors of successful ECIRS.

Until recently, many studies comparing clinical outcomes of ECIRS and PCNL for complex renal stones. Wen et al.[28] conducted systematic review and meta-analysis to analyze the clinical outcomes of simultaneous mini-PCNL combined with flexible ureteroscopy as single-stage therapy to treat the partial staghorn calculi, and compared with conventional mini-PCNL monotherapy. Sixty-seven patients with partial renal staghorn stone were randomly divided into mini-PCNL Group and ECIRS Group, respectively. The

ECIRS group showed a significantly higher one-step SFR (87.88% vs 58.82%,  $p=0.007$ ) than the mini-PCNL group. And there was no statistical difference in clinical complications between the two groups ( $p=0.409$ ). They concluded that the simultaneous combined mini-PCNL and flexible ureteroscopy was more effective and safer than conventional mini-PCNL monotherapy in the treatment of partial staghorn stones. Gauhar et al.[29] conducted a systematic review and meta-analysis comparing ECIRS and conventional PCNL in the treatment of renal stones. Analysis was conducted on a total of 2,054 patients from 17 studies, including 800 ECIRS patients and 1,254 conventional PCNL patients. A meta-analysis was performed on four categories of parameters: surgical time and length of stay, bleeding, infection complications, stone-free rate and retreatment. There was no significant difference in surgical time, mean postoperative length, blood transfusion rate, and postoperative sepsis between the two groups ( $p=0.20$ ,  $p=0.69$ ,  $p=0.15$ , and  $p=0.25$ , respectively). Mean hemoglobin drop and retreatment rate were lower in the ECIRS group ( $p=0.03$ ,  $p=0.002$ , respectively). And stone-free rate was significantly higher in the PCNL group ( $p<0.0001$ ). In conclusion, although conventional PCNL showed a higher stone-free rate, ECIRS showed a shorter operative time, lower complication rate, and retreatment, thus supporting the contention that ECIRS is an effective and safe treatment method.

There was a study comparing the treatment outcomes between standard ECIRS and mini-ECIRS. Usui et al.[30] retrospectively analyzed 77 patients who underwent single session mini ECIRS and 77 patients who underwent single session standard ECIRS between April 2009 and May 2016. The SFR was similar for mini-ECIRS and standard ECIRS at 61.1% and 52.0% ( $p=0.388$ ), respectively. In terms of postoperative complications, Clavien-Dindo classification grade 2 or higher complications were 19.5% and 26.0% in the mini-ECIRS and standard ECIRS groups, respectively ( $p=0.442$ ), and severe complications of Clavien-Dindo classification grade 3 or higher were 2.6% and 3.9%, respectively ( $p>0.99$ ), there was no significant difference between the two groups. In addition, there was no significant difference in bleeding-related complications at 2.6% and 6.5% ( $p=0.442$ ), respectively, but pseudoaneurysm and significant blood loss was not

observed in the mini-ECIRS group. They concluded that compared with standard ECIRS, mini-ECIRS has the potential to reduce surgery-related pain and bleeding-related complications without increasing perioperative complications while maintaining SFR.

In our study, ECIRS was performed under the intermediate-supine (GMSV) position. Recently, many studies have been conducted and reported on patient position when performing ECIRS. Wang et al.[31] reported their experience with ECIRS using a modified prone split-leg position. Analysis was conducted on a total of 96 patients from September 2017 to January 2021. The stone-free rate was 78.1%, and the complication rate was 7.1%. Most complications were Clavien-Dindo classification grade 1 or 2, and no Clavien-Dindo classification grade 3 complications were observed. They also analyzed the risk factors associated with the ECIRS stone-free rate and found that the number of calyces in which the stone was located was the only significant risk factor ( $p=0.01$ ). In conclusion, they stated that ECIRS using a modified prone split-leg position is effective and safe for the treatment of renal stones. In addition, the ease of retrograde approach was cited as an advantage of ECIRS through modified prone split-leg position.

In order to reduce complications when performing ECIRS, it is important to reduce intraoperative EBL. To reduce EBL during surgery, renal calyceal puncture process is important. In our study, renal calyceal puncture was performed under ultrasonic and fluoroscopic guidance. Inoue et al.[32] reported the wideband Doppler ultrasound-guided puncture for safe renal calyceal puncture. They performed a retrospective analysis on a total of 41 patients who underwent mini-ECIRS using wideband Doppler ultrasound for the treatment of renal stones larger than 30 mm. Renal calyceal puncture using wideband Doppler ultrasound was successfully performed in all cases. Initial stone-free rate was 73.2% and final stone-free rate after additional treatment was 97.5%. Mean hemoglobin drop was  $0.54 \pm 0.65$  g/dL and three patients developed fever after surgery. However, no major complications such as injury to adjacent organs occurred. Univariate analysis was performed to identify risk factors for a severe bleeding (hemoglobin drop of  $\geq 1$  g/dL), and as a result of the analysis, there were no significant risk factors. They concluded that mini-

ECIRS using wideband Doppler ultrasound-guided renal puncture could be a safe and effective treatment option for the treatment of large renal stones. Next, nephrostomy tract dilation is also an important process for reducing EBL during surgery. In our study, nephrostomy tract dilation was performed in one step using a balloon dilator or one-step dilator. The nephrostomy tract dilation procedure can be performed using either a multi-step serial dilation technique or a single step dilation technique. However, it has not yet been proven which of the two techniques is superior in terms of safety. Gupta et al.[33] reported a prospective study comparing the single-step dilation technique and serial dilation technique when performing PCNL. A study was conducted on a total of 100 patients, including 50 patients using the single-step dilation technique and 50 patients using the serial dilation technique. Mean access tract dilation time and mean total fluoroscopy time were significantly shorter in the one-step dilation group ( $p < 0.0001$ ). However, there was no significant difference in requirement of blood transfusion and post-operative complications rate between the two groups. They concluded that the single-step dilation technique showed clinically comparable results when compared to the conventional serial dilation technique and was therefore a safe, effective, and economical method. They also recommended that the method be selected according to the surgeon's preference and experience.

Recently, a method using an antegrade flexible ureteroscopy to increase the stone-free rate when performing ECIRS was reported. Yang et al.[34] introduced a new technique named 'Through-through' approach. The process of this technique is to first use a nephroscope to identify the orientation of the target calyx and then insert a flexible ureteroscope through the nephroscope instrument channel to identify and remove residual stones. They performed a retrospective analysis on a total of 68 patients who underwent ECIRS using a 'Through-through' approach. Mean operative time was  $100.1 \pm 18.0$  minutes, and Stone free rate was 91.2%. Complications of Clavien-Dindo classification grade 3 or higher were not observed, and no patients required blood transfusion after surgery. Ten patients developed fever after surgery, but none developed septic shock.

Based on these results, they concluded that ECIRS using a ‘Through-through’ approach is a safe and effective treatment method for the treatment of complex renal stones.

Recently, with the advancement of science and technology, many changes have occurred in medical care, and new medical technologies are being introduced as robotic technology is incorporated into medicine. Recently, with the advancement of science and technology, many changes have occurred in medical care, and new medical technologies are being introduced as robotic technology is incorporated into medicine. Robotic technology is also being introduced in endourology, and study on the treatment of renal stones using robotic technology is also being published. Tokatli et al.[35] reported a retrospective study on the treatment of complex renal stones by robot-assisted mini-ECIRS using Avicenna Roboflex™. The study was conducted on a total of 42 patients. The stone-free rate was 95.5%, and complications were observed in 7.1%, and all complications were Clavien-Dindo classification grade 1. In conclusion, they stated that robot-assisted mini-ECIRS is effective and safe in the treatment of complex renal stones. In addition, the possibility of endoscopic flexible ureteroscopic evaluation of the collecting system at the end of surgery was described as an important advantage of robot-assisted mini-ECIRS. Although additional research is necessary in the future, ECIRS using a robotic system is also considered to be a good option for the treatment of renal stones.

What is unique about our study is that this is the first study to compare surgical outcomes between standard ECIRS and mini-ECIRS. We analyzed surgical outcomes in the consecutive 200 cases of ECIRS and compared between standard ECIRS and mini-ECIRS by performing propensity score matching using MSL, MSD, SHI, VCSD and LCD. Also, we performed logistic regression analyses to identify factors significantly associated with successful treatment. This study result may be helpful for urologists in selection for the management of renal stones.

Our study had several limitations. First, complications between standard ECIRS and mini-ECIRS were not compared. Analysis of complications is an important element in evaluating the safety of each procedure, so analysis will be necessary in future studies.

Second, there was no classification or analysis of the location of the renal stones, which may affect the success rate. Instead, we evaluated S-ReSC and modified S-ReSC scores that can predict the SFR after PCNL and RIRS, respectively. Since the S-ReSC score indicates the distribution complexity of renal stones, it is thought that it can overcome this limitation to some extent. In addition to S-ReSC, there are several scoring systems such as Stone-Tract length-Obstruction-Number of involved calyces, and Essence of stone (S.T.O.N.E.) nephrolithometry, Guy's stone score (GSS), and the Clinical Research Office of the Endourological Society (CROES) nomogram, but according to previous studies, there was no significant difference in predictive accuracy[36]. Lastly, small sample size and retrospective analysis are limitations of our study. In the future, a prospective randomized trial with a larger sample size could further increase the reliability of our study results.

## V. CONCLUSION

In patients underwent mini-ECIRS, stones were relatively smaller and less complex and operation time was shorter. However, if the size of stones was similar, there was no difference in success rate, but EBL can be lower in mini-ECIRS than standard surgery.



## REFERENCES

1. Romero, V.; Akpınar, H.; Assimos, D.G. Kidney stones: a global picture of prevalence, incidence, and associated risk factors. *Rev Urol* **2010**, *12*, e86-96.
2. Türk, C.; Petřík, A.; Sarica, K.; Seitz, C.; Skolarikos, A.; Straub, M.; Knoll, T. EAU Guidelines on Interventional Treatment for Urolithiasis. *Eur Urol* **2016**, *69*, 475-482, doi:10.1016/j.eururo.2015.07.041.
3. Assimos, D.; Krambeck, A.; Miller, N.L.; Monga, M.; Murad, M.H.; Nelson, C.P.; Pace, K.T.; Pais, V.M., Jr.; Pearle, M.S.; Preminger, G.M.; et al. Surgical Management of Stones: American Urological Association/Endourological Society Guideline, PART I. *J Urol* **2016**, *196*, 1153-1160, doi:10.1016/j.juro.2016.05.090.
4. el-Nahas, A.R.; Eraky, I.; Shokeir, A.A.; Shoma, A.M.; el-Assmy, A.M.; el-Tabey, N.A.; Soliman, S.; Elshal, A.M.; el-Kappany, H.A.; el-Kenawy, M.R. Factors affecting stone-free rate and complications of percutaneous nephrolithotomy for treatment of staghorn stone. *Urology* **2012**, *79*, 1236-1241, doi:10.1016/j.urology.2012.01.026.
5. Ramón de Fata, F.; Pérez, D.; Resel-Folkersma, L.; Galán, J.A.; Serrano, A.; Servera, A.; Arrabal, M.; Álvarez-Ossorio, J.L.; Ballester, R.; Cao, E.; et al. Analysis of the factors affecting blood loss in percutaneous nephrolithotomy: a registry of the Spanish Association of Urology in the supine position. *Actas Urol Esp* **2013**, *37*, 527-532, doi:10.1016/j.acuro.2013.05.001.
6. Fernström, I.; Johansson, B. Percutaneous pyelolithotomy. A new extraction technique. *Scand J Urol Nephrol* **1976**, *10*, 257-259, doi:10.1080/21681805.1976.11882084.
7. Abdullatif, V.A.; Sur, R.L.; Abdullatif, Z.A.; Szabo, S.R.; Abbott, J.E. The Safety and Efficacy of Endoscopic Combined Intrarenal Surgery (ECIRS) versus Percutaneous Nephrolithotomy (PCNL): A Systematic Review and Meta-Analysis. *Adv Urol* **2022**, *2022*, 1716554, doi:10.1155/2022/1716554.

8. Li, J.K.; Teoh, J.Y.; Ng, C.F. Updates in endourological management of urolithiasis. *Int J Urol* **2019**, *26*, 172-183, doi:10.1111/iju.13885.
9. Scoffone, C.M.; Cracco, C.M.; Cossu, M.; Grande, S.; Poggio, M.; Scarpa, R.M. Endoscopic combined intrarenal surgery in Galdakao-modified supine Valdivia position: a new standard for percutaneous nephrolithotomy? *Eur Urol* **2008**, *54*, 1393-1403, doi:10.1016/j.eururo.2008.07.073.
10. Ping, H.; Zhang, J.H.; Wang, M.S.; Xing, N.Z. Endoscopic Combined Intrarenal Surgery for the Treatment of Postpercutaneous Nephrolithotomy Residual Stones. *Chin Med J (Engl)* **2016**, *129*, 2885-2887, doi:10.4103/0366-6999.194659.
11. Abouelgreed, T.A.; Abdelaal, M.A.; Amin, M.M.; Elatreisy, A.; Shalkamy, O.; Abdrabuh, A.M.; Ghoneimy, O.M.; Aboutaleb, H. Endoscopic combined intrarenal surgery in the prone split-leg position versus Galdakao-modified supine Valdivia position for the management of partial staghorn calculi. *BMC Urol* **2022**, *22*, 163, doi:10.1186/s12894-022-01115-3.
12. Esperto, F.; Juliebø-Jones, P.; Keller, E.X.; Tailly, T.; Mykoniatas, I.; V, D.E.C.; Pietropaolo, A.; Talso, M.; Sener, E.T.; Tonyali, S.; et al. Urolithiasis in anomalous kidneys: RIRS or PCNL, this is the question. May ECIRS be the answer? *Minerva Urol Nephrol* **2022**, *74*, 802-804, doi:10.23736/s2724-6051.22.05208-9.
13. Hamamoto, S.; Yasui, T.; Okada, A.; Taguchi, K.; Kawai, N.; Ando, R.; Mizuno, K.; Kubota, Y.; Kamiya, H.; Tozawa, K.; Kohri, K. Endoscopic combined intrarenal surgery for large calculi: simultaneous use of flexible ureteroscopy and mini-percutaneous nephrolithotomy overcomes the disadvantageous of percutaneous nephrolithotomy monotherapy. *J Endourol* **2014**, *28*, 28-33, doi:10.1089/end.2013.0361.
14. Jeong, C.W.; Jung, J.W.; Cha, W.H.; Lee, B.K.; Lee, S.; Jeong, S.J.; Hong, S.K.; Byun, S.S.; Lee, S.E. Seoul National University Renal Stone Complexity Score for Predicting Stone-Free Rate after Percutaneous Nephrolithotomy. *PLoS One* **2013**, *8*, e65888, doi:10.1371/journal.pone.0065888.

15. Jung, J.W.; Lee, B.K.; Park, Y.H.; Lee, S.; Jeong, S.J.; Lee, S.E.; Jeong, C.W. Modified Seoul National University Renal Stone Complexity score for retrograde intrarenal surgery. *Urolithiasis* **2014**, *42*, 335-340, doi:10.1007/s00240-014-0650-7.
16. Lieske, J.C.; Rule, A.D.; Krambeck, A.E.; Williams, J.C.; Bergstralh, E.J.; Mehta, R.A.; Moyer, T.P. Stone composition as a function of age and sex. *Clin J Am Soc Nephrol* **2014**, *9*, 2141-2146, doi:10.2215/cjn.05660614.
17. Austin, P.C. A critical appraisal of propensity-score matching in the medical literature between 1996 and 2003. *Stat Med* **2008**, *27*, 2037-2049, doi:10.1002/sim.3150.
18. Rubin, D.B.; Thomas, N. Matching using estimated propensity scores: relating theory to practice. *Biometrics* **1996**, *52*, 249-264.
19. Stamatelou, K.K.; Francis, M.E.; Jones, C.A.; Nyberg, L.M.; Curhan, G.C. Time trends in reported prevalence of kidney stones in the United States: 1976-1994. *Kidney Int* **2003**, *63*, 1817-1823, doi:10.1046/j.1523-1755.2003.00917.x.
20. Chen, Z.H.; Lee, K.H.; Tseng, W.H.; Su, C.C.; Hsieh, K.L.; Lim, C.Y.; Huang, S.K. Comparison of mini endoscopic combined intrarenal surgery and multitract minimally invasive percutaneous nephrolithotomy specifically for kidney staghorn stones: a single-centre experience. *BMC Urol* **2022**, *22*, 93, doi:10.1186/s12894-022-01030-7.
21. Jung, H.D.; Lee, J.Y.; Kang, D.H.; Ko, K.; Koh, D.H.; Kwon, O.; Koo, K.C.; Kim, K.T.; Kim, M.S.; Kim, B.S.; et al. Korean Society of Endourology and Robotics (KSER) recommendation on the diagnosis, treatment, and prevention of urolithiasis. *Investigative and clinical urology* **2023**, *64*, 325-337, doi:10.4111/icu.20230102.
22. Inoue, T.; Okada, S.; Hamamoto, S.; Yoshida, T.; Matsuda, T. Current trends and pitfalls in endoscopic treatment of urolithiasis. *Int J Urol* **2018**, *25*, 121-133, doi:10.1111/iju.13491.
23. Diri, A.; Diri, B. Management of staghorn renal stones. *Ren Fail* **2018**, *40*, 357-362, doi:10.1080/0886022x.2018.1459306.
24. Ibarluzea, G.; Scoffone, C.M.; Cracco, C.M.; Poggio, M.; Porpiglia, F.; Terrone, C.; Astobieta, A.; Camargo, I.; Gamarra, M.; Tempia, A.; et al. Supine Valdivia and

- modified lithotomy position for simultaneous antegrade and retrograde endourological access. *BJU Int* **2007**, *100*, 233-236, doi:10.1111/j.1464-410X.2007.06960.x.
25. Zhao, F.; Li, J.; Tang, L.; Li, C. A comparative study of endoscopic combined intrarenal surgery (ECIRS) in the galdakao-modified supine valdivia (GMSV) position and minimally invasive percutaneous nephrolithotomy for complex nephrolithiasis: a retrospective single-center study. *Urolithiasis* **2021**, *49*, 161-166, doi:10.1007/s00240-020-01207-5.
  26. Kwon, O.; Park, J.; Cho, M.C.; Son, H.; Jeong, H.; Cho, S.Y. Feasibility of single-session endoscopic combined intrarenal surgery for ipsilateral large renal stones and retrograde intrarenal surgery for contralateral renal stones: Initial experience. *Int J Urol* **2017**, *24*, 377-382, doi:10.1111/iju.13313.
  27. Jung, H.D.; Moon, Y.J.; Almujaalhem, A.J.; Alqahtani, A.A.; Alkhureeb, M.A.; Lee, J.Y. The First 100 Cases of Endoscopic Combined Intrarenal Surgery in Korea: Matched Cohort Analyses versus Shock-Wave Lithotripsy. *Yonsei Med J* **2022**, *63*, 440-445, doi:10.3349/ymj.2022.63.5.440.
  28. Wen, J.; Xu, G.; Du, C.; Wang, B. Minimally invasive percutaneous nephrolithotomy versus endoscopic combined intrarenal surgery with flexible ureteroscope for partial staghorn calculi: A randomised controlled trial. *Int J Surg* **2016**, *28*, 22-27, doi:10.1016/j.ijssu.2016.02.056.
  29. Gauhar, V.; Castellani, D.; Cracco, C.M.; Scoffone, C.M.; Lim, E.J.; Rubilotta, E.; Pretore, E.; Pirola, G.M.; Maggi, M.; Rice, P.; et al. Is endoscopic combined intrarenal surgery ready for primetime in endourology? Outcomes from a systematic review and meta-analysis. *Cent European J Urol* **2022**, *75*, 171-181, doi:10.5173/cej.2022.0049.
  30. Usui, K.; Komeya, M.; Taguri, M.; Kataoka, K.; Asai, T.; Ogawa, T.; Yao, M.; Matsuzaki, J. Minimally invasive versus standard endoscopic combined intrarenal surgery for renal stones: a retrospective pilot study analysis. *Int Urol Nephrol* **2020**, *52*, 1219-1225, doi:10.1007/s11255-020-02433-x.

31. Wang, D.; Sun, H.; Xie, D.; Liu, Z.; Yu, D.; Ding, D. Application of a new position in endoscopic combined intrarenal surgery: modified prone split-leg position. *BMC Urol* **2022**, *22*, 38, doi:10.1186/s12894-022-00994-w.
32. Inoue, T.; Kinoshita, H.; Okada, S.; Hamamoto, S.; Taguchi, M.; Murota, T.; Matsuda, T. Wideband Doppler Ultrasound-guided Mini-endoscopic Combined Intrarenal Surgery as an Effective and Safe Procedure for Management of Large Renal Stones: A Preliminary Report. *Urology* **2016**, *95*, 60-66, doi:10.1016/j.urology.2016.05.038.
33. Gupta, N.K.; Huda, N.A.; Pal, D.K. A comparative study between one step dilatation and serial dilatation technique in percutaneous nephrolithotomy. *Urologia* **2023**, 3915603231202839, doi:10.1177/03915603231202839.
34. Yang, L.; Yin, Z.; Liu, H.; Wu, X. Initial Experience of Simultaneous Combined use of Percutaneous Nephrolithotomy and Flexible Ureteroscopy in Complex Renal Calculi: A Novel Surgical Technique of 'Through-Through' Approach. *Surg Innov* **2023**, *30*, 428-431, doi:10.1177/15533506231165827.
35. Tokatli, Z.; Ibis, M.A.; Sarica, K. Robot-Assisted Mini-Endoscopic Combined Intrarenal Surgery for Complex and Multiple Calculi: What Are the Real Advantages? *J Laparoendosc Adv Surg Tech A* **2022**, *32*, 890-895, doi:10.1089/lap.2022.0124.
36. Al Adl, A.M.; Mohey, A.; Abdel Aal, A.; Abu-Elnasr, H.A.F.; El Karamany, T.; Noureldin, Y.A. Percutaneous Nephrolithotomy Outcomes Based on S.T.O.N.E., GUY, CROES, and S-ReSC Scoring Systems: The First Prospective Study. *J Endourol* **2020**, *34*, 1223-1228, doi:10.1089/end.2019.0856.

## APPENDICES

## ABSTRACT(IN KOREAN)

**내시경-결합 신장 내 결석 수술 200례: 표준 경피적 신절석술과 미니 경피적 신절석술의 비교**

&lt; 지도교수 이 주 용 &gt;

연세대학교 대학원 의학과

문 영 준

서론: 경피적 신절석술은 여전히 크고 복잡한 신장 결석에 대한 최적의 치료법이다. 내시경-결합 신장 내 결석 수술은 경피적 신절석술의 접근로 수를 최소화하는 동시에 무결석율을 향상시키는 것을 목표로 개발되었다. 본 연구의 목적은 한 기관에서 연속된 200례의 내시경-결합 신장 내 결석 수술에 대한 경험을 공유하고, 미니 내시경-결합 신장 내 결석 수술과 표준 내시경-결합 신장 내 결석 수술의 수술 결과를 분석하는 것이다.

방법: 2017년 7월부터 2020년 1월 사이에 성인 환자 200명을 대상으로 내시경-결합 신장 내 결석 수술을 시행하였다. 환자를 전신 마취 하에 중간 누운 자세로 내시경-결합 신장 내 결석 수술을 시행하였다. 튜브리스 기술을 사용하였고 요관 스텐트 삽입을 함께 시행하였다.

결과: 결석 특성 중 평균 최대 결석 길이, 결석 밀도 변동 계수, 선형 결석 밀도, 서울대학교 신장 결석 복잡성 점수 및 수정된 서울대학교 신장 결석 복잡성 점수와 수술 전후 결과 중 추정 혈액 손실, 수술 시간에서 두 군 간에 유의한 차이가 있었다. 성향-점수 매칭 결과에서는 두 군 간에 추정 혈액 손실에서만 유의한 차이가 있었다. 로지스틱 회귀 분석 모델에서, 평균 최대 결석 길이 [승산비 0.953; 95% 신뢰 구간 0.926-0.979;  $P<0.001$ ]와 선형 결석 밀도 (승산비 4.702; 95% 신뢰 구간 1.613-18.655;  $P=0.013$ )는 내시경-결합 신장 내 결석 수술 성공률에 있어 유의한 인자였다.

결론: 미니 내시경-결합 신장 내 결석 수술을 시행한 환자의 경우 결석의

크기가 상대적으로 작고 덜 복잡하며 수술 시간도 짧았다. 그러나 결석의 크기가 비슷할 경우 성공률에는 차이가 없었으나, 표준 내시경-결합 신장 내 결석 수술에 비해 미니 내시경-결합 신장 내 결석 수술에서 추정 혈액 손실은 더 낮을 수 있다.

---

핵심되는 말: 신장; 신장 결석; 쇄석술; 요로 결석



## PUBLICATION LIST

*Medicina* 2023, 59(11), 1971; <https://doi.org/10.3390/medicina59111971>