

# Improved Coronal Alignment Using the Preemptive Joint Line Convergence Angle Compensation Method in Medial Open Wedge High Tibial Osteotomy

## A Retrospective Propensity Score-Matched Analysis

Junwoo Byun,<sup>†‡</sup> MD , Min Jung,<sup>†§</sup> MD, PhD, Kwangho Chung,<sup>†||</sup> MD, Hyun-Soo Moon,<sup>†¶</sup> MD, PhD , Se-Han Jung,<sup>†§</sup> MD , Kyoung-Tak Kang,<sup>#\*\*††</sup> PhD , and Sung-Hwan Kim,<sup>\*†§</sup> MD, PhD 

*Investigation performed at the Department of Orthopedic Surgery, Yonsei University College of Medicine, Seoul, Republic of Korea*

**Background:** Accurate coronal alignment is crucial in medial open wedge high tibial osteotomy (MOWHTO). Postoperative changes in the joint line convergence angle (JLCA) can cause coronal alignment errors. A preemptive JLCA compensation method has been proposed to address this; however, its clinical value remains unclear.

**Purpose:** To assess the accuracy of the preemptive JLCA compensation method for correction angle calculation in MOWHTO.

**Study Design:** Cohort study; Level of evidence, 3.

**Methods:** Patients who underwent MOWHTO for varus osteoarthritis alignment between 2010 and 2024 were reviewed. The patients were classified into 2 groups based on whether the preemptive JLCA compensation method was applied (group 1: the conventional Miniaci method, group 2: the preemptive JLCA compensation method). After propensity score matching for sex, body mass index, hip-knee-ankle (HKA) angle, Kellgren-Lawrence grade, and correction angle on standing radiography, we conducted a comparative analysis of radiologic outcomes (weightbearing ratio) and functional outcomes (Lysholm score and International Knee Documentation Committee [IKDC] scores). Acceptable alignment was defined as a postoperative weightbearing line ratio between 55% and 70%.

**Results:** After propensity score matching, 20 patients were included in each group. The mean preoperative HKA angle was varus  $6.2^\circ \pm 2.2^\circ$  and  $6.3^\circ \pm 2.2^\circ$  ( $P = .922$ ), while the correction angle measured by the Miniaci method was  $10.0^\circ \pm 2.1^\circ$  and  $10.3^\circ \pm 1.7^\circ$  for the conventional and preemptive JLCA compensation method group, respectively ( $P = .855$ ). The mean postoperative weightbearing line ratio at 1 year postoperatively was  $64.1 \pm 7.4$  and  $62.5 \pm 4.7$  for the conventional and preemptive JLCA compensation methods, respectively, with no significant difference. However, 95.0% of patients in the JLCA modification group achieved acceptable target alignment, compared with 60.0% (12/20) in the conventional group ( $P = .014$ ). Despite the significant difference in coronal alignment accuracy, functional outcomes showed no significant difference between the 2 groups.

**Conclusion:** The preemptive JLCA compensation method significantly improved correction accuracy, with 95.0% of patients achieving acceptable target alignment, compared with 60.0% with the conventional Miniaci method, although no significant differences in clinical scores were found in the short-term follow-up period.

**Keywords:** correction angle; high tibial osteotomy; joint line convergence angle; osteoarthritis

a varus-malaligned knee in a mild valgus position, thereby reducing the load on the medial compartment and alleviating symptoms.<sup>6,14,18</sup> One key factor for successful outcomes is achieving appropriate postoperative alignment. Target coronal alignment is generally planned such that the weightbearing line passes through the Fujisawa point, which corresponds to 62.5% of the tibial plateau width from the medial edge.<sup>1,12,16</sup> Among various methods for calculating the correction angle, the Miniaci method is commonly used to determine the correction angle required to achieve the target alignment.<sup>2,21</sup>

However, even when the Miniaci method targeting the Fujisawa point is performed with precise radiographic calculations, not all patients achieve appropriate postoperative alignment. This may result from the fact that preoperative measurements consider only bony factors and neglect the condition of the soft tissues. Notably, after realignment of the lower extremity, altered biomechanics changes the force on the surrounding soft tissues of the knee, leading to unplanned, individual variations in the joint line convergence angle (JLCA, the angle between the 2 lines tangential to the femoral condyles and tibial plateau) depending on tissue tightness or laxity. Several studies have attempted to estimate the degree of change in JLCA postoperatively.<sup>9,19,20</sup> Recently, Jung et al<sup>5</sup> proposed a formula to predict postoperative JLCA change using preoperative JLCA, JLCA under valgus stress test, and preoperative joint line obliquity, which demonstrated superior predictive accuracy compared with the previous reports (Estimated JLCA change =  $0.493 \times (\text{JLCA under valgus stress}) - 0.727 \text{ Standing JLCA} + 0.189 \times (\text{Joint line obliquity}) - 1.587$ ).<sup>4</sup> Furthermore, a method for utilizing the estimated JLCA change for the calculation of the correction angle was proposed (preemptive JLCA compensation method). However, previous studies have only simulated the method using postoperative radiographs, and the actual correction-angle results calculated by the proposed preemptive JLCA compensation method have not been reported. Because the JLCA change prediction was based on a correction angle derived from the conventional Miniaci method rather than from the preemptive JLCA compensation method, validation of this new method is necessary. By evaluating the results of the preemptive JLCA compensation method, our study provides clinical evidence supporting a more accurate approach to correcting angle

calculations in MOWHTO. We hypothesized that this method would yield a superior clinical outcome.

## METHODS

This study was approved by the institutional review board, and the requirement for informed consent was waived because of its retrospective nature. The medical records of patients who underwent MOWHTO for advanced osteoarthritis, typically defined as Kellgren-Lawrence (KL) grades 3 and 4, with varus alignment from 2010 to 2024, were retrospectively reviewed. Patients who underwent MOWHTO using correction angle measurement with the Miniaci method targeting the Fujisawa point with or without preemptive JLCA compensation were included. The exclusion criteria were (1) follow-up period of <1 year, (2) hinge fracture confirmed on postoperative radiography, and (3) insufficient outcome data. The preemptive JLCA compensation method was initiated in April 2023, after which all eligible patients were treated using this technique. To apply the preemptive JLCA compensation method, the correction angles were measured on the supine radiograph, weightbearing radiograph with the conventional method, and weightbearing radiograph with the JLCA modification angle. When the preemptive JLCA compensation method yielded a smaller correction angle than that on the supine radiograph, it was considered an outlier and not applied. This was because the supine radiograph reflects lower-extremity alignment without lateral joint opening, a known cause of postoperative JLCA change; thus, the preemptive JLCA compensation method was not utilized. Eligible patients were classified based on whether the preemptive JLCA compensation method was used. Propensity score matching was performed on sex, body mass index (BMI), HKA angle, KL grade, and correction angle on standing radiographs between the 2 methods. A positive value of the HKA angle indicated varus alignment, and the KL grade was defined as a higher grade on full-extension weightbearing anteroposterior (AP) and 45° flexion posteroanterior (PA) views. The final sample included 2 groups based on the correction angle measurement method (group 1, conventional method; group 2, preemptive JLCA compensation method) (Figure 1).

\*Address correspondence to Sung-Hwan Kim, MD, PhD, Department of Orthopedic Surgery, Yonsei University College of Medicine, Severance Hospital, 50-1 Yonsei-ro, Seodaemun-gu, Seoul 03722, Republic of Korea (emails: orthohwan@gmail.com; orthohwan@yuhs.ac).

<sup>†</sup>Arthroscopy and Joint Research Institute, Yonsei University College of Medicine, Seoul, Republic of Korea.

<sup>‡</sup>Department of Orthopaedic Surgery, Mokdong Hospital, Ewha Womans University College of Medicine, Seoul, Republic of Korea.

<sup>§</sup>Department of Orthopedic Surgery, Severance Hospital, Yonsei University College of Medicine, Seoul, Republic of Korea.

<sup>||</sup>Department of Orthopedic Surgery, Yonjin Severance Hospital, Yonsei University College of Medicine, Yonjin, Republic of Korea.

<sup>¶</sup>Department of Orthopedic Surgery, Gangnam Severance Hospital, Yonsei University College of Medicine, Seoul, Republic of Korea.

<sup>#</sup>Clevion Co, LTD, Seoul, Republic of Korea.


<sup>\*\*</sup>Skyve R&D LAB, Seoul, Republic of Korea.

<sup>††</sup>Department of Mechanical Engineering, Yonsei University, Seoul, Republic of Korea.

Final version submitted November 26, 2025; accepted December 12, 2025.

One or more of the authors has declared the following potential conflict of interest or source of funding: Kyoung-Tak Kang is the Chief Executive Officer of Skyve and Clevion Co, Ltd. AOSSM checks author disclosures against the Open Payments Database (OPD). AOSSM has not conducted an independent investigation on the OPD and disclaims any liability or responsibility relating thereto.

Ethical approval for this study was obtained from Severance Hospital (IRB No. 4-2025-0285).



**Figure 1.** Flowchart for patient inclusion. JLCA, joint line convergence angle.

### Surgical Indication

At our institution, the indications for MOWHTO were as follows: (1) age <70 years; (2) advanced osteoarthritis defined as KL grade 3 or 4 on the medial compartment of the affected knee; (3) hip–knee–ankle (HKA) >5°, with positive values indicating varus alignment; (4) absence of advanced osteoarthritis in the lateral and patellofemoral compartment; and (5) no major limitation of range of motion, defined as flexion contracture >15° or active flexion angle <100°.

### Correction Angle Measurement by Preemptive JLCA Compensation Method


Before applying the preemptive JLCA compensation method, the estimated JLCA change was calculated using a previously reported formula consisting of JLCA on the 0° knee flexion valgus stress view AP radiograph obtained by applying 150 N using a Telos stress device (vgJLCA<sup>0</sup>), JLCA on a standing AP radiograph (preJLCA<sup>std</sup>), and pre-operative joint line obliquity (preJLO) (Figure 2). Both JLCA and preJLO were defined as positive values when the lateral apex or inclination was directed laterally.

The estimated JLCA change was calculated by applying the variables to the formula reported by the previous study (estimated  $\Delta$ JLCA =  $0.493 \times (\text{vgJLCA}^0) - 0.727 \times (\text{preJLCA}^{\text{std}}) + 0.189 \times (\text{preJLO}) - 1.587$ ).<sup>4</sup> After calculation,


the tibia and the fibula were rotated on the weightbearing whole lower extremity radiography using ImageJ software (National Institute of Health Image). Specifically, the tibia and fibula were rotated by the estimated JLCA change, using the midpoint of the tibial spine as the rotation center. The correction angle was measured on the rotated image using the Miniaci method. When the correction angle measured on the rotated image was between the correction angles on supine and standing radiographs, a 3-dimensional patient-specific instrument (3D PSI) gapper (Kneevigate, Skyve) was created after simulating the correction using Mimics software (Version 17; Materialize) for segmentation, and OnKnee-U software (Skyve) for simulation (Figure 3).

### Surgical Procedure and Rehabilitation

A single senior orthopaedic surgeon performed all procedures (S.-H.K.). Before osteotomy, diagnostic arthroscopy and concomitant procedures for cartilage and meniscal lesions were performed. After arthroscopy, an osteotomy was performed. A longitudinal incision was made midway between the anterior and medial borders of the tibia, and the pes anserinus was partially detached to expose the osteotomy site. The osteotomy level was set 4 cm below the joint line from the starting point, confirmed on fluoroscopy, with the hinge point located in the upper one-third of the proximal tibiofibular joint, as described in



**Figure 2.** Measurement of the (A) preoperative JLCA under valgus stress radiograph, and (B) preoperative standing JLCA (angle between the yellow lines) and JLO (angle between the blue lines, with the lower blue line representing the line parallel to the ground). JLCA, joint line convergence angle; JLO, joint line obliquity.



**Figure 3.** Correction angle measurement using the preemptive JLCA compensation method. Before the measurement, the estimated JLCA is calculated. (A) The midpoint of the tibial spine (tip of the yellow arrowhead), the Fujisawa point (62.5% of the tibia plateau, red asterisk), the target hinge area (tip of the blue arrow), and the ankle center (green circle) are marked. The area including the whole tibia and fibula is marked (red line). (B) The marked area is rotated to the degree of estimated JLCA change ( $0.493 \times \text{vgJLCA}^0 - 0.727 \times \text{preJLCA}^{\text{std}} + 0.189 \times \text{preJLO} - 1.587$ ) in accordance with the previously reported formula toward the direction of valgization, with the center of rotation being the midpoint of the tibial spine. (C) After the rotation, the correction angle is measured using the Miniaci method. (D) MOWHTO to the degree of the planned correction angle is simulated, and the 3D PSI gapper is printed according to the simulated data. Estimated  $\Delta\text{JLCA}$ , estimated change of joint line convergence angle; JLCA, joint line convergence angle; MOWHTO, medial open wedge high tibial osteotomy;  $\text{preJLCA}^{\text{std}}$ , preoperative joint line convergence angle on standing radiograph;  $\text{preJLO}$ , preoperative joint line obliquity;  $\text{vgJLCA}^0$ , joint line convergence angle on  $0^\circ$  flexion valgus stress radiograph.

the previous study.<sup>4</sup> Gap distraction was performed using the adjustable bone spreader. For the preemptive JLCA compensation method, a 3D PSI gapper was used to

maintain the gap during plate fixation.<sup>5</sup> The detached pes anserinus was reattached by embedding it under the locking plate.

## Patient Evaluation

Descriptive data, subjective functional scores, and radiological data were retrospectively analyzed. The descriptive data included age, sex, height, weight, and BMI.

Radiological variables included both pre- and postoperative variables. The HKA angle, weightbearing line ratio, KL grade of the medial compartment on full extension weightbearing AP and 45° flexion PA views at each follow-up period, and correction angle measured on weightbearing whole lower extremity radiography by the Miniaci method were included. For patients in group 2, additional measurements of the correction angle on supine whole-lower-extremity radiographs were performed. Two experienced physicians (J.B. and S.-H.J.), blinded to each other's results and group allocation, independently measured the radiological variables. The intraclass correlation coefficient (ICC) was calculated to assess interobserver reliability.

Subjective functional outcomes were assessed, including the visual analog scale (VAS) for Pain, Lysholm score, International Knee Documentation Committee (IKDC) subjective score, Knee injury and Osteoarthritis Outcome Score (KOOS), and Western Ontario and McMaster Universities (WOMAC). These scores were prospectively reported preoperatively, at 6 months postoperatively, and annually thereafter. Thresholds of minimal clinically important difference (MCID) for the VAS, IKDC subjective, Lysholm, KOOS symptoms, KOOS Pain, KOOS activities of daily Living (ADL), KOOS Sports/Recreation, KOOS Quality of Life (QoL), and WOMAC scores were derived from a distribution-based method, applying 0.5 standard deviation to define meaningful clinical improvement (14.8, 7.8, 10.2, 10.2, 9.7, 8.6, 13.6, 9.7, and 8.0, respectively).

## Ethical Approval

This study was approved by the institutional review board (IRB) of Severance Hospital, and the requirement for informed consent was waived owing to its retrospective nature (IRB No. 4-2025-0285).

## Statistical Analysis

An a priori power analysis was performed to determine the minimum sample size for each group. Based on the reported proportion of overcorrection patients from another method to reduce overcorrection,<sup>9</sup> and comparing with a previous study analyzing the change of JLCA, a minimal number of patients for adequate power was 19 patients for each group for the significance level (alpha) and power (1 – beta) set as .05 and 0.8, respectively.<sup>4</sup> For the propensity score matching, the MatchIt package in R Version 4.4.1 (R Core Team, R Foundation for Statistical Computing) was used. All statistical analyses, except propensity score matching, were conducted using SPSS Version 26 (IBM Inc). For continuous variables, normality was confirmed using the Shapiro-Wilk test. The student *t* test for normally distributed data and the Mann-Whitney *U* test for non-normally distributed data were used for

TABLE 1  
Comparison of Baseline Demographic Data  
and Intraoperative Data<sup>a</sup>

Variables	Group 1, n = 20	Group 2, n = 20	<i>P</i>
Descriptive data			
Age, years	54.1 ± 5.5	56.1 ± 5.6	.263
Sex, male/female	5/15	7/13	.731
Height, cm	159.7 ± 7.5	159.2 ± 8.7	.826
Weight, kg	71.8 ± 13.5	73.2 ± 14.3	.989
BMI, kg/m <sup>2</sup>	28.0 ± 4.0	28.7 ± 3.4	.341

<sup>a</sup>Values are presented as mean ± SD or number of patients. Group 1, the conventional Miniaci method; Group 2, the preemptive JLCA compensation method. BMI, body mass index.

between-group comparisons. For categorical variables, the Pearson chi-square test or the Fisher exact test was used. The Pearson correlation test was used to analyze the correlation. Statistical significance was set at *P* < .05.

## RESULTS

A total of 117 patients were included in this study after excluding 2 patients, resulting in a lower correction angle derived from the JLCA modification than from the correction angle on the supine radiograph. Among the included patients, 97 underwent correction using the conventional method (group 1), and 20 using the preemptive JLCA compensation method (group 2) (Figure 1). No significant differences were found in the descriptive data (Table 1). At 1-year postoperatively, 95.0% of group 2 patients achieved acceptable alignment, which was significantly higher than that in group 1 (60.0%), while other pre- and postoperative radiographic variables did not differ significantly between the groups (Table 2). The reliability of the radiological measurements for continuous variables was good to excellent, with ICC values<sup>10</sup> ranging from 0.883 to 0.936. For categorical variables, reliability was assessed<sup>11</sup> as having substantial agreement, with kappa values of 0.684 and 0.730.

Subjective functional outcomes were also not significantly different between the 2 groups. Furthermore, the proportion of patients who achieved improvement beyond the MCID did not differ significantly between the 2 groups (Table 3).

In group 2, the correlation between the estimated JLCA change and the actual change from the preoperative period to 1 year postoperatively was analyzed. The mean estimated JLCA change was 1.0° ± 0.9°, whereas the JLCA change at 1 year postoperatively was 1.1° ± 1.5°. No significant correlation was found between the estimated changes and the changes at 1 year postoperatively (*r* = 0.337; *P* = .146).

## DISCUSSION

The main finding of this study was that the preemptive JLCA compensation method achieved superior accuracy,

TABLE 2  
Comparison of Radiologic Outcomes<sup>a</sup>

Radiologic Variables	Group 1, n = 20	Group 2, n = 20	P
Preoperative			
HKA angle, deg	6.2 ± 2.2	6.3 ± 2.2	.922
Kellgren-Lawrence grade, 3/4	18/2	19/1	>.999
Weightbearing ratio, %	20.9 ± 8.7	20.6 ± 8.4	.738
MPTA, deg	85.6 ± 1.8	84.7 ± 1.9	.103
LDFA, deg	88.3 ± 1.7	88.0 ± 1.9	.588
JLCA, deg	3.8 ± 1.4	3.3 ± 1.6	.299
Correction angle, standing radiograph	10.0 ± 2.1	10.1 ± 2.1	.855
Correction angle, supine radiograph <sup>b</sup>		8.5 ± 1.7	
Correction angle, preemptive JLCA compensation method <sup>b</sup>		9.0 ± 1.7	
Postoperative 1-year			
HKA angle, deg	-3.1 ± 1.7	-3.6 ± 1.1	.344
JLCA, deg	2.8 ± 1.6	2.2 ± 1.6	.266
Weightbearing ratio, %	64.1 ± 7.4	62.4 ± 4.8	.394
Undercorrection	3	1	.014
Acceptable	12	19	
Overcorrection	5	0	

<sup>a</sup>Data are presented as mean ± SD or number of patients, unless otherwise indicated. Group 1, the conventional Miniaci method; Group 2, the preemptive JLCA compensation method.

HKA, hip-knee-ankle; JLCA; joint line convergence angle; LDFA, lateral distal femoral angle; MPTA, medial proximal tibial angle.

<sup>b</sup>The values are only measured for group 2.

with 95.0% of patients attaining acceptable postoperative alignment, compared with 60.0% with the conventional method.

One of the most crucial goals of MOWHTO is to achieve the target coronal alignment postoperatively, as previous studies have reported adverse outcomes under or overcorrection.<sup>7,12,15</sup> However, not all patients undergoing MOWHTO achieve acceptable alignment, and therefore several previous studies have reported methods for more accurate correction.<sup>9,20</sup> Among them, Shin et al<sup>20</sup> reported that, for the conventional method of correction angle measurement, using supine radiographs rather than standing radiographs resulted in better accuracy.<sup>20</sup> This could be attributed to changes in the weightbearing line after surgery; postoperative varus alignment causes medial dominant loading and increased JLCA during weightbearing, whereas lateral shift of the loading axis postoperatively reduces this effect.<sup>17</sup> Both the supine radiography method and the preemptive JLCA compensation method aim to adjust the correction angle by taking the effect of soft tissue into account. However, the preemptive JLCA compensation method has its strength in that it could reflect the effect of weightbearing adequately, whereas supine radiography neglects it as a whole. Nevertheless, the correction angle from supine radiography remains important for the preemptive JLCA compensation method. Considering that the supine radiography method entirely excludes the influence of soft tissue during weightbearing, any patient who had a lower degree of correction angle measured by the preemptive JLCA compensation method compared with supine radiography was regarded as an outlier in predicting JLCA changes; thus, the preemptive JLCA

compensation method was not applied in such cases. Further studies are required to achieve a more accurate realignment for patients with a smaller correction angle using the preemptive JLCA compensation method than using the supine radiography method.

A previous study that reported the estimated JLCA change used in our method retrospectively simulated postoperative radiographs based on the calculated JLCA change.<sup>4</sup> However, such retrospective simulations have limited clinical applicability because they cannot account for JLCA changes associated with reduced correction angle. Notably, the actual JLCA change and estimated value did not have a significant correlation when applying the correction angle of the preemptive JLCA compensation method. The mean difference of 1.1° between the correction angle from standing radiography and the preemptive JLCA compensation method could be a cause of this insignificance. Despite its insignificance, the proposed preemptive JLCA compensation method still has its strength for accurate correction, resulting in an acceptable range in 95.0% of patients. Therefore, the clinical implication of the current preemptive JLCA compensation method is to accurately correct coronal alignment rather than estimate the change in JLCA.

Another factor influencing correction accuracy is the use of 3D PSIs. Because the JLCA medication method adjusts the correction angle in 0.1° increments, 3D PSI plays a crucial part in the process. Although the superior outcome for group 2 patients in achieving target alignment could stem from the use of 3D PSI, 3D PSI itself did not result in superior postoperative coronal alignment. The proportion of patients achieving acceptable postoperative

TABLE 3  
Comparison of Clinical Scores<sup>a</sup>

Clinical scores	Group 1, n = 20	Group 2, n = 20	P
VAS score			
Preop	63.7 ± 18.4	67.3 ± 21.4	.565
6 months postop	35.4 ± 24.0	27.8 ± 18.7	.271
MCID achievement, yes/ no <sup>b,c</sup>	19/ 1	17/ 3	.605
Postop 1 year	23.8 ± 18.4	24.3 ± 17.9	.938
MCID achievement, yes/ no <sup>b,c</sup>	14/ 6	16/ 4	.716
IKDC subjective score			
Preop	37.4 ± 13.2	34.9 ± 14.8	.567
6 months postop	46.3 ± 13.3	45.9 ± 14.1	.923
MCID achievement, yes/ no <sup>b,c</sup>	15/ 5	13/ 7	.731
Postop 1 year	54.4 ± 15.6	54.6 ± 18.4	.779
MCID achievement, yes/ no <sup>b,c</sup>	11/9	12/8	>.999
Lysholm score			
Preop	37.1 ± 19.7	47.8 ± 17.0	.075
6 months postop	50.1 ± 18.7	58.0 ± 22.4	.263
MCID achievement, yes/ no <sup>b,c</sup>	16/ 4	12/ 8	.301
Postop 1 year	63.0 ± 21.5	64.1 ± 17.5	.863
MCID achievement, yes/ no <sup>b,c</sup>	11/ 9	9/ 11	.752
KOOS Symptom			
Preop	47.0 ± 22.7	48.0 ± 17.2	.866
6 months postop	63.6 ± 19.9	60.7 ± 19.1	.645
MCID achievement, yes/ no <sup>b,c</sup>	14/ 6	15/ 5	>.999
Postop 1 year	67.4 ± 18.2	67.1 ± 17.6	.957
MCID achievement, yes/ no <sup>b,c</sup>	14/ 6	11/ 9	.514
KOOS pain			
Preop	43.9 ± 22.1	51.2 ± 16.5	.245
6 months postop	64.7 ± 19.8	66.9 ± 17.6	.711
MCID achievement, yes/ no <sup>b,c</sup>	17/ 3	14/ 6	.451
Postop 1 year	70.8 ± 19.6	68.9 ± 17.7	0.744
MCID achievement, yes/ no <sup>b,c</sup>	14/ 6	12/ 8	.741
KOOS ADL			
Preop	49.4 ± 21.6	55.0 ± 17.7	.378
6 months postop	66.8 ± 18.4	70.4 ± 19.4	.560
MCID achievement, yes/ no <sup>b,c</sup>	19/ 1	14/ 6	.091
1 year postop	76.1 ± 15.0	74.2 ± 15.9	.697
MCID achievement, yes/ no <sup>b,c</sup>	16/ 4	13/ 7	.480
KOOS Sport/Rec			
Preop	27.0 ± 26.9	18.3 ± 22.4	.211
6 months postop	32.5 ± 23.6	34.8 ± 29.0	.789
MCID achievement, yes/ no <sup>b,c</sup>	11/ 9	9/ 11	.752
1 year postop	34.5 ± 26.1	36.5 ± 33.2	.883
MCID achievement, yes/ no <sup>b,c</sup>	8/ 12	9/ 11	>.999
KOOS QoL			
Preop	22.8 ± 15.8	26.9 ± 16.8	.434
6 months postop	40.0 ± 18.5	37.2 ± 21.9	.662
MCID achievement, yes/ no <sup>b,c</sup>	15/ 5	12/ 8	.501
1 year postop	47.2 ± 24.4	46.2 ± 25.4	.907
MCID achievement, yes/ no <sup>b,c</sup>	12/ 8	11/ 9	>.999
WOMAC			
Preop	49.5 ± 20.4	43.0 ± 16.0	.269
6 months postop	32.4 ± 17.6	29.4 ± 17.8	.602
MCID achievement, yes/ no <sup>b,c</sup>	20/0	18/2	.487
1 year postop	23.3 ± 14.1	25.6 ± 15.3	.617
MCID achievement, yes/ no <sup>b,c</sup>	19/1	18/2	>.999

<sup>a</sup>Data are presented as mean ± SD, unless otherwise indicated. Group 1, the conventional Miniaci method; Group 2, the preemptive JLCA compensation method. ADL, activities of daily living; IKDC, International Knee Documentation Committee; KOOS, Knee injury and Osteoarthritis Outcome Score; MCID, minimal clinically important differences; Postop, postoperative; Preop, preoperative; QoL, quality of life; Sport/Rec, sports/recreation; VAS, visual analog scale; WOMAC, Western Ontario and McMaster Universities.

<sup>b</sup>Values are presented as the number of patients.

<sup>c</sup>Comparison with the corresponding preoperative variables.

alignment did not significantly differ between group 1, which included patients with and without the application of 3D PSI, and the patients enrolled in the previous study, which only included patients with 3D PSI, showed no significant differences (Appendix table A3). Considering these results, the superior outcome of the preemptive JLCA compensation method is more likely to stem from the method itself rather than the utilization of the 3D PSI.

## Limitations

This study had some limitations. First, its retrospective nature introduced a potential risk of selection bias owing to the study design. Second, although the proposed method demonstrated high accuracy, a few patients showed a lower correction angle with the preemptive JLCA compensation method than with supine radiography. Thus, the preemptive JLCA compensation method was not utilized. Further studies are needed to identify alternative treatment methods for these patients. Third, although the discrepancy and nonsignificant correlation between the estimated and actual JLCA change could be attributed to the degree of modification after applying the preemptive JLCA compensation method, the potential clinical implications warrant further studies. Fourth, this study included only short-term follow-up, and further research should assess long-term outcomes. Fifth, although the number of patients per group exceeded the minimum required by the a priori power analysis, the overall sample size remained relatively small.

## CONCLUSION

The preemptive JLCA compensation method significantly improved correction accuracy, with 95.0% of patients achieving acceptable target alignment, compared with 60.0% with the conventional Miniaci method. However, no significant differences in clinical scores were found in the short-term follow-up period.

## ORCID iDs

Junwoo Byun  <https://orcid.org/0000-0002-8372-5247>  
 Hyun-Soo Moon  <https://orcid.org/0000-0001-8678-2960>  
 Se-Han Jung  <https://orcid.org/0000-0001-8422-093X>  
 Kyoung-Tak Kang  <https://orcid.org/0000-0002-6752-4576>  
 Sung-Hwan Kim  <https://orcid.org/0000-0001-5743-6241>

## REFERENCES

- Alemayehu DG, Zhang Z, Tahir E, et al. Undercorrection: the undesired effect of compression on the osteotomy gap of the medial opening wedge high tibial osteotomy and its clinical significance. *Arch Orthop Trauma Surg.* 2022;142(6):937-946.
- Aygün Ü, Bölükbaşı M, Yamak K, Çiçek AC. Comparison of the Miniaci and Dugdale techniques on functional outcomes in medial opening wedge high tibial osteotomy. *J Exp Orthop.* 2023;10(1):86.
- Han JH, Jung M, Chung K, et al. Effects of concurrent cartilage procedures on cartilage regeneration in high tibial osteotomy: a systematic review. *Knee Surg Relat Res.* 2024;36(1):13.
- Jung SH, Jung M, Chung K, et al. Preoperative joint line obliquity, a newly identified factor for overcorrection, can be incorporated into a novel preoperative planning method to optimise alignment in high tibial osteotomy. *Knee Surg Sports Traumatol Arthrosc.* 2024;32(1):64-77.
- Jung SH, Jung M, Chung K, et al. Factors causing unintended sagittal and axial alignment changes in high tibial osteotomy: comparative 3-dimensional analysis of simulation and actual surgery. *Am J Sports Med.* 2024;52(6):1543-1553.
- Jung SH, Park H, Jung M, et al. Implantation of hUCB-MSCs generates greater hyaline-type cartilage than microdrilling combined with high tibial osteotomy. *Knee Surg Sports Traumatol Arthrosc.* 2024;32(4):829-842.
- Kang BY, Lee DK, Kim HS, Wang JH. How to achieve an optimal alignment in medial opening wedge high tibial osteotomy? *Knee Surg Relat Res.* 2022;34(1):3.
- Kim HJ, Shin JY, Lee HJ, et al. Does preoperative bone marrow edema affect clinical outcomes after medial open-wedge high tibial osteotomy? *Knee Surg Relat Res.* 2024;36(1):9.
- Kim MS, Koh IJ, Sung YG, et al. Alignment adjustment using the Valgus stress technique can increase the surgical accuracy of novice surgeons during medial opening-wedge high Tibial osteotomy. *BMC Musculoskelet Disord.* 2021;22(1):585.
- Koo TK, Li MY. A Guideline of selecting and reporting intraclass correlation coefficients for reliability research. *J Chiropr Med.* 2016;15(2):155-163.
- Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics.* 1977;33(1):159-174.
- Lee SS, Kim JH, Kim S, et al. Avoiding overcorrection to increase patient satisfaction after open wedge high tibial osteotomy. *Am J Sports Med.* 2022;50(9):2453-2461.
- Mabrouk A, Risebury M, Yasen S. High survivorship and low complication rate in a single-centre series of 651 medial opening wedge high tibial osteotomy cases with a mean follow-up of 13 years. *Knee Surg Sports Traumatol Arthrosc.* 2024;32(3):736-749.
- Mabrouk A, Risebury M, Yasen S. Medial opening wedge high tibial osteotomy performs similarly irrespective of body mass index. *Knee Surg Sports Traumatol Arthrosc.* 2024;32(9):2328-2341.
- Machado A, Micicoi L, Ernat J, et al. Normo- or slightly overcorrection show better results after medial closing wedge high tibial osteotomy. *Knee Surg Sports Traumatol Arthrosc.* 2023;31(10):4276-4284.
- Murray R, Winkler PW, Shaikh HS, Musahl V. High tibial osteotomy for varus deformity of the knee. *J Am Acad Orthop Surg Glob Res Rev.* 2021;5(7).
- Noyes FR. Editorial commentary: measurements for successful high tibial osteotomy: understanding supine versus standing and intraoperative fluoroscopic alignment is required. *Arthroscopy.* 2020;36(6):1665-1669.
- Park JY, Kim JH, Cho JW, Kim MS, Choi W. Clinical and radiological results of high tibial of osteotomy over the age of 65 are comparable to that of under 55 at minimum 2-year follow-up: a propensity score matched analysis. *Knee Surg Relat Res.* 2024;36(1):10.
- Ryu DJ, Lee SS, Jung EY, et al. Reliability of preoperative planning method that considers latent medial joint laxity in medial open-wedge proximal tibial osteotomy. *Orthop J Sports Med.* 2021;9(10):23259671211034151.
- Shin KH, Jung JK, Nam JJ, Jang KM, Han SB. Preoperative supine radiographs are more accurate than standing radiographs for preoperative planning in medial open-wedge high tibial osteotomy. *Arthroscopy.* 2020;36(6):1655-1664.
- Sivertsen EA, Vik J, Meland ASV, Nerhus TK. The Dugdale planning method for high tibial osteotomies underestimates the correction angle compared to the Miniaci method. *Knee Surg Sports Traumatol Arthrosc.* 2023;31(4):1507-1514.

## APPENDIX

TABLE A1  
Variables Included in the Propensity Score Matching<sup>a</sup>

Variables	Before Matching		After Matching		SMD <sup>b</sup>
	Group 1	Group 2	Group 1	Group 2	
Sex, male/female <sup>c</sup>	53/ 44	7/ 13	5/ 15	7/ 13	.20
BMI, kg/m <sup>2</sup>	26.8 ± 3.4	28.7 ± 3.4	28.0 ± 4.0	28.7 ± 3.4	.20
HKA angle, deg	7.6 ± 2.8	6.3 ± 2.2	6.2 ± 2.2	6.3 ± 2.2	.03
Kellgren-Lawrence grade, 3/4 <sup>c</sup>	76/21	19/1	18/2	19/ 1	.16
Correction angle, standing radiograph, deg	11.3 ± 2.7	10.1 ± 2.1	10.0 ± 2.1	10.1 ± 2.1	.06

<sup>a</sup>Data are presented as mean ± SD, unless otherwise indicated. Group 1, the conventional Miniaci method; Group 2, the preemptive JLCA compensation method. BMI, body mass index; HKA, hip-knee-ankle; SMD, standardized mean difference.

<sup>b</sup>Values are presented with 2 decimal places.

<sup>c</sup>Values are presented as the number of patients.

TABLE A2  
Results of the normality test on Continuous Variables

Variables	P <sup>a</sup>	Variables	P <sup>a</sup>
Age, years	.286	6 months postop	
Height, cm	.618	VAS score	.111
Weight, kg	.009	IKDC subjective score	.957
BMI, kg/m <sup>2</sup>	.038	Lysholm score	.604
Preop		KOOS Symptom	.883
HKA angle, deg	.112	KOOS Pain	.796
Weightbearing ratio, %	.038	KOOS ADL	.591
MPTA, deg	.547	KOOS Sport/Rec	.092
LDFA, deg	.059	KOOS QoL	.692
JLCA, deg	.122	WOMAC	.630
Correction angle, standing radiograph	.145	Postoperative 1 year	
Correction angle, supine radiograph	.464	VAS score	.067
Correction angle, JLCA modification	.993	IKDC subjective score	.045
1 year postop		Lysholm score	.846
HKA angle, deg	.377	KOOS Symptom	.556
JLCA, deg	.111	KOOS Pain	.353
Weightbearing ratio, %	.305	KOOS ADL	.635
Preop		KOOS Sport/Rec	.016
VAS score	.025	KOOS QoL	.065
IKDC subjective score	.946	WOMAC	.817
Lysholm score	.997		
KOOS Symptom	.329		
KOOS Pain	.794		
KOOS ADL	.973		
KOOS Sport/Rec	.001		
KOOS QoL	.396		
WOMAC	.995		

<sup>a</sup>Values are presented as the result of the Shapiro-Wilk test for normality. Group 1, the conventional Miniaci method; Group 2, the preemptive JLCA compensation method. ADL, activities of daily living; BMI, body mass index; HKA, hip-knee-ankle angle; IKDC, International Knee Documentation Committee; JLCA, joint line convergence angle; KOOS, Knee injury and Osteoarthritis Outcome Score; LDFA, lateral distal femoral angle; MPTA, medial proximal tibial angle; Postop, postoperative; Preop, preoperative; QoL, quality of life; Sport/Rec, sports/recreation; VAS, visual analog scale; and WOMAC, Western Ontario and McMaster Universities Osteoarthritis Index.

TABLE A3  
Comparison of Postoperative Coronal Alignment With Regard to the Utilization of 3D PSI<sup>a</sup>

Radiologic Variables <sup>b</sup>	Group 1 <sup>c</sup> n = 20	Previous Study <sup>d</sup> n = 61	<i>P</i>
Postop coronal alignment <sup>e</sup>			.879
Undercorrection	3	7	
Acceptable	10	29	
Overcorrection	7	25	

<sup>a</sup>Group 1, the conventional Miniaci method; Group 2, the preemptive JLCA compensation method. MOWHTO, medial open wedge high tibial osteotomy; Postop, postoperative; PSI, patient-specific instrument; 3D, 3-dimensional.

<sup>b</sup>Values are presented as the number of patients.

<sup>c</sup>Including patients both with and without the application of 3D PSI.

<sup>d</sup>All included patients had undergone MOWHTO using 3D PSI.

<sup>e</sup>Acceptable alignment defined as  $62.5\% \pm 6.25\%$ , to compare with the result of previous study.