

Comparison of double pants with single pants on satisfaction with colonoscopy

Sook Hee Chung, Soo Jung Park, Jong Suk Hong, Jee Young Hwang, Sin Ae Lee, Kyung Ran Kim, Hye Sun Lee, Sung Pil Hong, Jae Hee Cheon, Tae Il Kim, Won Ho Kim

Sook Hee Chung, Soo Jung Park, Jong Suk Hong, Jee Young Hwang, Sin Ae Lee, Sung Pil Hong, Jae Hee Cheon, Tae Il Kim, Won Ho Kim, Department of Internal Medicine and Institute of Gastroenterology, Yonsei University College of Medicine, Seoul 120-752, South Korea

Kyung Ran Kim, Department of Psychiatry, Yonsei University College of Medicine, Seoul 120-752, South Korea

Hye Sun Lee, Department of Biostatistics, Yonsei University College of Medicine, Seoul 120-752, South Korea

Author contributions: Chung SH and Park SJ contributed equally to this work; Chung SH and Park SJ contributed equally to this work; Chung SH and Park SJ analyzed the data and wrote the thesis. Hong JS, Hwang JY and Lee SA collected the data; Kim KR provided consultation about psychological interview; Lee HS provided consultation about statistics; Hong SP, Cheon JH and Kim TI enrolled the patients and reviewed the manuscript; Kim WH designed the research and reviewed the manuscript.

Correspondence to: Won Ho Kim, MD, PhD, Department of Internal Medicine and Institute of Gastroenterology, Yonsei University College of Medicine, 50 Yonseiro, Seodaemun-gu, Seoul 120-752, South Korea. kimwonho@yuhs.ac

Telephone: +82-2-22281951 Fax: +82-2-3936884

Received: February 14, 2013 Revised: April 24, 2013

Accepted: June 5, 2013

Published online: July 14, 2013

Abstract

AIM: To increase satisfaction and diminish anxiety and shame during colonoscopy, we developed novel double pants (NDP) which consist of doubled fabrics with an inner hole. The aim of study was to compare satisfaction, anxiety and shame between NDP and conventional single pants (CSP).

METHODS: Total 160 consecutive examinees were randomly divided into NDP and CSP group. Before colonoscopy, questionnaires identifying state and trait anxiety were completed. After colonoscopy, questionnaires for overall satisfaction (Group Health Association of

America 9) and pants-specific satisfaction (5-20), state anxiety (20-80), and shame (6-24) were interviewed.

RESULTS: Pants-specific satisfaction scores regarding willingness to repeat colonoscopy using same pants (3.3 ± 0.8 vs 2.1 ± 0.9 , $P < 0.001$) and recommendation of same pants to other people (3.3 ± 0.7 vs 2.0 ± 1.0 , $P < 0.001$) were significantly higher in NDP than CSP groups. State anxiety (33.0 ± 7.0 vs 35.4 ± 6.9 , $P = 0.028$) and shame (6.6 ± 1.5 vs 8.1 ± 3.2 , $P = 0.001$) after colonoscopy was lower in NDP group compared with CSP group.

CONCLUSION: The NDP contribute to increase satisfaction and decrease anxiety and shame after colonoscopy.

© 2013 Baishideng. All rights reserved.

Key words: Pants; Colonoscopy; Satisfaction; Shame; Anxiety

Core tip: We developed novel double pants (NDP) those are consisted of double fabrics with an inner hole. We compared the satisfaction, anxiety and shame between NDP and conventional single pants (CSP). The examinees in NDP group responded higher pants specific satisfaction, lower state anxiety after colonoscopy and lower shame score compared to those in CSP group. NDP developed in our institute may contribute to increase satisfaction and decrease anxiety and shame after colonoscopy.

Chung SH, Park SJ, Hong JS, Hwang JY, Lee SA, Kim KR, Lee HS, Hong SP, Cheon JH, Kim TI, Kim WH. Comparison of double pants with single pants on satisfaction with colonoscopy. *World J Gastroenterol* 2013; 19(26): 4177-4184 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v19/i26/4177.htm> DOI: <http://dx.doi.org/10.3748/wjg.v19.i26.4177>

INTRODUCTION

Colonoscopy has recently increased in importance worldwide due to its use in screening for colon polyps and colorectal cancer^[1]. Even in institutions where conscious sedated colonoscopy is available, colonoscopy without sedation is still performed due to patient comorbidities, the economic burden of sedation, and examinee preference. There is considerable global variation in the prevalence of sedative endoscopy^[2]. Most colonoscopies in the United States are performed as sedated procedures, but some countries rarely use sedation in colonoscopy^[3,4].

Consideration of the factors affecting satisfaction during colonoscopy has also recently increased because satisfaction may represent an important quality indicator for colonoscopy^[5,6]. Moreover, the satisfaction of examinees may be reduced by anxiety, shame, discomfort, and embarrassment^[7-11]. Many factors have been identified which determine satisfaction, anxiety, and shame, including the circumstance of the endoscopy room, the clothing for the procedure, endoscopist's skill, and the unfamiliarity of the medical staff with whom examinees interact during the procedure.

During colonoscopy at our center, examinees previously wear conventional single pants (CSP), where there is no hole for insertion of the scope. Examinees remove the CSP to below the level of the buttocks and expose their buttocks area during colonoscopy. Exposing the buttocks during colonoscopy can make examinees feel shameful and anxious, which can diminish their satisfaction with the colonoscopy. To decrease the shame and anxiety induced by exposing the buttocks of examinees wearing CSP during colonoscopy, we developed novel double pants (NDP). Examinees wearing NDP can undergo colonoscopy without taking off the inner pants. The smaller area of the buttock exposed by using NDP could minimize shame and anxiety during colonoscopy. Therefore we hypothesized that NDP could decrease shame and anxiety and increase satisfaction. We aimed to assess satisfaction, anxiety, and shame of patients wearing NDP compared with patients wearing CSP during colonoscopy through a prospective randomized single-center study. We also investigated the factors associated with patient satisfaction, anxiety, and shame during colonoscopy.

MATERIALS AND METHODS

Participants and study design

The study included examinees over 20 years old who agreed to undergo colonoscopy without sedation in the endoscopy unit of Severance Hospital, Yonsei University College of Medicine, Seoul, South Korea from January 2012 and July 2012. All of included people were the patients who visited the clinic for routine health check-up or for evaluating their mild gastrointestinal symptoms. Because we tried to minimize the selection bias of tertiary referral medical center, we excluded the patients who were referred by the physicians of primary or secondary




Figure 1 Consort diagram. ASA: American Society of Anesthesiologists.

medical center and needed more special care due to their objective medical problems. Before they signed at written informed consent, the nurse in outpatient clinic explained the purpose of the study. They got the information about not only the purpose of the study, but also the study design, the types of pants, the possible adverse events of procedure, and the contents of interview. Examinees under 20 years of age, those who had undergone colonoscopy within the past three years, those who were treated with an emergent colonoscopic procedure, those with a history of bowel resection surgery, inflammatory bowel disease (IBD), cancer, colostomy, ileostomy, psychotic disease including depression, anxiety disorder, or obsessive compulsive disorder, women who were pregnant or lactating, illiterate patients, foreigners, or examinees with American Society of Anesthesiologists (ASA) Scores of ≥ 3 were excluded. We used our preliminary survey data to perform a power calculation due to the lack of the prior published reports to guide this analysis. We calculated that a sample size of 80 participants was sufficient to detect an effect value of 0.5 (mean difference/common SD) at a significance level of 0.05% (two-sided) with 80% power and 20% drop-out rate. A total of 185 colonoscopy examinees were enrolled in this study between February 2012 and July 2012. Eight examinees were excluded because of colon cancer found during the colonoscopy, ten examinees were excluded because of psychologic disorder, three examinees were excluded because of a history of bowel resection, and four examinees were excluded because of ASA scores ≥ 3 , as shown in Figure 1. Finally a total of 160 consecutive examinees were randomly divided into NDP and CSP groups. Participants in this study were randomly assigned to a “CSP group” or a “NDP group” using a permuted four-block randomization method. Biostatistician made permuted four-block randomization table and calculated the numbers of participants. All random code was contained in a closed box. The nurse in outpatient clinic enrolled the examinees. Before colonoscopy, participants, outcome assessors, and care providers (endoscopists and nurses) were blinded to assignment to




Figure 2 Drawings of the conventional single pants and novel double pants. A: The conventional single pants were composed of a single layer of fabric; B: If the examinees wearing conventional single pants dropped the outer pants below hips, the total area of the hip is exposed; C: The novel double pants were composed of a double layer of fabric from the hip to the thigh and a single layer of fabric below the thigh; D: In the inner layer of the double pants, there is a hole for insertion of scope in the back. If the examinees wearing novel double pants dropped the outer pants below hips, the only buttock area is exposed through the hole (25 cm × 15 cm) for insertion of scope.

the allocation because the third party examiner allocated the enrolled subjects regarding to the prepared permuted-block randomization table. Written informed consent was provided by all participants in the study.

Pants

In CSP, there is no hole for insertion of the scope. Examinees remove the CSP to below the level of the buttocks and expose their buttocks area during colonoscopy, as shown in Figure 2A and B. NDP consist of single fabric only below the thigh, and doubled fabric from the hip to the thigh with a hole in the inner pants at the level of the buttocks. The hole is 25 cm wide and 15 cm long. Examinees wearing NDP can undergo colonoscopy without taking off the inner pants, as shown in Figure 2C and D. We supplied the 80 NDP (Bobo trading, Seoul, South Korea) to NDP group and 80 CSP (Seodaemun uniform, Seoul, South Korea) to CSP group. All patients got the pants in the hospital just before the procedure according to the randomized allocation by the nurse in outpatient clinic. The pants had been prepared with marking “A

type” (CSP) and “B type” (NDP).

Interview

In this study the only one nurse, as the third party examiner, interviewed the patients before and after the procedure to exclude interviewer’s influence on the answers in this study. Before colonoscopy, examinees underwent a one-on-one interview in a quiet, separated room with a third-party examiner in which they completed questionnaires identifying state and trait anxiety, marital status, education, and residence. After colonoscopy, examinees had a similar one-on-one interview to complete the questionnaires, which included the Group Health Association of America 9 (GHAA 9) scales and questions regarding pant-specific satisfaction, state anxiety, and shame. After colonoscopy, examinees’ pain during colonoscopy was assessed after the procedure and scored from 0 to 10 using a face pain scale (0-very happy, no pain, 2-hurts just a little bit, 4-hurts a little more, 6-hurts even more, 8-hurts a lot, 10-hurts as much as you can imagine; don’t have to be crying to feel this much pain)^[12].

Colonoscopy

Professional endoscopists who had performed more than 1000 colonoscopies performed all of the study colonoscopy procedures with a standard colonoscope (CF Q240L, CF Q240I, CF H260AI, CF Q260AI; Olympus Optical Co, Ltd, Tokyo, Japan). The indication for colonoscopy, ASA status, Ottawa quality scale, procedural time, number of polyps, number of examinees with polyps, method of polyp removal, rates of adverse events after polyp removal, rates of successful cecal approach, and gender difference between examinees and endoscopists in the NDP and CSP groups were investigated.

Outcome measurements

Satisfaction: The questionnaires measuring satisfaction included 14 questions regarding the examinees' colonoscopy experience. Nine of the questions regarding overall satisfaction were derived from a previously validated GHAA 9 satisfaction survey^[13]. These questions used a 5-point scale to grade satisfaction (1: poor, 2: fair, 3: good, 4: very good, 5: excellent). A score of more than 3 was considered a favorable response. Five questions were asked to elicit pants-specific satisfaction as related to the following: difficulty of defecation wearing the pants; difficulty in position change during the colonoscopy; worryness about the exposed buttock area; willingness to try the same pants for the next exam; and recommendation of the same pants to other people. The five pants-specific satisfaction questions used a 4-point scale to grade satisfaction (1, not at all; 2, somewhat; 3, moderately so; 4, very much so). Three of the five questions were negative questions about satisfaction and thus scored in reverse. Scores on pants-specific satisfaction ranged between 5 and 20.

Anxiety: The trait anxiety questionnaire has been shown to reflect the general disposition of patients or stable tendency for anxiety, while the state questionnaire reflects a patient's anxiety related to a particular set of circumstances^[9,14]. The trait and state questionnaires each consists of 20 statements, and all answers were graded using a 4-point scale (1, not at all; 2, somewhat; 3, moderately so; 4, very much so). Seven of the 20 questions from the STAI-Trait anxiety and 10 of the 20 questions from the STAI-State anxiety were negative questions and scored in reverse. The scores of the trait and state questionnaire range between 20 and 80, and higher scores reflected higher anxiety.

Shame: Among the experienced shame scale, 4 questions were used to measure body shame during colonoscopy^[15]. We designed another 2 questions to measure shame following the exposure of the body during position change for the colonoscopy and exposure of the buttock area while walking to the colonoscopy room. In total, 6 different questions were used to measure shame. All answers were graded using a 4-point scale (1, not at all; 2, somewhat; 3, moderately so; 4, very much so). Scores on the body shame questionnaire ranged between 6 and 24, and

higher scores reflected higher level of shame.

Ethical considerations

Informed consent was obtained from each patient included in the study. The study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki (6th revision, 2008) as reflected in a prior approval by the institution's human research committee. The study protocol was approved by the ethical committee of Yonsei University College of Medicine. The study protocol was also approved by clinicaltrial.gov (NCT 01524042).

Statistical analysis

All differences in CSP and NDP were examined using SPSS Statistics (version 18.0.0, IBM Corp., Armonk, NY, United States). Continuous variables were compared using Student's *t* test, while categorical data were analyzed using the χ^2 test (Fisher's exact test) between two groups. Univariate and multivariate linear regression analysis were used to assess independent predictive factors associated with satisfaction, state anxiety after colonoscopy, and shame. For multivariate analysis, variables with $P < 0.1$ by univariate analysis were included. All statistical tests were two-tailed and considered statistically significant with a P value < 0.05 .

RESULTS

Baseline characteristics and endoscopic characteristics of the study population

There was no significant difference in baseline characteristics including age, gender, marital status, education, and residence in the study population and in the NDP and CSP groups in Table 1. There were no significant differences in the endoscopic characteristics of study population including indication for colonoscopy, Ottawa quality scale, procedural time, number of polyps, number of examinees with polyps, method of polyp removal, rates of adverse events after polyp removal, rates of successful cecal approach, pain scale, and gender difference between examinees and endoscopists in the NDP and CSP groups, as shown in Table 2.

Outcomes of satisfaction, state anxiety, trait anxiety and shame

There was no significant difference in GHAA9 score (Likert scale, 0-5) between the NDP and CSP groups. There were high favorable response rates (FRRs) which were greater than 90% for waiting time, waiting on procedure day, personal manner of physician and support staff, technical skills, adequacy of explanation, overall rating of the visit, and willingness to have the procedure repeated by the same physician and at the same facility in Table 3. In terms of pants-specific satisfaction in Table 3, the CSP group worried more about exposing the buttock area during the colonoscopy than did the NDP group (3.3 ± 0.7 vs 2.9 ± 0.7 , $P < 0.001$). The NDP group was more willing to wear same pants when they undergo their

Table 1 Baseline characteristics of study population *n* (%)

	CSP (<i>n</i> = 80)	NDP (<i>n</i> = 80)	<i>P</i> value
Age (yr)	59.1 ± 11.4	59.4 ± 11.7	0.847
20 ≤ age < 40	3 (3.8)	5 (6.2)	
40 ≤ age < 60	38 (47.5)	43 (53.8)	
age ≥ 60	39 (48.7)	32 (40)	
Gender			0.210
Male	41 (51.3)	45 (56.3)	
Female	39 (48.7)	35 (43.7)	
Marital status			> 0.999
Married	75 (93.8)	76 (96.0)	
Single or divorced	5 (6.2)	4 (4.0)	
Education			0.320
Middle school	20 (25.0)	13 (16.2)	
High school	31 (38.8)	28 (35.0)	
University	22 (27.5)	32 (40.0)	
Graduate school	7 (8.7)	7 (8.8)	
Residence			0.609
Urban	73 (91.3)	70 (87.5)	
Rural	7 (8.7)	10 (12.5)	
ASA status			0.896
Class I	32 (40)	36 (45.0)	
Class II	48 (60)	44 (55.0)	
Class III	0 (0.0)	0 (0.0)	
Class IV	0 (0.0)	0 (0.0)	
Class V	0 (0.0)	0 (0.0)	

CSP: Conventional single pants; NDP: Novel double pants; ASA: American Society of Anesthesiologists.

next colonoscopy than the CSP group (3.3 ± 0.8 vs 2.1 ± 0.9 , $P < 0.001$). The NDP group was also more willing to recommend other people wear the same pants when they undergo their own colonoscopies than the CSP group (3.3 ± 0.7 vs 2.0 ± 1.0 , $P < 0.001$). A significantly lower shame score was estimated in the NDP group compared with CSP group (6.6 ± 1.5 vs 8.1 ± 3.2 , $P < 0.001$), which is shown in Table 3.

Predictive factors for satisfaction, state anxiety, and shame after colonoscopy

To investigate the predictive factors related to satisfaction, state anxiety, and shame after colonoscopy, univariate and multivariate regression analysis was performed (Table 4). In the multivariate analysis of pants-specific satisfaction, unmarried examinees had less pants-specific satisfaction than married examinees [B (SE) = -1.82 (0.61), $P = 0.004$], and the NDP group had higher pants-specific satisfaction than the CSP group [B (SE) = 2.72 (0.28), $P < 0.001$]. In multivariate analysis of state anxiety after the colonoscopy, female participants had higher state anxiety score after the procedure than males [B (SE) = 3.52 (1.14), $P = 0.002$]. Unmarried examinees had higher state anxiety score after colonoscopy than married examinees [B (SE) = 4.45 (2.22), $P = 0.047$]. Urban examinees had higher state anxiety score after colonoscopy than rural examinees [B (SE) = 4.68 (1.70), $P = 0.007$]. The NDP group had a tendency of lower state anxiety score after colonoscopy than the CSP group [B (SE) = -1.80 (1.04), $P = 0.086$]. In the multivariate analysis of shame, female examinees had higher shame score than male examinees [B (SE) = 1.25

Table 2 Endoscopic characteristics of study population *n* (%)

	CSP (<i>n</i> = 80)	NDP (<i>n</i> = 80)	<i>P</i> value
Indication of colonoscopy			0.260
Abnormality on other study	2 (2.5)	4 (5.0)	
Stool occult blood positive	9 (11.2)	13 (16.2)	
Screening	47 (58.8)	42 (52.5)	
Anemia	0.0 (0)	1.0 (1.2)	
Diarrhea	3(3.8)	4 (5.0)	
Surveillance after polyp removal	6 (7.5)	11 (13.8)	
Abdominal pain	13 (16.2)	5 (6.3)	
Ottawa Quality Scale	3.9 ± 2.5	4.3 ± 2.8	0.382
Procedural time (min)			
Insertion	10.7 ± 8.5	8.6 ± 5.5	0.059
Withdrawal	12.8 ± 6.2	11.8 ± 5.5	0.309
Number of polyp	1.1 ± 1.8	0.8 ± 1.5	0.266
Number of examinees with polyps	47 (58.8)	42 (52.5)	0.525
Method of polyp removal			0.521
Biopsy	25 (52.1)	22 (56.4)	
Snaring polypectomy	15 (31.3)	8 (20.5)	
Endoscopic mucosal resection	8 (16.6)	9 (23.1)	
Adverse events			
Bleeding	0 (0.0)	0 (0.0)	-
Perforation	0 (0.0)	0 (0.0)	-
Cecal approach	80 (100)	80 (100)	-
Pain scale ¹	3.5 ± 2.7	3.7 ± 2.4	0.719
Genders between examinee and endoscopist			0.525
Same gender	42 (52.5)	47 (58.8)	
Different gender	38 (47.5)	33 (41.2)	

Data are expressed as absolute numbers (percentage) or mean ± SD. ¹Examinees' pain during colonoscopy was assessed after the procedure and scored from 0 to 10 using a face pain scale (0-very happy, no pain, 2-hurts just a little bit, 4-hurts a little more, 6-hurts even more, 8-hurts a lot, 10-hurts as much as you can imagine; don't have to be crying to feel this much pain). CSP: Conventional single pants; NDP: Novel double pants.

(0.37), $P = 0.001$]. Unmarried examinees also had higher shame score than married examinees [B (SE) = 2.78 (0.81), $P = 0.001$]. The NDP group had lower shame score than the CSP group [B (SE) = -1.37 (0.37), $P = 0.001$].

DISCUSSION

In this study, we compared the satisfaction, anxiety and shame between NDP and CSP with prospective randomized control trial. The examinees in NDP group responded with higher pants-specific satisfaction, lower state anxiety after colonoscopy, and lower shame scores compared to those in CSP group. Thus, the NDP developed at our institution may contribute to increased satisfaction and decreased anxiety and shame after colonoscopy.

Although there have been a wide range of studies regarding satisfaction^[1,16-20] and anxiety^[9,21-23] during colonoscopy, our study was unique because it specifically addressed colonoscopic pants and investigated the differences in emotional change by the type of colonoscopic pants participants wore. Colonoscopic pants were designed considering the maximization of the efficiency of colonoscopies and hygiene aspects. Various types of colonoscopic pants have been used at different institutions. In some centers, examinees wear CSP or pants with

Table 3 Comparison of satisfaction rated by Group Health Association of America 9 survey by favorable response rate and likert scale, pants specific satisfaction, state anxiety and shame after colonoscopy between novel double pants and conventional single pants

	FRR, <i>n</i> (%)			Likert scale (mean ± SD)		
	CSP	NDP	<i>P</i> value	CSP	NDP	<i>P</i> value
GHAA9						
Appointment wait time	73 (91.3)	72 (90.0)	> 0.999	3.7 ± 0.7	3.6 ± 0.9	0.351
Waiting on procedure day	74 (92.5)	76 (95.0)	0.746	3.8 ± 0.8	3.9 ± 0.8	0.649
Personal manner of physician	78 (97.5)	79 (98.8)	> 0.999	4.4 ± 0.6	4.4 ± 0.6	0.712
Technical skills of physician	74 (92.5)	76 (95.0)	0.746	4.1 ± 0.8	4.1 ± 0.7	0.564
Personal manner of support staff	77 (96.3)	78 (97.5)	> 0.999	4.3 ± 0.7	4.0 ± 0.5	0.546
Adequacy of explanation of what was done	74 (92.5)	78 (97.5)	0.276	4.0 ± 0.6	4.0 ± 0.5	0.902
Overall rating of visit	75 (93.8)	74 (92.5)	> 0.999	3.9 ± 0.6	3.9 ± 0.6	0.806
Yes/No questions						
Would have procedure by same physician: Yes	78 (97.5)	77 (96.3)	> 0.999			
Would have procedure at same facility: Yes	79 (98.8)	79 (98.8)	> 0.999			
Pants specific satisfaction						
Difficulty in defecation				3.7 ± 0.5	3.8 ± 0.4	0.106
Difficulty in position change				3.7 ± 0.5	3.8 ± 0.3	0.175
Worriens about exposing buttock area in procedure				3.3 ± 0.7	2.9 ± 0.7	< 0.001
Will to wear same pants at next colonoscopy				2.1 ± 0.9	3.3 ± 0.8	< 0.001
Will to recommend to other people to wear same pants				2.0 ± 1.0	3.3 ± 0.7	< 0.001
Before colonoscopy						
Total scores of trait anxiety				39.2 ± 8.8	39.4 ± 8.2	0.904
Total scores of state anxiety				37.9 ± 8.2	39.9 ± 8.8	0.144
After colonoscopy						
Total scores of state anxiety				35.4 ± 6.9	33.0 ± 7.0	0.028
Total scores of shame				8.1 ± 3.2	6.6 ± 1.5	< 0.001
Change of state anxiety before and after the procedure				-2.4 ± 7.6	-6.9 ± 8.4	0.001

The scores on pants-specific satisfaction ranged between 5 and 20, and the scores of the trait and state anxiety questionnaire range between 20 and 80. The scores on the body shame questionnaire ranged between 6 and 24. A score of more than 3 of Likert scale was considered a FRR (Excellent: 5; Very Good: 4; Good: 3; Fair: 2; Poor: 1). Likert scale: Excellent: 5; Very Good: 4; Good: 3; Fair: 2; Poor: 1. Scores of pants specific satisfaction, Not at all: 1; A little: 2; Moderately: 3; Very much: 4. CSP: Conventional single pants; NDP: Novel double pants; GHAA9: Group Health Association of America 9; FRR: Favorable response rate.

Table 4 Univariate and multivariate analysis of pants specific satisfaction after colonoscopy

Predictive factors	Outcome (pants specific satisfaction)				Outcome (state anxiety after colonoscopy)				Outcome (shame after colonoscopy)			
	Univariate		Multivariate		Univariate		Multivariate		Univariate		Multivariate	
	B (SE)	<i>P</i> value	B (SE)	<i>P</i> value	B (SE)	<i>P</i> value	B (SE)	<i>P</i> value	B (SE)	<i>P</i> value	B (SE)	<i>P</i> value
Sex												
Male	Ref.				Ref.		Ref.		Ref.		Ref.	
Female	0.04 (0.36)	0.913			4.29 (1.06)	0.913	3.52 (1.14)	0.002	1.31 (0.40)	0.001	1.25 (0.37)	0.001
Age												
Age	-0.02 (0.01)	0.104			0.01 (0.04)	0.104			-0.01 (0.01)	0.928		
Marital status												
Married	Ref.		Ref.		Ref.		Ref.		Ref.		Ref.	
Unmarried	-1.6 (0.7)	0.033	-1.82 (0.61)	0.004	4.48 (2.40)	0.033	4.45 (2.22)	0.047	2.79 (0.87)	0.002	2.78 (0.81)	0.001
Education												
Middle school	Ref.				Ref.				Ref.			
High school	-0.04 (0.49)	0.931			-1.81 (1.53)	0.931			-0.01 (0.57)	0.998		
University	0.27 (0.50)	0.506			-2.64 (1.55)	0.506			0.22 (0.58)	0.697		
Graduate school	0.57 (0.73)	0.430			-2.50 (2.24)	0.430			0.15 (0.84)	0.853		
Type of pants												
CSP	Ref.		Ref.		Ref.		Ref.		Ref.		Ref.	
NDP	2.70 (0.29)	< 0.001	2.72 (0.28)	< 0.001	-2.43 (1.10)	< 0.001	-1.80 (1.04)	0.086	-1.47 (0.39)	< 0.001	-1.37 (0.37)	0.001
Residence												
Rural area	Ref.				Ref.		Ref.		Ref.			
Urban area	-0.46 (0.58)	0.430			4.2 (1.7)	0.430	4.68 (1.70)	0.007	0.93 (0.67)	0.165		
Genders between examinee and endoscopist												
Same gender	Ref.				Ref.		Ref.		Ref.			
Different gender	-0.36 (0.36)	0.308			3.90 (1.07)	< 0.001	1.45 (1.15)	0.209	0.65 (0.41)	0.115		

CSP: Conventional single pants; NDP: Novel double pants.

a hole in the buttock area with a movable flap, which can potentially expose buttock area during walking. In other centers, examinees wear single pants with hole in the buttocks and outer gown to hide the hole, which is likely inferior to the NDP we describe in terms of repair and maintenance expenses.

In our study there was no significant difference in GHAA 9 patient satisfaction between NDP and CSP groups as the contents of GHAA 9 were not closely connected with satisfaction regarding colonoscopic pants. However, in the survey of pants-specific satisfaction, the NDP group showed increased satisfaction than the CSP group in terms of their willingness to wear the same pants in the next time and to recommend the same pants to other people. High patient satisfaction during colonoscopy may result in a higher rate of compliance with screening and clinical surveillance programs^[1]. The American Society for Gastrointestinal Endoscopy and the American College of Gastroenterology recommends assessment of patient satisfaction during colonoscopy to evaluate the quality of colonoscopy^[5,6].

The sensitivity related to the psychological aspects including anxiety and shame might partly depends on the different characteristics including age, gender, behavior, social environment and sensitivity of the society. As example lower anxiety scores was reported to be associated with older age, male sex, lower income, experience of previous colonoscopy and lower education^[9]. During gastrointestinal endoscopic procedures, female examinees have been reported to have higher state and trait anxiety levels than male examinees^[10,22]. In our study, female, unmarried, urban examinees and CSP group had more state anxiety after colonoscopy than that of their male, married, rural examinees and NDP group counterparts. The differences of shame and state of anxiety after colonoscopy between the two groups were moderate in most of the items. But, the difference in change of state of anxiety before and after the procedure might be clinically significant. Because all participants are healthy persons in psychological aspect at baseline, even moderate change of anxiety level after colonoscopy could result in decreased satisfaction of the procedure and reduced compliance to next examinations in clinical practice.

There were some limitations to our study. Endoscopists could not be completely blinded to the types of colonoscopic pants worn because they ultimately saw which pants they were wearing during the colonoscopic procedure. However, the outcome assessors were blinded during the total period of study. In addition, all the endoscopists who performed the colonoscopy were excluded from the outcome assessors. And there was no questionnaire for endoscopists in this study. Examinees with cancer or IBD were suspected to have higher anxiety levels during colonoscopy, and thus were excluded. Therefore understanding the satisfaction, anxiety, and satisfaction of these particular examinees would require further investigation. Most of examinees in this study were over 40 years old and more than half of examinees underwent colonoscopy for screening purposes; the num-

ber of young unmarried examinees was very small in this study, and the level of anxiety, shame and satisfaction in this group may be significantly different. For a more ideal comparison of anxiety, shame and satisfaction during colonoscopy between the NDP and CSP group, it would be better for one examinee to wear single pants and double pants during colonoscopy, but this was not logistically feasible. To compensate for this limitation in state anxiety we measured the change in state anxiety before and after colonoscopy in the same examinees.

In conclusion, the examinees in the NDP group had higher pants-specific satisfaction and lower state anxiety and lower shame after colonoscopy compared to CSP group. Therefore NDP could help to increase satisfaction and decrease anxiety and shame after colonoscopy. Future studies should continue to investigate factors for anxiety, shame and satisfaction.

ACKNOWLEDGMENTS

We thank Mr. Dong-Su Jang for help in preparing the specimens and figures.

COMMENTS

Background

Exposing buttocks during colonoscopy can make examinees feel unsatisfied, anxious, and shameful. To increase satisfaction and diminish anxiety and shame during colonoscopy, the authors developed novel double pants (NDP) which consist of doubled fabrics with an inner hole.

Research frontiers

Pants-specific satisfaction scores regarding willingness to repeat colonoscopy using same pants and recommendation of same pants to other people were significantly higher in NDP than conventional single pants (CSP) groups. State anxiety and shame after colonoscopy was lower in NDP group compared with CSP group.

Innovations and breakthroughs

Although there have been a wide range of studies regarding satisfaction and anxiety during colonoscopy, this study was unique because it specifically addressed colonoscopic pants. The authors developed NDP, which consist of single fabric only below the thigh, and doubled fabric from the hip to the thigh with a hole in the inner pants at the level of the buttocks. Examinees wearing NDP can undergo colonoscopy without taking off the inner pants.

Applications

Through these findings, the NDP developed at our institution may contribute to increased satisfaction and decrease anxiety and shame after colonoscopy.

Terminology

NDP are single fabric only below the thigh, and doubled fabric from the hip to the thigh with a hole in the inner pants at the level of the buttocks. The hole is 25 cm wide and 15 cm long.

Peer review

Satisfaction studies are important in relationship with the compliance of colorectal cancer screening programs and less so in the group of patients with specific suspicion of diseases affecting the anus, rectum or colon. This complex study is well designed and analyzed. The topic of the manuscript is interesting and the work performed is ambitious.

REFERENCES

- 1 Seip B, Bretthauer M, Dahler S, Friestad J, Huppertz-Hauss G, Høie O, Kittang E, Nyhus S, Pallenschat J, Sandvei P, Stallemo A, Svendsen MV, Hoff G. Patient satisfaction with on-demand sedation for outpatient colonoscopy. *Endoscopy* 2010; 42: 639-646 [PMID: 20669075 DOI: 10.1055/s-0030-1255612]

- 2 **Porostocky P**, Chiba N, Colacino P, Sadowski D, Singh H. A survey of sedation practices for colonoscopy in Canada. *Can J Gastroenterol* 2011; **25**: 255-260 [PMID: 21647459]
- 3 **Cohen LB**, Wechsler JS, Gaetano JN, Benson AA, Miller KM, Durkalski V, Aisenberg J. Endoscopic sedation in the United States: results from a nationwide survey. *Am J Gastroenterol* 2006; **101**: 967-974 [PMID: 16573781 DOI: 10.1111/j.1572-0241.2006.00500.x]
- 4 **Froehlich F**, Harris JK, Wietlisbach V, Burnand B, Vader JP, Gonvers JJ. Current sedation and monitoring practice for colonoscopy: an International Observational Study (EPAGE). *Endoscopy* 2006; **38**: 461-469 [PMID: 16767580 DOI: 10.1055/s-2006-925368]
- 5 **Faigel DO**, Pike IM, Baron TH, Chak A, Cohen J, Deal SE, Hoffman B, Jacobson BC, Mergener K, Petersen BT, Petrini JL, Rex DK, Safdi MA. Quality indicators for gastrointestinal endoscopic procedures: an introduction. *Gastrointest Endosc* 2006; **63**: S3-S9 [PMID: 16564906 DOI: 10.1016/j.gie.2006.02.017]
- 6 Quality improvement of gastrointestinal endoscopy: guidelines for clinical application. From the ASGE. American Society for Gastrointestinal Endoscopy. *Gastrointest Endosc* 1999; **49**: 842-844 [PMID: 10343248]
- 7 **Parker D**. Human responses to colonoscopy. *Gastroenterol Nurs* 1992; **15**: 107-109 [PMID: 1472554]
- 8 **Brandt LJ**. Patients' attitudes and apprehensions about endoscopy: how to calm troubled waters. *Am J Gastroenterol* 2001; **96**: 280-284 [PMID: 11232665 DOI: 10.1111/j.1572-0241.2001.03508.x]
- 9 **Jones MP**, Ebert CC, Sloan T, Spanier J, Bansal A, Howden CW, Vanagunas AD. Patient anxiety and elective gastrointestinal endoscopy. *J Clin Gastroenterol* 2004; **38**: 35-40 [PMID: 14679325]
- 10 **Trevisani L**, Sartori S, Putinati S, Gaudenzi P, Chiamenti CM, Gilli G, Grassi L, Abbasciano V. [Assessment of anxiety levels in patients during diagnostic endoscopy]. *Recenti Prog Med* 2002; **93**: 240-244 [PMID: 11989128]
- 11 **Tønnesen H**, Puggaard L, Braagaard J, Ovesen H, Rasmussen V, Rosenberg J. Stress response to endoscopy. *Scand J Gastroenterol* 1999; **34**: 629-631 [PMID: 10440615]
- 12 **Warden V**, Hurley AC, Volicer L. Development and psychometric evaluation of the Pain Assessment in Advanced Dementia (PAINAD) scale. *J Am Med Dir Assoc* 2003; **4**: 9-15 [PMID: 12807591 DOI: 10.1097/01.JAM.0000043422.31640.F7]
- 13 **Del Río AS**, Baudet JS, Fernández OA, Morales I, Socas Mdel R. Evaluation of patient satisfaction in gastrointestinal endoscopy. *Eur J Gastroenterol Hepatol* 2007; **19**: 896-900 [PMID: 17873615 DOI: 10.1097/MEG.0b013e3281532bae]
- 14 **Spielberger CD**, Gorsuch RL, Lushene RE. STAI manual for the state-trait anxiety inventory ("self-evaluation questionnaire"). Palo Alto: Consulting Psychologists Press, Inc., 1970: 1-23
- 15 **Andrews B**, Qian M, Valentine JD. Predicting depressive symptoms with a new measure of shame: The Experience of Shame Scale. *Br J Clin Psychol* 2002; **41**: 29-42 [PMID: 11931676]
- 16 **Nijjar UK**, Edwards JA, Short MW. Patient satisfaction with family physician colonoscopists. *J Am Board Fam Med* 2011; **24**: 51-56 [PMID: 21209344 DOI: 10.3122/jabfm.2011.01.100112]
- 17 **Lin OS**, Kozarek RA, Arai A, Gluck M, Jiranek GC, Kowdley KV, McCormick SE, Schembre DB, Soon MS, Dominitz JA. The effect of periodic monitoring and feedback on screening colonoscopy withdrawal times, polyp detection rates, and patient satisfaction scores. *Gastrointest Endosc* 2010; **71**: 1253-1259 [PMID: 20598251 DOI: 10.1016/j.gie.2010.01.017]
- 18 **Ko HH**, Zhang H, Telford JJ, Enns R. Factors influencing patient satisfaction when undergoing endoscopic procedures. *Gastrointest Endosc* 2009; **69**: 883-891, quiz 891.e1 [PMID: 19152911 DOI: 10.1016/j.gie.2008.06.024]
- 19 **Chartier L**, Arthurs E, Sewitch MJ. Patient satisfaction with colonoscopy: a literature review and pilot study. *Can J Gastroenterol* 2009; **23**: 203-209 [PMID: 19319384]
- 20 **Bytzer P**, Lindeberg B. Impact of an information video before colonoscopy on patient satisfaction and anxiety - a randomized trial. *Endoscopy* 2007; **39**: 710-714 [PMID: 17661246 DOI: 10.1055/s-2007-966718]
- 21 **Ersöz F**, Toros AB, Aydoğan G, Bektaş H, Özcan O, Arikan S. Assessment of anxiety levels in patients during elective upper gastrointestinal endoscopy and colonoscopy. *Turk J Gastroenterol* 2010; **21**: 29-33 [PMID: 20533109]
- 22 **Mitsonis C**, Dimopoulos N, Zavrou M, Psarra V, Giofkos C, Fiorakis C, Dimitriadis A, Valavanis D, Voursora E, Zervas I, Papavassiliou E. Panic Attack during Elective Gastrointestinal Endoscopy. *Gastroenterol Res Pract* 2011; **2011**: 162574 [PMID: 22007196 DOI: 10.1155/2011/162574]
- 23 **Ovayolu N**, Ucan O, Pehlivan S, Pehlivan Y, Buyukhatipoglu H, Savas MC, Gulsen MT. Listening to Turkish classical music decreases patients' anxiety, pain, dissatisfaction and the dose of sedative and analgesic drugs during colonoscopy: a prospective randomized controlled trial. *World J Gastroenterol* 2006; **12**: 7532-7536 [PMID: 17167846]

P- Reviewers Blom J, Bordas JM, Bujanda L **S- Editor** Zhai HH
L- Editor A **E- Editor** Ma S





百世登

Baishideng®

Published by **Baishideng Publishing Group Co., Limited**

Flat C, 23/F., Lucky Plaza,

315-321 Lockhart Road, Wan Chai, Hong Kong, China

Fax: +852-65557188

Telephone: +852-31779906

E-mail: bpgoffice@wjgnet.com

<http://www.wjgnet.com>



ISSN 1007-9327



9 771007 932045